

PROPERTY Granite Mountain
 GRID _____

CANEX AERIAL EXPLORATION LTD.
 DIAMOND DRILL LOG

012945

HOLE No. RD 11-3
 SHEET 1 OF 8

LOCATION Line 00100, A20+00N BEARING _____ LATITUDE _____ CORE SIZE B.G. wireline LOGGED BY D. Howard
 DATE COLLARED July 5, 1967 LENGTH 426' DEPARTURE _____ SCALE OF LOG 1"=10' DATE July 14, 1967
 DATE COMPLETED July 8, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE STRUCTURE			WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
		0	Casing to 12 feet. No core								
White fine grain porphyritic rhyolite Phenocrysts are quartz No alteration	None	10	Pale yellow to light brown limonite along all joint planes. Questionable cubic boxwork on some of the joint planes. Intensely jointed; Very blocky 5-30°, 7-60° 6-80° 2-20°								
Same as above	None	20	Same as above Joints 2-30°, 9-60°, 4-70°, 8-80°, 4-20°								
Same as above	None	30	Same as above except some disseminated limonite is present. 1-15°, 5-60°, 14-70°, 4-80°, 2-80°								
Same as above	None	40	Same as above Many joints with the same angles to the core axis as above								
Same as above	None	50	Same as above except a few of the joints do not contain limonite								

PROPERTY Granite Mountain
 GRID _____

CANEX AERIAL EXPLORATION LTD.
DIAMOND DRILL LOG

HOLE No. 00H-3
 SHEET 2 OF 6

LOCATION Line 00100 A 20100N BEARING _____ LATITUDE _____ CORE SIZE 5.0 wireline LOGGED BY D. Howard
 DATE COLLARED July 5, 1967 LENGTH 426' DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 11, 1967
 DATE COMPLETED July 8, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE			WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
White fine grain porphyritic rhyolite. Quartz phenocrysts No alteration	None	60	Pale yellow to light brown limonite along joint plane. Several joint with dark brown limonite Intense jointed and blocky Several plates of MoS ₂ Joint angles - 20°, 30°, 60°, 70°, 80°								Tr-MoS ₂
Same as above	None	70	Same as above with the following exception Ferromolybdate assoc. with qtz on 70° joint 1/16" vein of FeS ₂ on 60° joint No visible MoS ₂								Tr Mo
Same as above	None	80	Same as above except for mineralization								0
Same as above	None	90	Same as above								0
Same as above	None	100	Same as above with the exception of one joint coated with FeS ₂ (164)								0
Same as above	None	110	Same as above except for two MoS ₂ (164) joints at 113-114. Spotty coating. This section contained very little limonite, compared to rest of section. Some ferromoly along joints not containing MoS ₂ (113-14)								.01 MoS ₂

PROPERTY Granite Mountain
GRID _____

CANEX AERIAL EXPLORATION LTD.
DIAMOND DRILL LOG

HOLE No. COH-3
SHEET 3 OF 8

LOCATION Line COH-3 2012001 BEARING _____ LATITUDE _____ CORE SIZE P.B. wireline LOGGED BY D. Howard
DATE COLLARED July 5, 1967 LENGTH 426' DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 11, 1967
DATE COMPLETED July 8, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE			WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
White fine grain porphyritic rhyolite Quartz phenocrysts No alteration		120	Pale yellow to dark brown limonite along most joints. Intensely jointed and blocky. Joint angles 20°, 30°, 60°, 70°, 80°. 2 hairline MoS ₂ coatings on 80° joints. 1- FeS ₂ (1/8") coated joint. Some finely disseminated MoS ₂ between 127-128, enough to give the rhyolite a gray appearance. Mineralized sect. contains little limonite.								.02 MoS ₂
Same as above		130	Same limonite coated joints and intensity of jointing as above. 4-joints (60°+20°) containing finely disseminated coating of bornite coated pyrite. Bornite coated FeS ₂ grains are all approx. 1/64" in diameter.								.02 Cu
Same as above		140	Hairline MoS ₂ vein with some disseminated bornite stained FeS ₂ . Same limonite coated joints and intensity of jointing as above.								Tr Cu MoS ₂
Same as above		150	Same limonite coated joints and intensity of jointing as above. 1- Hairline MoS ₂ along on 30° joint. 1- 1" zone of bornite (1) coated pyrite - very fine grained. - associated with a very siliceous zone.								Tr Cu MoS ₂
Same as above		160	Same limonite coats and intensity of jointing as above. 1- Hairline bornite coated 80° joint (disseminated)								Tr-Cu
Same as above		170	Same limonite coated and intensity of jointing as above. 2- Hairline disseminated MoS ₂ coatings along 70° joints - very fine grained.								Tr - MoS ₂

PROPERTY Granite Mountain
 GRID _____

CANEX AERIAL EXPLORATION LTD.
 DIAMOND DRILL LOG

HOLE No. AB-8-9
 SHEET 4 OF 6

LOCATION Line 00+00 AB-8-9 BEARING _____ LATITUDE _____ CORE SIZE 8.9 wireline LOGGED BY D. Howard
 DATE COLLARED July 5, 1967 LENGTH 426' DEPARTURE _____ SCALE OF LOG 1"=10' DATE July 12, 1967
 DATE COMPLETED July 8, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE			WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
White fine grain porphyritic rhyolite. Quartz phenocrysts No alteration	None	180	Sale yellow to dark brown limonite along most joints. Intensely jointed and blocky. Joint 20, 30, 60, 70, 80, 101.5. Pyrite + Chalcocite vein (24" - 20") 182 - Hairline MoS ₂ (disseminated on 80° joint plane) First occurrence of chalcocite 2 - disseminated bornite coated pyrite veins along 80° joint 1 disseminated MoS ₂ vein along 70° joint								Tr - Cu .01 MoS ₂
Same as above	None	190	Same limonite coated joints and intensity of jointing 1 - Hairline MoS ₂ vein along 80° joint (disseminated) 1 - Hairline bornite coated pyrite vein along 60° joint 1 - disseminated pyrite vein along 60° joint								Tr MoS ₂ Cu
Same as above	None	200	Same limonite coated joints and intensity of jointing as above 2 - Hairline bornite coated pyrite veins along 80° joint (disseminated) 1 - Hairline bornite coated chalcocite (finely disseminated)								Tr - Cu
Same as above	None	210	Same limonite coated joints and intensity of jointing as above Extremely blocky Tr of MoS ₂ along 70° joint - assoc. with hairline grain Tr of bornite coated pyrite on 5° joint								Tr MoS ₂ + Cu
Same as above	None	220	Same limonite coated joints but jointing is less intense than above finely disseminated pyrite with Tr bornite coatings Approx 2' section missing 223 - 225 Tr of bornite + chalcocite along 60° joint Hairline MoS ₂ along 80° joint								Tr MoS ₂ + Cu
Same as above (Box 11) Alteration of QM indicates that the rhyolite is intrusive into the QM. Fine to med grain granitic gray to tan porphyritic 8.4. monzonite. Intense K-feldspar + chlorite alteration	None	230	Same limonite coated joints and intensity of jointing as above Very fine widely disseminated MoS ₂ , bornite coated pyrite Some on 70° joint surfaces Very irregular contact - Definitely intrusive but asc relationships not clear. Disseminated MoS ₂ , FeS ₂ - vein + finely								Tr MoS ₂

LOCATION Loc 00110 A 20+000' BEARING _____ LATITUDE _____ CORE SIZE B. Swivel LOGGED BY D. Howard
DATE COLLARED July 5, 1967 LENGTH 476' DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 12, 1967
DATE COMPLETED July 8, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	ROCK TYPE ALTERATION	GRAPHIC LOG FOOTAGE STRUCTURE	MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
					WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
<i>Fine to med. gr. greenish gray gte monzonite intermixed with rhyolite. Contact probably brecciated on a macro scale. Q.M. intensely altered. Matrix all altered to chlorite. Plag. to kersin, kersin shows little alteration. Kersin phenocrysts are white to gray</i>		240	<i>Some of the joints are limonited coat. Jointing is very intense and the ground is very blocky. Joint angles - 50, 60, 20, 70 Very fine unevenly disseminated pyrite, bornite coated pyrite & chalcopyrite and chalcopyrite, (?) moly Some along a few joints</i>								.01 Cu Tr. MoS ₂
<i>Fine to medium grain med gray biotite gte monzonite. Biotite is fine grained and clotted (secondary?) Plag is intensely alt. to kersin Some of the plag has a faint greenish cast. (Ferrocite?)</i>		250	<i>Same as above</i>								.01 Cu Tr. MoS ₂
<i>Same as above with the following exception: At 265 all biotite has disappeared, replaced by chlorite. Alteration is more intense. Kersin is gradually becoming more pink toward 270</i>		260	<i>Same as above</i>								.01 Cu Tr. MoS ₂
<i>Fine to med. gr. dark greenish gray, slightly porphyritic gte monzonite. Very intense chlorite & kersin alteration. 4" dark gray fine grain andesite dike (60°)</i>		270	<i>Approximately 4 feet of last core lost Very blocky. Only minor limonite staining on joints Very spotty disseminated chalcopyrite & pyrite</i>								Tr. Cu
<i>Same as above except it contains short sections of less altered Q.M. In these the biotite is only slightly altered</i>		280	<i>Extremely blocky 1/2 x 1/8 plate of bornite coated Chalco Very spotty disseminated chalcopyrite & pyrite Several spotty bornite coated chalcopyrite veins along 30° joints (Hornblende)</i>								.01 Cu Tr. MoS ₂
<i>Same as above except color is lighter due to slight increase in gte content Narrow rhyolite dike (6") at 290</i>		290	<i>Very blocky. Very little limonite on joint surfaces. Trace amount of epidote assoc. with some joints Some disseminated FeS₂ Several near vertical joints with a thin spotty coating of bornite coated chalcopyrite (chalcocite?)</i>								.01 Cu Tr. MoS ₂

PROPERTY Granite Mountain
GRID _____

CANEX AERIAL EXPLORATION LTD.
DIAMOND DRILL LOG

HOLE No. DDH-3
SHEET 6 OF 6

LOCATION Line 00400 380+00N BEARING _____ LATITUDE _____ CORE SIZE B. Q. wireline LOGGED BY D. Howard
DATE COLLARED July 5, 1967 LENGTH 426' DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 13, 1966
DATE COMPLETED July 8, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE			STRUCTURE	WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag
Fine to medium grain greenish gray, slightly porphyritic quartz monzonite becoming slightly lighter in color at 308. Pink oplitic section at 310 - not a slide. Intense chlorite and kaolin alteration. Light pink K-spar not altered.	300	300	Extremely blocky; joint angles 0, 30, 60, 90, 70, 20 Tr. of Epidote on joint planes 303-304 Fresh disseminated MoS ₂ , CuFeS ₂ + FeS ₂ Most of section is not mineralized 309-310 Very finely disseminated bornite coated pyrite								Tr. MoS ₂ .02 Cu
Fine to med. gr. light greenish gray slightly porphyritic gte. monzonite, i.e. like the section from 308-310. Alteration same as above.											
See K-spar envelope (sample taken)	320	310	Extremely blocky - joint angles - no change Pyrite with chloropyrite on MoS ₂ along several vertical fractures Pyrite on 50° joints Epidote assoc. with joints carrying pyrite secondary 1/2 vein of chlorite (?) coated chloropyrite with small K-spar zone								.01 Cu
Fine to med. gr. greenish gray slightly porphyritic gte. monzonite. Same as section between 300-308. Change from lighter at 321. Change in color is gradational.											
Same as above	330	320	Several high-angle joints with disseminated pyrite, bornite coated pyrite + chloropyrite Vertical joints containing Tr. CuFeS ₂ chloropyrite shales coated with chlorite (?) Pyrite rich zone, Tr. of disseminated MoS ₂								Tr. MoS ₂ .02 Cu
Same as above											
Same as above	340	330	1/2-20 pyrite vein with (?) chlorite coatings Disseminated bornite with Tr. MoS ₂ Near vertical joint with disseminated FeS ₂ + Tr. chal. Epidote assoc. with most joints Extremely blocky								Tr. Cu Tr. MoS ₂
Same as above											
Same as above	350	340	Thinly disseminated FeS ₂ throughout section Tr. Epidote along most joints Tr. CuFeS ₂ + FeS ₂ on 50° joint Section extremely blocky Tr. bornite stained FeS ₂								Tr. Cu

FORM NO. 110

PROPERTY Granite Mountain
GRID _____

CANEX AERIAL EXPLORATION LTD.
DIAMOND DRILL LOG

HOLE No. DDH-3
SHEET 7 OF 8

LOCATION Line 02+00 to 20+00 BEARING _____ LATITUDE _____ CORE SIZE B.O. wingline LOGGED BY D. Howard
DATE COLLARED July 5, 1967 LENGTH 426' DEPARTURE _____ SCALE OF LOG 1"=10' DATE July 13, 1967
DATE COMPLETED July 3, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	ROCK TYPE ALTERATION	FOOTAGE	STRUCTURE	MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
						WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
360-362 same as previous interval 362 - Gradational zone - To medium to coarse grain dark bluish green gray porphyritic qtz monzonite to granodiorite. Bluish gray. Sharp reduction in quartz. Biotite not altered. Blue clay probably indicates sericite (Sample)	Sericite	360		Pyrite coated pyrite - Very thin Pyrite zones with sec. k-spar envelope with Tr. MoS ₂ + Tr. CuFeS ₂ Pyrite slick on 20° joint Magnetite + pyrite vein (20°) Sharp increase in pyrite - Barite coated pyrite vein (0°) 60° pyrite slick Very blocky								.01 Cu Tr. MoS ₂
Same as above (362-370)	Sericite	370		Very blocky. Some Epidote 4 foot zone of numerous 50, 60 & 20° joints coated with thin slicks of pyrite. Tr. MoS ₂ with FeS ₂								Tr Cu Tr. MoS ₂
Same as above	Sericite	380		Extremely blocky and broken. At least 5 foot missing from section Pyrite + chalcocite coated chalcocypite on joint surfaces. Ground core prevented determining orientation.								.01 Cu
Same as above	Sericite	390		Tr. of CuFeS ₂ with FeS ₂ Fault zone is 1' of sand Several pyrite zones - along 60 & 20° joints A little disseminated FeS ₂ throughout section with some chalcocypite Very blocky								Tr Cu
Same as above	Sericite	400		Numerous joints with thin slicks of FeS ₂ and Tr. of CuFeS ₂ . No one joint set is predominant								Tr Cu
Same as above	Sericite	410		Same as above. All joint are the same age ie No offset relationships. This applies to the entire hole								Tr Cu

HOLE No. 004-3
SHEET 8 OF 8

FORM NO. 110

PROPERTY Granite Mountain
 GRID

CANEX AERIAL EXPLORATION LTD.
 DIAMOND DRILL LOG

HOLE No. DDH-6
 SHEET 1 OF 5

LOCATION Line 7-50 E A 12+00 N BEARING LATITUDE CORE SIZE B. & wire line LOGGED BY D. Howard
 DATE COLLARED July 14, 1967 LENGTH 383 DEPARTURE SCALE OF LOG 1" = 10' DATE July 23, 1967
 DATE COMPLETED July 16, 1967 DIP 00° ELEVATION REMARKS

ROCK TYPES AND ALTERATION	ROCK TYPE ALTERATION	GRAPHIC LOG FOOTAGE STRUCTURE	MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
					WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
Cased to 91 feet with no core recovery. Partially overburden and partially intensely altered bed rock.		0	Note interval change at 90 feet.								
Fine to medium grain light gray to green porphyritic gte monzonite or granite. Feldspars are so intensely altered and weathered that identification is impossible Mafics - biotite and/or chlorite		90	Intense limonite staining along all joints & fractures and some disseminated. Trace amounts of disseminated FeS ₂ . Extremely pumy and broken due to intense weathering								
Same as above		100	Same as above								
Same as above		110	Same as above								
Same as above		120	Same as above								
Same as above		130	Same as above								

LOCATION Line 7405 AIRCON BEARING _____ LATITUDE _____ CORE SIZE BQ. wireline LOGGED BY D. Howard
 DATE COLLARED July 14, 1967 LENGTH 383 DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 22, 1967
 DATE COMPLETED July 16, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	ROCK TYPE ALTERATION	GRAPHIC LOG FOOTAGE	STRUCTURE	MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
						WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
Fine to medium grain light gray to green porphyritic biotite gte monzonite or granite. Feldspars are so intensely altered and weathered that identification is impossible		140		Intense limonite staining along all joints, fractures and disseminated. Tr amounts of disseminated FeS ₂ Extremely punky and broken due to intense weathering								
Same as above		150		Same as above								
Same as above except there is no more chlorite alteration. Rock now has a vague banded appearance.		160		Extremely punky and broken. Very little limonite staining and no pyrite								
Same as above		170		Same as above with more limonite								
Fine to medium grain greenish gray slightly porphyritic gte. monzonite. Intense alteration still make identification questionable. All feldspars have a greenish cast which could indicate sericite alteration. Sample at 188		180		Same as above								
Same as above		190		Same as above								

PROPERTY Garnet Mountain

GRID _____

CANEX AERIAL EXPLORATION LTD.

DIAMOND DRILL LOG

HOLE No. DDH-6SHEET 3 OF 5
 LOCATION Line 7505 A 184000N BEARING _____
 DATE COLLARED July 14, 1967 LENGTH 383
 DATE COMPLETED July 16, 1967 DIP 90°

 LATITUDE _____
 DEPARTURE _____
 ELEVATION _____

 CORE SIZE B. Q. wireline
 SCALE OF LOG 1"=10'
 REMARKS _____

 LOGGED BY D. Howard
 DATE July 29, 1967

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE			WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
Fine to medium grain, greenish gray slightly porphyritic Ote Monzonite(?) Feldspars have a greenish cast, but biotite is not altered - may be secondary. Alteration very intense.	Karl + Co. Inc. Hole 7505 A	200	Both limonite and pyrite occur along most joints, fractures. Section intensely broken due to weathering. Limonite - yellow brown. Some epidote present on a few joint planes.								
Same as above.		210	Same as above.								
Same as above although weathering is not quite as intense. Sample at 229		220	Same as above.								
Same as above.		230	Same as above.								
Same as above.		240	Same as above. 0.1 - 0.2 disseminated FeS ₂								
Same as above.		250	Same as above.								

PROPERTY Granite Mountain
GRID _____

CANEX AERIAL EXPLORATION LTD.
DIAMOND DRILL LOG

HOLE No. DDH-6
SHEET 4 OF 5

LOCATION Line 7450E A15100N BEARING _____ LATITUDE _____ CORE SIZE 8.0 wireline LOGGED BY D. Howard
DATE COLLARED July 14, 1967 LENGTH 383 DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 23, 1967
DATE COMPLETED July 16, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	GRAPHIC LOG		MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
	ROCK TYPE ALTERATION	FOOTAGE			WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MnS ₂	Oz/Ton Ag	EST. GRADE
Fine to medium grain greenish gray slightly porphyritic gte monzonite Feldspars have less of a given cast than before and the biotite is slightly altered. Weathering effects are less severe Rock has vague band appearance	Granite	260	Both limonite and pyrite are present along most joints and fractures. Disseminated FeS ₂ is also present in varying amounts. Some sections contain up to 1% Section intensely shattered								
Same as above	Granite	270	Same as above								
Same as above	Granite	280	Same as above								
Same as above Sample at 300'	Granite	290	Same as above. 7 feet of core is missing from 290-297 297-300 Relatively massive and not broken 1/2 block of CuFeS ₂ Several narrow FeS ₂ (30-40')								.01 Cu
Same as above	Granite	300	Horine CuFeS ₂ (60') Disseminated FeS ₂ throughout 4xline x 30 FeS ₂ Massive section 4xline x 20 FeS ₂ 1/2 x 80 FeS ₂ with Fe of CuFeS ₂ coated with chalcocite (?) 2-4 x 30 FeS ₂ 2" white aplite dike containing disem. FeS ₂ 1/2 x 20 x 10 FeS ₂								.01 Cu
Same as above	Granite	310	Most joints barren except for a few coated with limonite 1/2 x 20 FeS ₂ 1/2 x 10 CuFeS ₂ Very massive section Finely disseminated FeS ₂ throughout section								.02 Cu

PROPERTY Granite Mountain
 GRID _____

CANEX AERIAL EXPLORATION LTD.
 DIAMOND DRILL LOG

HOLE No. DDH-6
 SHEET 5 OF 5

LOCATION Line THESE 412-300N BEARING _____ LATITUDE _____ CORE SIZE B.O. winding LOGGED BY A. Howard
 DATE COLLARED July 14, 1967 LENGTH 383 DEPARTURE _____ SCALE OF LOG 1" = 10' DATE July 26, 1967
 DATE COMPLETED July 16, 1967 DIP 90° ELEVATION _____ REMARKS _____

ROCK TYPES AND ALTERATION	ROCK TYPE ALTERATION	FOOTAGE	STRUCTURE	MINERALIZATION AND STRUCTURES	SPECIFIC GRAVITY	REC. CORE		ASSAY RESULTS				
						WT. IN GRAMS %	SAMPLE No.	% Cu	% Cu	% MoS ₂	Oz/Ton Ag	EST. GRADE
Fine to medium grain, greenish gray slightly porphyritic etc. Monzonite. Some of the feldspars have a slight greenish cast. Very little pink Ksp. Some of the biotite is slightly altered to chlorite.		320		$\frac{1}{32}$ X 20° FeS ₂ , CuFeS ₂ + biotite coated CuFeS ₂ Disseminated FeS ₂ throughout section with trace amounts of CuFeS ₂ and magnetite 1/8 x 6 magnetite Very broken section								.01 Cu
Same as above		330		Disseminated FeS ₂ throughout limonite only on joints Very broken section.								Tr Cu
6" white aplite dike containing multiple parallel veins of magnetite + FeS ₂				Intense chlorite alteration from 338-340								
Same as above with intense chlorite alteration with some epidote May be a different intrusion although the compositions are similar.		340		Same as above								Tr Cu
Same as above		350		Same as above								Tr. MoS
				Trace amounts of MoS ₂ along vert. fault.								
Same as above		360		Same as above								0
Same as above		370		Same as above								0
		383		Hole bottomed at 383								0