

DIAMOND DRILL LOG

HOLE NUMBER	04 KEL 5
DATE DRILLED	July 29-30, 2004
AZIMUTH	210°
DIP OF HOLE	-65°
CASING DEPTH	3.35 meters
BEDROCK DEPTH	5.45 meters
LENGTH OF HOLE	30.48 meters (100 feet)
CORE SIZE	BQTW
NORTHING	6824632N
EASTING	573381E
UTM ZONE	7
UTM DATUM	NAD 83
LOCATION	Lower Canyon, Reed Creek, Whitehorse Mining District
NTS	115-G-12
LOGGED BY	Jim McFaull
CLIENT	Kelli Creek Group
DRILLED BY	E. Caron Diamond Drilling Ltd.

BOX 1

0.0 -2.00 m No recovery-casing placer tailings.

2.00- 5.45 m Placer tailings.

5.45- 8.30 m Pale grey/white quartz sericite schist cut by occasional narrow white quartz and carbonate veinlets. No HCl reaction on the sericite schist and a weak HCl reaction from the carbonate veinlets. Trace very fine grained disseminated pyrite. Minor interbeds of black limey graphitic schist with a strong HCl reaction and trace very fine grained disseminated pyrite @ 8.10- 8.21m. Narrow white quartz veins cut the graphitic schist, one of which has a small vug lined with grey quartz crystals @ 8.40m. core loss of 0.10m.

BOX 2

8.30- 10.67 m Pale grey/white quartz sericite schist cut by occasional narrow white quartz and carbonate veinlets. No HCl reaction on the sericite schist and a weak HCl reaction from the carbonate veinlets. Trace very fine grained disseminated pyrite. Minor interbeds of black limey graphitic schist with strong HCl reaction and trace very fine grained disseminated pyrite @ 8.30- 8.50m and @ 9.45- 10.51m.

10.67- 13.05 m Black limey graphitic schist with strong HCl reaction. Trace very fine grained disseminated pyrite. Schist is cut by narrow white quartz and carbonate veinlets. Foliation variable but mostly @ 70° TCA.

BOX 3

13.05- 17.20 m Black limey graphitic schist with strong HCl reaction. Trace very fine grained disseminated pyrite. Schist is cut by narrow white quartz and carbonate veinlets. Foliation variable but mostly @ 70° TCA.

17.20- 18.29 m Medium green chloritic schist (metavolcanics) with a weak HCl reaction from narrow carbonate veinlets. Trace very fine grained disseminated pyrite. Hangingwall contact conforms to foliation @ 70° TCA.

BOX 4

18.29- 21.60 m Medium green chloritic schist (metavolcanics) with weak HCl reaction from narrow carbonate veinlets. Trace very fine grained disseminated pyrite. Footwall contact conformable with foliation @ 60° TCA.

21.60- 21.76 m Black graphitic schist bed grades rapidly into quartz sericite schist.

21.76- 23.54 m Tan/pale green/grey quartz sericite schist with no HCl reaction. Trace very fine grained disseminated pyrite. Cut by narrow quartz veins.

BOX 5

23.54- 26.16 m Tan quartz sericite schist faulted to gouge @ 25.20- 25.42m with minor black graphitic schist in the fault. Core loss of 0.07m. Footwall contact @ 26.16m.

26.16- 30.48 m Black limey graphitic schist with strong HCl reaction. Trace very fine grained disseminated pyrite. Foliation @ 70° TCA.

END OF HOLE 04 KEL 5

CORE RECOVERY

30.48 m drilled = 100.00%

28.31 m recovered = 92.88%

2.17 m lost = 7.12%

ASSAY SAMPLES 04 KEL 5

SAMPLE #	INTERVAL
K 089	5.45 – 9.14 m
K 090	9.14 -12.19
K 091	12.19 -15.24
K 092	15.24 -18.29
K 093	18.29 -21.34
K 094	21.34 -24.38
K 095	24.38 -27.43
K 096	27.43 -30.48
End of Hole	