

17.20- 18.29 m Medium green chloritic schist (metavolcanics) with a weak HCl reaction from narrow carbonate veinlets. Trace very fine grained disseminated pyrite. Hangingwall contact conforms to foliation @ 70° TCA.

BOX 4

18.29- 21.60 m Medium green chloritic schist (metavolcanics) with weak HCl reaction from narrow carbonate veinlets. Trace very fine grained disseminated pyrite. Footwall contact conformable with foliation @ 60° TCA.

21.60- 21.76 m Black graphitic schist bed grades rapidly into quartz sericite schist.

21.76- 23.54 m Tan/pale green/grey quartz sericite schist with no HCl reaction. Trace very fine grained disseminated pyrite. Cut by narrow quartz veins.

BOX 5

23.54- 26.16 m Tan quartz sericite schist faulted to gouge @ 25.20- 25.42m with minor black graphitic schist in the fault. Core loss of 0.07m. Footwall contact @ 26.16m.

26.16- 30.48 m Black limey graphitic schist with strong HCl reaction. Trace very fine grained disseminated pyrite. Foliation @ 70° TCA.

END OF HOLE 04 KEL 5

CORE RECOVERY

30.48 m drilled = 100.00%

28.31 m recovered = 92.88%

2.17 m lost = 7.12%

ASSAY SAMPLES 04 KEL 5

SAMPLE #	INTERVAL
K 089	5.45 – 9.14 m
K 090	9.14 -12.19
K 091	12.19 -15.24
K 092	15.24 -18.29
K 093	18.29 -21.34
K 094	21.34 -24.38
K 095	24.38 -27.43
K 096	27.43 -30.48
End of Hole	



SAMPLE#	Au** ppb	Sample kg
K067	4	4.61
K068	<2	4.90
K069	2	4.16
K070	7	2.31
K071	191	3.91
K072	10	2.26
K073	4	3.45
K074	4	3.81
K075	5	3.51
K076	4	4.10
K077	8	4.35
K078	17	4.30
K079	12	5.05
K080	8	6.69
RE K080	6	-
RRE K080	9	-
K081	65	4.02
K082	110	4.26
K083	21	3.44
K084	135	4.27
K085	41	3.15
K086	150	3.84
K087	169	3.56
K088	172	3.65
K089	5	5.44
K090	3	3.75
K091	8	4.76
K092	14	4.70
K093	16	5.39
K094	14	3.56
K095	96	3.50
K096	25	4.30
STANDARD AU-R2	589	-

Sample type: CORE R150 60C. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.