

**SYNOPTIC LOG**  
**NORDAC RESOURCES LTD.**

Property: SIMPSON

Hole: SM-97-03 Section: 10,000 N

Page 1 of 1

Easting: 468,259.42 Northing: 6,735,859.01 Elevation: 1,551.31 Depth: 96.62 Logger: Sefika Lesnikoff  
Drilling Dates: July 21-23, 1997

Depth	0.00	96.62			
Azimuth	40	40			
Dip	-55	-58			
Method	Brunton	Acid			

From (m)	To (m)	Interval (m)	*	Unit	Comments	From (m)	To (m)	Interval (m)	Sample No.	REC %	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Au (ppb)
0.00	3.38	3.38		CSDH	casing										
3.38	14.93	11.55	T	CMSH	quartz-muscovite-chlorite schist										
14.93	23.70	8.77	T	QCSH	clay altered quartz-chlorite schist										
23.70	39.93	16.23	T	CMSH	quartz-chlorite-muscovite schist										
					-trace pyrite, chalcopyrite and galena										
39.93	48.32	8.39	S	CMSH	mineralized quartz-chlorite-muscovite schist										
					-sphalerite & galena occurs as bands or as disseminations	39.56	41.56	2.00	59307	87	56	56	2670	0.6	n/a
					-disseminated chalcopyrite	41.86	43.86	2.00	59308	100	34	348	1670	1.4	n/a
						41.56	41.86	0.30	59309	100	1890	594	7.82%	31.0	n/a
						43.86	46.32	2.46	59310	100	11	28	490	<0.2	n/a
						46.32	48.97	2.65	59311	100	95	442	1835	1.2	n/a
48.32	50.90	4.58	T	CMSH	quartz-chlorite-muscovite schist										
50.90	54.34	3.44	T	CMSH	quartz-chlorite-muscovite schist										
					-gouges at 53.30m-53.50m and 54.14m-54.34m										
54.34	63.65	9.31	T	CMSH	quartz-chlorite-muscovite schist										
63.65	64.89	1.24	T	QFMT	metatuff										
64.89	66.33	1.44	T	QZVN	quartz vein										
					-trace galena										
66.33	73.20	6.87	T	CMSH	quartz-muscovite chlorite schist with relict quartz phenocrysts										
73.20	79.85	6.65	S	CMSH	coarse grained and oxidized quartz-chlorite-muscovite schist with relict quartz and feldspar phenocrysts										
79.85	96.62	16.77	F	CMSH	quartz-muscovite-chlorite schist with coarse grained relict quartz phenocrysts										
					EOH										

\*S-strong weathering, T-transitional weathering, F-fresh