


HOLE TH13  
 LOCATION: CONTACT  
 LENGTH: 312'  
 DIP: -50° AZIMUTH: 030°  
 COLLAR ELEVATION: 5000'



- [1] SYENITE tan weathering, medium to coarse grained nepheline syenite often with abundant sulphides and purple fluorite
- [2] PSEUDOLEUCITE TINGUAITE 5-30%, 3-20mm pseudoleucite phenocrysts 1-10mm orthoclase phenocrysts in a medium grey, aphanitic groundmass
- [3] SHEARED TINGUAITE foliated and/or auto-brecciated tinguaites often biotite altered
- [4] MIXED TINGUAITE lenses or bands of pseudoleucite phenocrysts in sheared tinguaites groundmass
- [5] FINE GRAINED TINGUAITE medium to dark grey tinguaites without pseudoleucite phenocrysts, some 1-3 mm orthoclase phenocrysts. Usually as dykes
- [6] INTERMEDIATE TINGUAITE medium grey tinguaites with pseudoleucite orthoclase phenocrysts. Pseudoleucite phenocrysts smaller than usual

B,b BIOTITE ALTERATION . intense, moderate  
 C,c CHLORITE ALTERATION intense, moderate

— — — — — geologic contact: sharp, gradational




FIG. U-TG28  
 ARCHER, CATHRO & ASSOCIATES LTD  
**SECTION TH13**  
 TOMBSTONE PROPERTY  
 UKON JOINT VENTURE

