

R.G. MILNER
CONSULTING GEOLOGIST
WHITEHORSE YUKON TERRITORY

PROPERTY <u>TINTINA SILVER</u>	Claim No. <u>Eng 5</u>	Strike <u>S 50° W</u>	Lat. <u>79° 22'</u> <u>2357.3</u>	Hole No. <u>74-D, 13</u>
Date <u>Aug 9</u> 19 <u>74</u>	Section No. <u>23+000</u>	Dip <u>-40°</u>	Dep. <u>2309.8</u> <u>5451.4</u>	Total Depth <u>153-80</u>
Logged By <u>G. Carlson</u>	Plan No. <u>74-D</u>	Level <u>3-D7</u>	Elev. <u>5510</u>	Page No. <u>1</u>

[illegible]

R.B. MILNER
CONSULTING GEOLOGIST
WHITEHORSE YUKON TERRITORY

PROPERTY <u>TINTINA SILVER</u>	Claim No. _____	Strike _____	Lat. _____	Hole No. <u>74-713</u>
Date <u>Aug 9</u> 19 <u>74</u>	Section No. _____	Dip _____	Dep. _____	Total Depth <u>15880</u>
Logged By <u>C. Carlson</u>	Plan No. _____	Level _____	Elev. _____	Page No. <u>2</u>

FOOTAGE		ROCK CLASSIFICATION EPID. DIOP. GARN. SERP. QTZ/SIL. ACTINO. TREMO. CHLO. CRYSTALLINE. SHEARING. VEINS. FRACTURING. FOLIATION. GRAIN SIZE. TEXTURE	MINERALIZATION		ASSAY DATA							
From	To		TYPE	%	SAMPLE NO.	WIDTH	RECOV.	%CU	%FE	MOLY	AU/AG	IN.
32.5	62	LIMESTONE - mainly light grey, mottled, local argillaceous sections. - po + py appears to be associated mainly with argillaceous material - foliation quite variable, 10-50°, average 30° → change in argillaceous rock.	po, py	0-2								
62	80	LAMPROPHYRE - fine to medium grained - typical, biotitic, and white patches of variable size and concentration throughout, mainly less than 1/4" diameter - biotite shows light foliation, 0-20° to core down.										

80

END OF HOLE