




SILVER ARROW VEIN

- 1969 reserves of 2558 tonnes grading 695 g/ton Ag and 42.5% Pb. (22.3 oz./ton)
- mined in 1979 by Silver Arrow Syndicate.
- average surface value from 4 samples collected in 1987 was 54.15 oz./ton Ag.


LEGEND

- CRETACEOUS AND (NEARLY) TERTIARY**
- KTap Dark green, fine grained biotite-bearing mafic dikes.
 - Kg Homogenous, medium grained, biotite quartz monzonite.
- LATE DEVONIAN AND MISSISSIPPIAN**
- Mv Undifferentiated felsic and mafic volcanics, hornblende syenite, and black shale.
 - uDMs Black shale, chert, and chert conglomerate.
- SILURIAN, EARLY AND MIDDLE DEVONIAN**
- Sd Buff, grey, and red weathering dolomite, with lenses of massive quartz ore.
 - Ss Grey weathering platy, thinly laminated dolomitic siltstone.
 - Sq Massive grey weathering quartz arenite.
- ORDOVICIAN AND SILURIAN**
- OSsl Black, graphitic shale, minor chert.
- LATE CAMBRIAN AND EARLY ORDOVICIAN**
- uEOsl Grey-buff weathering thinly laminated calcareous phyllite, lufaceous phyllite, with some mafic luffs, and flows.
 - uEOB Resistant dark green mafic flow or sill.
- EARLY CAMBRIAN**
- Kcsl Grey weathering calcareous mica schist and marble.

SYMBOLS

- Geological contact: defined, approx., assumed.
- Bedding: inclined, vertical.
- Steep dipping fault: sense of movement.
- Unknown: defined approx.
- Normal fault: defined, approx.
- Thrust fault: defined, approx.
- Outcrop.
- Claim post, claim boundary.
- Main road, 4x4 road, cat trail.
- Rock sample site and number.

SAMPLE NUMBER	WIDTH (cm)	Ag (oz./ton)	%Pb	%Zn
18800	grab	49.0		
18801	grab	47.9		
18802	grab	43.8		
18804	float	13.9		
18824	float	36.5		
18836	grab	8.08	11.1	0.03
18879	grab	501.61		



092007 Geology after J.G. Abbot (1986).

YUKON MINERALS - PERREX J.V.

HV PROPERTY 092010

PROPERTY GEOLOGY - NORTH

GROUNDHOG CREEK, YUKON TERRITORY

NTS: 105 F/10	TECH: G.D.	DATE: NOV., 1987
SCALE: 1:10,000	DRAFTING: <i>HYDROGRAPHICS LTD.</i>	FIGURE: 7