LEGEND

UNIT No.	LITHOLOGY	PRISMA COLOR	AGE
12	Unconsolidated sands and gravels	927	Pleistocene
11	Quartz Diorite, Feldspar Porphyry	930	Tertiary
10	Anvil Batholith (Granite, Granodiorite, Quartz Monzonite)	929	Cretaceous
9	Basic Intrusive – Ultrabasic Rocks: Peridotite, Serpentine	932	Mississippian or later
	Diorite, Gabbro	934	
8	Basic Extrusive – Flows,	911	"
	Fragmental,	909	"
	Amphibolite, Hornblende schist (metamorphosed equivalent of above)	935	"
7	Conglomerate (Fanglomerate?) composed of pebbles of chert, quartz and volcanics	916	"
6	Limestone	904	"
5	Argillite	936	"
4	Chlorite schist, Chloritic Phyllite (sedimentary in origin)	920	"
3	Graphite Schist, Graphitic Phyllite	937	"
2	Biotitic Phyllite, Biotite schist	941	"
1	Quartzites, Quartz schist, Sericite schist	942	"

NOTE:

a Mye formation
b Vangorda formation
c Rose formation
d Un-named conglomerate (Fanglomerate?)
e Lap formation

GEOLOGICAL CONTACT

Firm or observed
Probable contact
Inferred contact
Contact very uncertain

AIRPHOTO LINEARS

Firm
Probable
Possible

BEDDING
SCHISTOCITY
FOLIATION
JOINT

FAULT

Observed
Probable

DIAMOND DRILL HOLES ●
ROTARY DRILL HOLES ○

MINERALIZATION x Cu, Pb, Zn, Po
SULPHIDE BODY — FARO

inches 0 1 2
centimetres 0 1 2 3 4 5
YUKON GEOLOGICAL SURVEY
This reference scale bar was prepared from the original mosaic & will scale at the same rate as the mosaic which can be used as a reference for the original size.

133° 30' LONG

N

A N V I L C R E E K

COVERED C R E E K

R O S E C R E E K

P E L L Y R I V E R

VANGORDA CREEK

UNMAPPED SEDIMENTS

NO OUTCROP

FAULTED CONTACT DIORITE / CHERT

RIBEL BLEACHED & ALTERED FRACTURED FAULTED BRACCIA SILICIFIED GRAPHITIC SERICITE, CHLORITE SCHIST ZONE. ORIENTED ALONG MOVEMENT.

62° 18' LAT.

LONG 133° 30'

LAT. 62° 18'

ANVIL MINING CORP.	
WHITEHORSE	
REGIONAL GEOLOGY SOUTH WEST ANVIL DIST. N.W. AREA	
DATE: 3 APRIL 1967	DRAWING NR. GWA-31
SCALE: 1" = 1/2 MILE	FILE /4
DRAWN: JG	
GEOLOGY: D. MAYES, R.S. ADAMSON, ET AL	

REVISED	BY	
19 NOV. 68	A.M.	

000331

DATE PRINTED: