

006729

DIAMOND DRILL RECORD

PROPERTY YAGORDA

Handwritten:
 April 15, 1977
 D.S.

SHEET NUMBER 1

SECTION FROM _____ TO _____

STARTED September 10, 1953

LATITUDE 30,034.63 9145.53N

DATUM _____

COMPLETED September 27, 1953

DEPARTURE 30,012.38 E 9147.77 E

BEARING _____

ULTIMATE DEPTH 198'

ELEVATION 3,999.50^{17.9} 1298.0 m

DIP = 90°

PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
						Ag(oz/t)	% Cu	% Pb	% Zn	Ag(oz/t)	% Cu	% Pb	% Zn	
4.3 0-11	Casing													
14-18 ^{5.5}	Ground core - sludges show heavy lead, zinc & iron sulphides	4051			4					1.5	0.64	3.92	3.69	
18-20 ^{1.0}	Ground core - sludges show heavy lead, zinc and iron sulphides	4052			2					2.24	0.98	4.47	8.57	
20-25 ^{7.4}	Ground core - sludges show as above	4053			5					2.34	0.91	5.56	6.38	
25-30 ^{2.1}	No core - sludges as above	4054			5					3.40	0.71	8.39	10.67	
30-35 ^{11.7}	No core - sludges as above	4055			5					12.80	0.71	9.92	8.87	
35-40 ^{12.2}	No core - sludges as above	4056			5					3.20	0.78	7.63	6.98	
40-45 ^{13.7}	No core - appears higher in lead, zinc in the sludges	4057			5					3.08	0.91	7.69	8.37	
45-50 ^{15.2}	Ground core - some gravelled core retained in barrel - consists of quartzite heavily mineralized with lead, zinc and iron sulphides - zinc is in form of resinous sphalerite - bedding is horizontal	4058			5					2.06	0.37	3.92	6.68	
50-55 ^{16.8}	Ground core - 1.0' coarse gravelled portions consisting of quartzite with bedding horizontal - sulphides of copper, lead, zinc, iron present - sludges consist of Pb, Zn & Fe sulphides	4059			5					0.98	0.25	2.29	1.99	
55-58 ^{18.7}	Six inches of gravelled core recovered - pyrite, sphalerite and some galena - approximately 25% pyrite, 10% combined lead-zinc	4060			3					1.28	0.44	2.29	2.39	
		4068			3	1.34	0.22	3.46	3.29					

Campbell

DIAMOND DRILL RECORD,

HOLE NO. 7-1

PROPERTY VANGORBA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
						Ag(oz/t)	% Cu	% Pb	% Zn	Ag(oz/t)	% Cu	% Pb	% Zn	
18.3														
58-60	1.8' recovered gravelled core - bluish quartzite - 20% sulphides - principally pyrite - some galena & sphalerite - sludge appears to contain mostly pyrite	4061 4009			2	0.72	0.25	1.76	1.89 ✓	0.76	0.29	1.53	1.79	
18.9														
60-62	0.7' partially ground core recovered - consists of quartzite and fine beds of schist imparting a gneissic appearance to the core sulphide replaced 40% of rock - principally pyrite with some sphalerite and the occasional crystal of galena	4062 4010			2	0.56	0.22	1.01	2.09 ✓	0.72	0.54	1.62	2.29	
19.3														
62-65	2.6' core recovered - consists of quartzite and narrow horizons of schist as described above - the quartz predominates and in places has a milky massive appearance as opposed to the granular quartzite exhibited in the surface showing - this may be due to the degree of replacement and type (i.e.) which is greater in the latter - the schistose layers are graphitic in places - sulphides have replaced 30% of core - predominately pyrite with minor amounts of galena and sphalerite.	4063 4011			3	0.50	0.25	.95	1.30 ✓	0.62	0.32	0.93	0.80	
20.3														
65-66.5	1.5' partially ground core recovered - principally quartzite with some schistose layers - 20% replaced with pyrite - no sphalerite or galena noted	4064 4012			1.5	1.00	.17	3.30	2.79 ✓	0.84	0.27	2.94	5.19	

DRILLED BY _____

SIGNED F.A. CAMPBELL

DIAMOND DRILL RECORD,

HOLE NO. Yan # 1

PROPERTY YANGONDA

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

1

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag(oz/t)	% Cu	% Pb	% Zn	Ag(oz/t)	% Cu	% Pb	% Zn
21.3 66.5-70	3.0' of core recovered - quartzite & schist - some minor drag folding exhibited - 15% replaced by sulphides principally pyrite with sphalerite and some galena.	4065								1.30	0.67	2.86	3.39
22.9 70-75	2.3' core recovered - consists of quartzite with schistose layers - 15% replaced with sulphides - mainly pyrite & sphalerite with some galena	4013 4066 4014				1.20	0.15	3.43	2.89 ✓	1.20	0.22	3.68	3.29 ✓
23.8 75-78	2.8' core recovered - lost water & sludge in slip planes - core consists of quartzite with schistose horizons - 10% sulphide replacement mainly pyrite & sphalerite, some galena	4015			3	1.56	0.15	4.30	5.28 ✓				
24.7 78-81	2.6' core recovered - quartzite & schist - 10% sulphides - mainly pyrite & sphalerite with minor amount of galena	4067 4016			3	.94	0.15	3.16	2.49 ✓	1.08	1.08	3.05	3.59
25.3 81-83	1.5' core recovered - quartzite & schist - 10% sulphides - mainly pyrite & sphalerite with some galena	4068 4017				1.86	0.13	2.40	2.79 ✓	0.68	0.25	2.40	3.58
25.9 83-85	1.3' core recovered - more schistose material with the quartzite - 5% sulphides - mostly pyrite with minor amount of sphalerite - no galena noted - lost water & sludge	4018				0.66	0.15	1.84	1.00 ✓				
26.8 85-88	3.0' core recovered - quartzite & schist in almost equal proportions - 5% sulphides between 85 and 86.5 mainly pyrite with some sphalerite - massive pyrite 86.5 to 88.0'	4069 4019 4020	85 86.5	86.5 88.0		1.10 2.60	0.22 1.32	2.76 6.23	4.88 ✓ 8.27 ✓	0.98	0.63	2.51	4.18

4.50 6.58

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 1

PROPERTY VANGORDA

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

#1

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
						Ag(oz/t)	% Cu	% Pb	% Zn	Ag(oz/t)	% Cu	% Pb	% Zn	
27.4 88-90	Leached zone - lost core & sludge					LORE								
28.9 90-95	Sericite schist - talcose in places - some quartz bearing fine pyrite - no sphalerite or galena noted - 90° may be lower contact of the favourable quartz zone	4070 4021												
30.5 95-100	2.2' recovered - sericite schist - talcose in part - some fine pyrite - no galena or sphalerite noted - some quartz interlaminated with schist.	4071 4022												
31.1 100-102	1.4' recovered - sericite schist - 0.2' quartzite bearing fine pyrite (BIOGEN) & galena - bedding at 25° to core.	4072 4023	100	105	5'					0.46	0.15	0.81	1.89	
32.0 102-105	1.7' recovered - 0.9' quartzite containing some pyrite & sphalerite - remainder is sericite schist	4024												
33.2 105-109	Four feet of partially ground core recovered - sericite schist - no visible sulphides - bedding at 10° to core	4073 4025												
34.4 109-113	3.8' recovered - sericite schist - some partially leached pyrite present - bedding at 50° to core - lost water - not able to drive Excasing	4026												
36.0 113-118	2.6' recovered - sericite schist - talcose in part - some quartz present - no visible sulphides	4027												

DRILLED BY _____

SIGNED P.A. CAMPBELL

DIAMOND DRILL RECORD,

HOLE NO. Van # 1

PROPERTY VANGORDA

SHEET NUMBER 5 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
						Ag(oz/t)	% Cu	% Pb	% Zn	Ag(oz/t)	% Cu	% Pb	% Zn	
37.5 118-123	1.0' recovered - sericite schist - talcose in part - some quartz present - no visible sulphides	4028 C												
38.4 123-126	1.3' recovered - sericite schist - considerable contortion in the bedding showing minor drag folds	4029 C												
39.3 126-129	2.5' recovered - sericite schist - graphitic in part - bedding at 25° to core - some fine pyrite present	4030 C												
39.9 129-131	2.0' recovered - sericite schist with quartz increasing in lower foot - bedding at 25° to core	4031 C												
40.5 131-133	1.7' recovered - quartzite - (80 to 90%) replaced by sulphides - principally pyrite with some sphalerite & only very little galena	4032 C			2	2.96	0.07	5.05	11.35	✓				
41.6 133-136	5 One foot of partially ground core recovered quartzite - (70%) replaced by sulphides - mostly pyrite and some sphalerite - no galena noted	4033 C			3.5	3.30	nil	6.58	8.97	✓				
42.7 136.5-140	3.3' recovered - quartzite 90% replaced by sulphides - very fine galena & sphalerite up to 10% combined - remainder is pyrite	4034 C			3.5	2.04	0.32	4.50	5.78	✓				
44.2 140-145	5.0' recovered - quartzite 90% replaced by sulphides - principally pyrite with up to 10% lead-zinc sulphides	4035 C			5	1.62	0.32	2.97	4.78	✓				

DRILLED BY _____

SIGNED E. A. CAMPBELL

DIAMOND DRILL RECORD,

HOLE NO. VVI

PROPERTY VANGORDA

SHEET NUMBER 6 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

#1

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag/oz/t	% Cu	% Pb	% Zn	Ag/oz/t	% Cu	% Pb	% Zn
45.7 145-50	4.8' recovered - 2.6' 90% replaced quartzite - the remainder consists of quartzite with some fine disseminated cubic pyrite - in the highly replaced quartzite the galena and sphalerite are present up to 10% combined	4036				0.80	0.49	1.45	1.99				
47.2 150-155	4.2' recovered - quartzite 90% replaced by sulphides from 150 to 151.8' - 151.8 to 155 consists of bands of quartzite sericite schist & sulphides - the latter having replaced up to 50% - sulphides principally pyrite with up to 10% combined lead-zinc sulphides	4037				0.74	0.29	1.26	1.49				
48.1 155-158	0.9' core recovered - quartzite with some sericite - 30% replaced - up to 10% lead-zinc combined	4074 (155-157)								tr.	0.13	0.08	0.55
49.1 158-161	0.4' badly ground core recovered - consists of quartzite, graphite & fine pyrite	4038 (157-159)				0.32	0.12	0.15	0.35				
49.7 161-163	No core - sludge black & muddy - may be graphitic - some fine pyrite	4076 (159-161)											
50.3 163-165	No core - sludge as above	4077											
50.9 165-167	No core - sludge as above	4078											
51.5 167-169	No core - sludge as above	4079											
52.1 169-171	No core - sludge as above	4080 4081											

DRILLED BY _____

SIGNED F. A. CAMPBELL

DIAMOND DRILL RECORD,

HOLE N. Van # 1

PROPERTY VANGORDA

SHEET NUMBER 7 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____ # /
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag(oz/t)	% Cu	% Pb	% Zn	Ag(oz/t)	% Cu	% Pb	% Zn
^{52.7} 171-173	No core - sludge as above	4082											
^{53.3} 173-175	No core - sludge as above	4083											
^{53.9} 175-177	No core - sludge as above	4084											
^{54.5} 177-179	No core - sludge as above	4085											
^{54.9} 179-180	0.9' recovered - dark gray graphitic schist silicified along bedding planes - fine pyrite present - bedding at 500.	4086											
^{55.5} 180-182	1.7' recovered - graphitic schist - highly contorted - contains quartz and fine pyrite	4087											
^{56.4} 182-185	2.4' recovered - graphitic schist - quartz and pyrite present - minor drag folding	4088											
^{57.0} 185-187	1.5' recovered - graphitic schist - silky white quartz - fine pyrite present - highly contorted	4089 (185-187)											
^{57.6} 187-189	1.7' recovered - graphite schist - quartz present imparts gneissic appearance - some fine pyrite	(187-189)											
^{58.1} 189-190.5	1.4' recovered - graphite schist with quartz and fine pyrite	4090 (189-198)											
^{58.7} 190.5-192.5	1.9' recovered - graphite schist - quartz and fine pyrite - highly contorted												
^{59.4} 192.5-195	2.4' recovered - graphite schist - some quartz and fine pyrite												
^{60.3} 195-198	2.6' recovered - schist becoming more sericitic - some quartz and pyrite												

End of Hole 198'

DRILLED BY _____

SIGNED E.A. CAMPBELL

DIAMOND LOG RECORD

PROPERTY - VANGORDA

HOLE NO. Van #2

Sheet 1

Latitude 30,066.86N 7154.56
 Departure 29,788.12E 7077.42
 Elevation 4,009.83+259 1301.14

Dip 90°

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
0.0 - 14.0 (A.2.1.4)		Casing
14.0 - 16.0	1.2	Thinly-bedded black graphitic schist. Soft. Rusty surfaces common. Bedding flat. No mineral.
16.0 - 18.0	0.7	As above.
18.0 - 20.0	1.0	As above.
20.0 - 23.0	1.2	As above - bedding varies to 25° to core.
23.0 - 25.0	0.7	As above.
25.0 - 28.0	0.6	As above - bedding flat to 10° to core.
28.0 - 30.0	0.8	As above.
30.0 - 32.0	0.5	As above - some contortion in bedding.
32.0 - 34.0	0.6	As above.
34.0 - 36.0	1.2	As above.
36.0 - 38.0	1.1	As above - bedding contorted and drag folded.
38.0 - 40.0	0.8	Soft black graphitic sericite schist. Bedding flat to 10° to core.
40.0 - 42.0	0.6	As above - bedding locally 18° to core.
42.0 - 44.0	0.8	As above.
44.0 - 46.0	0.9	As above.
46.0 - 48.0	1.0	As above.

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
48.0 - 50.0	0.4	As above. Bedding contorted.
50.0 - 52.0	0.8	As above. Bedding flat to 10° to core. Scattered grains pyrite
52.0 - 54.0	1.0	As above.
54.0 - 56.0	1.2	Graphitic sericite schist. Bedding 10° to core. Silicified. Narrow threads quartz along shearing. Some brassy pyrite and pyrrhotite with quartz. Also in scattered grains and tiny fracture fillings.
56.0 - 59.0	1.4	Black schist. Bedding contorted. Some quartz with brassy pyrite.
59.0 - 61.0	0.6	Black graphitic schist. Brassy pyrite in fractures.
61.0 - 62.0	0.5	As above. quartz veinlets.
62.0 - 64.0	0.6	As above. Some pyrite.
64.0 - 66.0	0.6	As above.
66.0 - 67.0	0.8	As above - bedding contorted. Some at 10° to 15° to core. Injected lit-par-lit by narrow quartz veinlets. Minor brassy pyrite.
67.0 - 68.0	1.0	Graphitic sericite schist. Contorted and brecciated. Injected with quartz and carbonate. Fine pyrite common. 1" heavy pyrrhotite and pyrite at 68 feet.
68.0 - 71.0	1.7	Graphitic schist thinly interbedded with sericite schist. Bedding regular at 20° to core. Some pyrite and pyrrhotite along shear planes.
71.0 - 73.0	1.1	As above. Bedding contorted ranging from 15° to 30° to core. Locally brecciated and cemented with white carbonate. Some pyrite.
73.0 - 75.0	1.8	As above. Bedding regular at 10° to 20° to core. 1/4" massive pyrite with quartz at 73.7'
75.0 - 77.0	1.0	Dark gray schist.
77.0 - 79.0	1.4	Graphitic sericite schist. Bedding 25° to core.

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
79.0 - 81.0	0.6	As above. Bedding highly contorted.
81.0 - 83.0	0.4	Dark schist and white quartz.
83.0 - 85.0	0.4	Black graphitic schist.
85.0 - 87.0	1.6	As above - bedding 15° to core.
87.0 - 89.0	1.0	As above - Quartz stringers.
89.0 - 91.0	1.1	As above. 4" quartz vein at 90.7'. No mineral.
91.0 - 93.0	1.1	As above. Bedding 5° to 20° to core.
93.0 - 95.0	1.4	As above - 1" quartz stringer at 94.5' contains brown pyrrhotite
95.0 - 97.0	2.0	Black graphitic schist. Bedding 5° to core. Brown pyrrhotite in scattered narrow quartz stringers.
97.0 - 99.0	1.2	As above. Bedding 10° to core. Locally brecciated and cemented with white carbonate. Some pyrrhotite with the carbonate.
99.0 - 101.0	1.1	As above.
101.0 - 103.0	1.3	As above. Bedding 10° to core. Brown pyrrhotite replacement along bedding locally.
103.0 - 105.0	1.0	As above. locally brecciated and cemented by white carbonate. Brown pyrrhotite with the carbonate and locally along bedding.
105.0 - 107.0	1.5	As above. Bedding flat to 10° to core.
107.0 - 110.0	1.3	Black graphitic schist with interbedded sericite schist. Bedding 5° to core. Locally silicified. Brown pyrrhotite and some pyrite with quartz stringers and along shear planes.

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
110.0 - 112.0	1.0	Graphitic schist. Pyrite in fractures and along bedding.
112.0 - 114.0	2.0	As above. Shearing 10° to 15° to core but locally contorted and drag folded.
114.0 - 119.0	4.9	Graphitic sericite schist. Bedding at 10° to core where regular. Locally contorted drag folded and brecciated. Injected lit-par-lit by quartz and white carbonate. Minor amounts of pyrite and pyrrhotite.
119.0 - 124.0 -	4.8	Dark grey sericite schist. Bedding regular at 10° to core. Scattered narrow quartz stringers with minor amounts of brassy pyrite.
124.0 - 130.0	5.0	Dark grey sericite schist. Bedding contorted. Section cut by quartz veins and stringers. Quartz white and barren except for minor brassy pyrite.
130.0 - 132.0	0.7	Dark schist. Silicified and injected with quartz stringers. Some pyrrhotite.
132.0 - 136.0	3.4	Graphitic sericite schist with white quartz stringers. Bedding at 10° to core. Pyrite and pyrrhotite in some quartz stringers and locally along schist planes.
136.0 - 139.0	2.6	Graphitic sericite schist. Bedding flat to 10° to core. Three quartz stringers. Pyrite and pyrrhotite along bedding and in quartz.
139.0 - 143.0	3.0	Sericite schist. Bedding 10° to core. Quartz stringers but no mineral.
143.0 - 148.0	4.5	As above. Bedding 20° to 25° to core. Some quartz stringers and blebs.
148.0 - 154.0	4.7	As above. 40% replaced by white quartz. Brown pyrrhotite in much of the quartz. Some pyrite.
154.0 - 156.0	1.1	Sericite schist. Bedding 30° to core. Some quartz stringers with pyrite.
156.0 - 160.0	4.0	Graphitic sericite schist with scattered quartz stringers. Bedding 10° to 15° to core but locally drag folded and contorted. Brown pyrrhotite and brassy pyrite common in contorted sections and in quartz veins.
160.0 - 165.0	2.2	As above. 50% replaced by quartz. Some pyrite and pyrrhotite.

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
165.0 - 170.0	3.6	Sericite schist with graphitic sections. Bedding generally contorted but some at 16° to core. Some quartz. Some pyrrhotite and pyrite.
170.0 - 174.0	2.6	As above
174.0 - 178.0	3.1	Graphitic sericite schist. Bedding flat to 10° to core --locally contorted. Brown pyrrhotite common in small blebs and tiny veinlets thruout section. Less than 1% of volume made up by sulphides.
178.0 - 181.0	2.8	Black graphitic schist. Bedding at 10° to 15° to core, although locally contorted and drag folded. Brown pyrrhotite common in tiny blebs and veinlets. Some pyrite.
181.0 - 186.0	5.0	As above.
186.0 - 192.0	2.8	As above. Scattered quartz stringers with brown pyrrhotite.
192.0 - 195.0	2.3	As above.
195.0 - 197.0	1.8	As above.
197.0 - 200.0	2.2	As above. Pyrrhotite in one quartz stringer.
200.0 - 204.0	2.9	As above. Bedding at 15° to core. Some quartz stringers. Brown pyrrhotite in threads and blebs along shear planes.
204.0 - 207.0	3.0	As above.
207.0 - 210.0	1.8	As above. Bedding 20° to core. 3" quartz vein at 209.6 with some brown pyrrhotite.
210.0 - 213.0	2.0	As above. Bedding contorted. Local veinlets brown pyrrhotite.
213.0 - 216.0	2.1	As above. Bedding 15° to 25° to core. Brown Pyrrhotite in one quartz stringer & along bedding.
216.0 - 219.0	1.7	As above. Bedding 15° to core. Scattered quartz-carbonate veinlets with pyrite.
219.0 - 224.0	4.8	Graphitic sericite schist. Bedding locally contorted & drag folded. Where regular - at 15° to core. Injected with white carbonate carrying brown pyrrhotite. Barren blue-white quartz vein from 222.8 to 224.0
224.0		End of Hole.

(Signed) "H. D. McLeod"

DIAMOND DRILL RECORD

PROPERTY - VANGORDA

HOLE NO. Van #3

Latitude 29,965.15N 9133.38
 Departure 29,832.59E 9092.97
 Elevation 3,976.94 1291.11 Dip 90⁰

Started October 8, 1953
 Completed October 10, 1953
 Ultimate Depth 111.0'

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
0.0 - 11.0 ^{3.35m}		<u>Casing</u>
11.0 - 12.0	0.5	<u>Schist</u> - Fine-grained, black, graphitic sericite schist. Some carbonate veinlets. Shearing well developed but obscured by broken core. Angle indicated to be very steep. No mineral.
12.0 - 13.0	0.3	<u>Schist</u> - As above
13.0 - 15.0	-	<u>Ground core</u> - sludge black schist.
15.0 - 17.0	0.5	<u>Schist</u> - as above
17.0 - 20.0	0.5	<u>Schist</u> - as above
20.0 - 30.0	-	<u>Ground core</u> - sludge black schist
30.0 - 32.0	0.6	<u>Schist</u> - as above
32.0 - 34.0	0.7	<u>Schist</u> - Fine black graphitic sericite schist. Bedding obscured by broken core but indicated to be flat.
34.0 - 36.0	0.6	<u>Schist</u> - as above
36.0 - 38.0	1.0	<u>Schist</u> - as above
38.0 - 40.0	0.8	<u>Schist</u> - as above
40.0 - 41.0	0.8	<u>Schist</u> - as above. Bedding flat.
41.0 - 42.0	0.8	<u>Schist</u> - Black graphitic schist with quartz carbonate veinlets. Some brassy pyrite. Bedding 17 ⁰ to core. Some warping and one well-developed small dragfold.
42.0 - 44.0	-	<u>Ground core</u> - Sludge black schist.
44.0 - 45.0	0.5	<u>Schist</u> - as above.

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
45.0 - 27.0	0.7	<u>Schist</u> - as above
47.0 - 50.0	1.2	<u>Schist</u> - graphitic schist with some light coloured sericite - talc sections. Shearing flat.
50.0 - 52.0	0.4	<u>Schist</u> - as above
52.0 - 54.0	0.5	<u>Schist</u> - as above.
54.0 - 55.0	0.5	<u>Schist</u> - Sericite - talc schist with some graphite. Shearing 20° to 25° to core.
55.0 - 60.0	2.0	<u>Sericite schist</u> - quartz vein from 55.5 to 57.0. Quartz barren, and white. Searing at 10° to 15° to core.
60.0 - 62.0	1.8	<u>Schist</u> - light coloured sericite - talc schist with a quartz vein from 60.5 to 60.9. Shearing flat. No mineralization.
62.0 - 64.0	1.0	<u>Schist</u> - Talc sericite schist. Bedding flat to 10° to core.
64.0 - 65.0	0.5	<u>Schist</u> - as above. Bedding warped but at 15° to core.
65.0 - 70.0	0.6	<u>Schist</u> - as above
70.0 - 71.0	0.6	<u>Schist</u> - as above. 1" quartz stringer at 70.3 contains some brassy pyrite.
71.0 - 73.0	0.6	<u>Schist</u> - as above. Bedding flat to 10° to core.
73.0 - 75.0	0.6	<u>Schist</u> - darker in colour.
75.0 - 77.0	0.3	<u>Schist</u> - as above.
77.0 - 78.0	0.5	<u>Schist</u> - Sericite - talc variety with some graphite. Shearing highly contorted and drag folded.
78.0 - 80.0	0.7	<u>Schist</u> - As above. White carbonate veinlets along shear planes contain some pyrite.
80.0 - 82.0	1.0	<u>Schist</u> - Dark grey sericite - talc schist. Shearing warped but flat to 10° to core.

<u>Depth Feet</u>	<u>Recovery</u>	<u>Formation</u>
82.0 - 85.0	0.6	<u>Schist</u> - as above
85.0 - 86.0	0.4	<u>Schist</u> - as above
86.0 - 88.0	1.8	<u>Schist</u> - Sericite talc schist with some graphitic sections. Cut by quartz and quartz-carbonate veinlets carrying minor amounts of brassy pyrite. Shearing locally warped and drag folded. Rest at 25° to 30° to core.
88.0 - 89.0	1.0	<u>Schist</u> - Sericite variety. Shearing at 5° to 10° to core. Some indication here that the shearing and original bedding in the rock are at radically different angles. One piece of core has wavy banding along its length or vertical. This banding is warped and offset by the well-developed and almost flat shearing. One quartz stringer with minor pyrite.
89.0 - 91.0	1.6	<u>Schist</u> - Sericite - talc schist. One quartz stringer with some brown pyrrhotite. Shearing warped and drag folded.
91.0 - 95.0	4.4	<u>Schist</u> - Sericite - talc variety shearing at 10° to 15° to core. Some well developed drag folding. Scattered quartz carbonate veinlets with some pyrrhotite.
95.0 - 99.0	3.3	<u>Schist</u> - as above
99.0 - 100.0	0.6	<u>Schist</u> - as above
100.0 - 103.0	1.1	<u>Schist</u> - as above
103.0 - 105.0	1.1	<u>Schist</u> - Sericite - talc variety. Shearing well developed and regular at 10° to core.
105.0 - 107.0	1.0	<u>Schist</u> - as above.
107.0 - 111.0	3.3	<u>Schist</u> - Sericite - talc variety. Shearing at 10° to 15° to core- locally warped and drag folded. Some evidence of vertical bedding but almost destroyed by the flat shearing. Pyrrhotite in some small quartz carbonate veinlets.
111.0	-	End of hole.

(Signed) "H. D. McLeod"

DIAMOND DRILL RECORD,

HOLE NO. Van # 4

PROPERTY VANGORDA

the B.
 Date 15, 1977

SHEET NUMBER 1.

SECTION FROM _____ TO _____

STARTED Oct 13, 1953

LATITUDE 29.981.61 N 9138.4 N

DATUM _____

COMPLETED Oct 15, 1953

DEPARTURE 29.919.02 E 9119.32 E

BEARING _____

ULTIMATE DEPTH _____

ELEVATION 3987.23 1136.2 M

DIP -90°

PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
							Ag	Cu	Pb	Zn	Ag g/t	Cu %	Pb %	Zn %	
0.0-5.0	(15")	Casing													
5.0-7.0	0.3 chips	Massive brassy pyrite in quartzite. Minor amounts of sphalerite - possibly some galena.	4040 na	2	4143					2.94	0.35	3.92	3.19		
7-9	0.6 chips	Massive pyrite in quartzite. End of section consists of massive resinous sphalerite and pyrite with some galena	4041 na	2	4144					2.82	0.95	3.98	10.28		
9-10	0.6 chips	Massive sulphides. Some chips consist massive sphalerite, pyrite & galena - others of pyrite in quartzite. This suggests banding of the sulfides within the domain.	4042 na	1	4145	9-11				3.16	0.83	5.30	10.57		
10-11.5	0.8 Chips	Pyrite in massive blebs and stringers in quartz. Minor amounts of sphalerite, galena & chalcopyrite.	4043 na	1.5	4146	11-13				2.26	1.13	4.03	8.28		
11.5-14	0.7 chips	Same as above except pyrite is more abundant.. Banding of sulphides evident in one piece of core where a band of sphalerite rich sulphide occurs in massive pyrite. Some galena.	4044 na												
14-15	0.3 chips	Massive pyrite in quartzite: No recognisable sphalerite or galena.	4045 na	1	4147	13-15				3.62	1.75	4.11	8.77		
15-16	0.7	Massive sulphides replacing quartzite comprise 80% of volume and consist of sphalerite, pyrite & galena in order of abundance. Banding in sulphides evident. Galena not nearly so evident as in Hole Van #1.	4046 ✓	1	4148	15-17	4.14	1.66	6.81	15.38	4.02	0.93	6.73	13.46	

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Ym # 4

PROPERTY VANCOEBA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
							Ag	Cu	Pb	Zn	Ag %	Cu %	Pb %	Zn %
16-17	1.0	Massive sulphides with 10% to 15% quartz. Sulphides consist of sphalerite, pyrite & galena.	4047 ✓	1			5.10	0.28	8.90	17.32				
17-19 19-20	0.6 0.9	Same as above Massive sulphides with 20 to 25% quartz. Sphalerite, pyrite & galena in order of abundance. Galena much more abundant than before occurring in fine fractures throughout the quartz.	4048 ^{ma} 4049 ✓	2	4149		5.60	1.86	10.88	19.01	3.46	2.71	4.97	8.87
20-21 21-23	0.8 1.5	Massive sulphides as above Massive sulphides - sphalerite, pyrite & galena in order of abundance. 10% to 15% quartz. Galena in very fine grains and fracture fillings. Last section of core some schist and mineral, largely pyrite. One quartz veinlet cutting the banding in schist, contains some pyrite & sphalerite.	4050 ✓ 4201 ✓	1 2	4150 4151	19-21	5.20 4.36	0.58 0.78	9.99 8.57	17.53 13.42	8.90 3.24	1.13 2.61	10.60 4.87	17.35 8.87
23-25	0.8 chips	Massive pyrite with 20% to 25% quartzite. Bedding apparent in dark quartzite but angle indefinite. Minor amounts of sphalerite.	4202 ^{ma}	2	4152						1.34	2.10	1.02	3.59
25-27	0.8 chips	First half of section massy brassy pyrite with 15% to 20% quartz. Second half bedded quartzite 40% replaced by brassy pyrite. Minor amounts of sphalerite. Some galena in fine fractures in the quartz.	4203 ^{ma}	2	4153						0.46	0.60	nil	1.60

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 4

PROPERTY VANGORDA

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE		SLUDGE SAMPLE				
							Ag %	Cu %	Pb %	Zn %			
27-30	0.2 Chips	Bedded quartzite partly replaced by pyrite, sphalerite & galena	MA 4204	x 3	4154	27-29							
30-31	0.3 Chips	Same as above	MA 4205	x 1	4155	29-31							
31-32	0.4 Chips	Same as above	MA 4206		4156	31-33			0.46	0.63	Tr.	0.90	
32-33	0.6 Chips	Massive pyrite 75% replacing quartzite	MA 4207										
33-35		GROUND CORE - Sludge comprised of sulphides & quartz			4157				0.22	0.65	Tr	0.10	
35-36	0.5 Chips	Brassy pyrite. Quartzite 75% replaced.	MA 4208		4158	35-37			0.20	0.33	Tr	nil	
36-37	0.4 Chips	Quartzite 35% replaced by pyrite. Fine galena in tiny fractures in quartz.	MA 4209										
37-39	1.5	Massive pyrite with 10% to 15% quartz. No recognizable sphalerite or galena	4210	for Cu	4159		0.58		0.48	0.68	Tr	nil	
39-40	0.6 Chips	White quartzite 60% replaced by brassy pyrite. One chip dark gray schistose quartzite	MA 4211		4160	39-41			0.44	0.63	nil	0.50	
40-41	0.6 Chips	Quartzite 75% replaced by brassy pyrite. Minor amounts of galena in fractures in the quartz	MA 4212										

x 4204 & 4205 grouped together to provide enough material for assay.

DIAMOND DRILL RECORD,

HOLE NO. Van # 4

PROPERTY VANGORDA

SHEET NUMBER 4 SECTION FROM _____ TO _____

LATITUDE _____ DATUM _____

DEPARTURE _____ BEARING _____

ELEVATION _____ DIP _____

STARTED _____

COMPLETED _____

ULTIMATE DEPTH _____

PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
											Ag %	Cu %	Pb %	Zn %
41-45	0.3 Chips	Core recovered consists of brassy pyrite replacing quartzite. Sludges indicate sulphides extend to 42.0. Remainder of section black graphitic schist from which no core was recovered.	41213	41-2	4161	41-43					0.76	0.93	0.36	0.81
					4162	43-45					0.16	0.30	Tr	nil
45-50		<u>GROUND CORE</u> - Sludge black graphitic schist.			4163	45-47					0.24	0.23	nil	nil
50-53	0.2 chips	Chips recovered are half quartzite 30% replaced by pyrite & half black schist. Recovery came from last 6" of section. Remainder black graphitic sericite schist as indicated by sludge.	41214		4164	47-49					0.28	0.28	Tr	0.10
					4165	49-51					Tr	0.20	nil	nil
					4166	51-53					Tr	0.20	nil	nil
53-55		<u>GROUND CORE</u> - Sludge largely black graphitic schist but some pyrite - bearing quartzite interbedded. Schist & quartzite indicated to be interbedded in thin horizons.			4167						Tr	0.20	nil	0.40
55-57	0.4 Chips	Graphitic sericite schist with white quartz veins. Minor amounts brassy pyrite. Banding well developed but angle to core indefinite.			4168						Tr	0.30	nil	0.30
57-60	0.6	Dark grey to black graphitic schist. Banding highly contorted. Some pyrite in scattered quartz-carbonate stringers			4169						Tr	0.18	nil	0.10

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 4

PROPERTY YANGORRA

SHEET NUMBER 5

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core Sample NO.	Footage Sludge Footage		CORE SAMPLE	SLUDGE SAMPLE	
				FROM	TO		WIDTH	
60-62	0.3 chips	Black graphitic sericite schist		4170				
62-67		GROUND CORE Sludge - graphitic schist		4171				
67-71	0.7 chips	Black Graphitic schist		4172				
71-76	0.6	Black graphitic schist with a barren white quartz vein		4173				
76-81	0.9	Black graphitic schist with scattered quartz-carbonate veinlets. One 1" section silicified & mineralised with brassy pyrite. Banding well developed but angle indefinite due to broken nature of core. Indicated to have a low dip		4174				
81-86		GROUND CORE		4175				
86-91	0.7 chips	Black graphitic schist						
91-96	0.5 "	Same as above						
96-101	0.7 "	Same as above						
101-106	1.0	Black graphitic schist - shearing well developed - flat to 15° to core due to warping. Minor amounts brassy pyrite in carbonate veinlets.						
106-111	0.8	Black graphitic schist. Shearing flat. Brown pyrrhotite & pyrite in quartz-carbonate veinlets.						

DRILL RECORD,

HOLE NO. Van # 4

PROPERTY VANGORDA

SHEET NUMBER 6

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage			Sludge Footage			CORE SAMPLE			SLUDGE SAMPLE				
				FROM	TO	WIDTH											
111-115	0.9	Black graphitic schist. Shearing 20° to 25° to core.															
115-118	0.3	Few chips and sludge recovered. Black graphitic sericite schist.															
118-121	0.1 chips	Graphitic schist															
121-124	0.4 chips	Graphitic schist															
124		END OF HOLE.															

DRILLED BY _____

SIGNED [Signature]

DIAMOND DRILL RECORD,

HOLE NO. Van # 5

PROPERTY YANGORDA

SHEET NUMBER 1

SECTION FROM _____ TO _____

STARTED October 16, 1953

LATITUDE 28,481.06N

DATUM _____

COMPLETED _____

DEPARTURE 31,308.62E

BEARING N 43° 50 E

ULTIMATE DEPTH _____

ELEVATION 4059.07

DIP 44° 15 at surface, at 350' 46°

PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
							Ag %	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn	
0-22	(C.1) ¹⁰	Casing													
22-24 M	0.8	Massive brassy pyrite - leached at beginning of section	4215	6	4191	22-24	^{29.5} 0.86	0.38	1.11	1.69	1.44	0.60	2.08	2.18	
24-26 M	0.8	Massive pyrite - some highly leached	4216	7	4192	24-26	^{23.4} 0.80	0.43	0.97	2.29	1.18	0.96	0.99	1.49	
26-28 M	1.3	One 1/2" streak galena & sphalerite Massive sulphides - pyrite with galena some quartz & carbonate throughout - the galena occurs in very fine grains throughout the pyrite. Percentage impossible to estimate. End of section becomes massive pyrite.	4217	7	4193	26-28	^{44.6} 1.30	0.25	1.96	3.79	1.58	0.55	2.74	3.09	
28-33 M	3.8	28.0-31.7 Massive pyrite with galena in scattered narrow streaks and in white carbonate veinlets with sphalerite	4218	6	4194	28-30	^{37.7} 1.10	0.50	1.36	1.30	1.06	0.50	1.21	1.29	
					4195	30-32					2.14	0.28	3.73	4.28	
					4196	32-34					2.16	0.30	3.08	5.08	
		31.7-33.0 Massive galena with 15% pyrite - no sphalerite - galena very fine grained dull lead-gray variety	4219	1.3	4197	31.7-33	^{96.0} 2.80	0.38	4.40	7.08					
33-37 M	4.0	Massive sulphides with 20% white carbonate - sulphides 30% galena with the remaining being pyrite & some sphalerite	4220	4	4198	33-37	^{40.5} 2.64	0.18	4.83	7.48					
37-40 M	3.0	Massive pyrite with rounded blue to white quartz eyes. Pyrite and some chalcopyrite in fine fractures in the quartz. White carbonate blebs and stringers appear at the end of the section. Streaks of galena rich sulphides here also.	4221	3	4198	37-39	^{16.8} 2.24	0.18	4.28	6.98					

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 7m 45

PROPERTY YANGORDA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage			CORE SAMPLE				SLUDGE SAMPLE			
				FROM	TO	WIDTH	Ag %	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn
40-42 M	1.0	Blebs & stringers of massive pyrite in white quartz carbonate. Pyrite comprises 40% of section. Galena in fine fractures in the carbonate.	4222		4199	39-41	0.32 11.0	0.15	0.36	1.19	0.34	0.25	0.44	1.19
42-45 P	3.0	Bluish quartzite with thin streaks of sericite schist throughout. Section shattered & mineralised with brassy pyrite in massive stringers & scattered grains. Chalcopyrite common as tiny fracture fillings. Some rounded grains of a granular silvery mineral possibly arsenopyrite. Sulphides comprise 10% of the section.	4223	4200	41-43	0.88 30.2	0.73	0.47	1.10					
45-48 M	3.0	Massive pyrite with white carbonate and scattered narrow sections of bluish quartzite. Chalcopyrite common in scattered grains & tiny fracture fillings. Some sphalerite in the quartz. Very minor amounts of galena.	4224	4302	45-47	0.96 32.7	0.35	1.22	0.90					
48-52 M	4.0	48.0-51.0 Massive pyrite with fractured carbonate stringers. Pyrite fills in around fragments. Some chalcopyrite throughout in grains & tiny fracture fillings.	4225	4303	47-49	0.72 24.7	0.38	0.69	0.70					
		51.0-52.0 Bluish quartzite fractured & 20% replaced by sulphides - pyrite with some chalcopyrite & reddish sphalerite.		4304	49-51									

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Yan # 5

PROPERTY YANGORDA

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
							Ag %	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn	
52-60	6.2	Bluish quartzite 30% replaced by sulphides - sphalerite, pyrite & galena in a 5:1:2 ratio. Some chalcopyrite. Percentage of galena difficult to estimate as it occurs in a very fine dissemination throughout the sphalerite. Quartzite well bedded but highly contorted & fractured. Same as above - White quartz & yellow carbonate stringer cuts across the mineral.	4226	52-55	4305	52-53	1.76	0.18	3.63	7.57	1.62	0.25	2.96	6.18	
					4306	53-55	60.3				2.32	0.25	4.83	7.67	
				4227	55-60	4307	55-57	1.90	0.18	4.18	9.26	1.32	0.20	2.31	5.28
						4308	57-59	65.12				1.52	0.23	2.75	5.68
60-66	1.2	Same as above. Mineral less abundant at end of section. One piece sericite at 67'. Ground core - sludge light - gray sericite schist	4228	60-64	4309	59-61	2.66	0.20	5.28	10.68	2.26	0.25	4.94	9.87	
						4310	61-63	91.2				1.40	0.20	3.30	7.48
64-67	3.0	Same as above. Shearing not too pronounced due to contortion and brecciation	4229	64-65	4311	63-65	2.86	0.28	5.61	11.75	2.28	0.23	4.83	9.97	
					4312	65-67	98.0								
67-69						4313	67-69					4.24	0.25	0.55	1.89
69-75	0.6	Same as above. Banding highly contorted. Some white quartz.													
75-79	1.2														
79-82	1.5														
82-85	2.0														
85-90	2.5	Same as above. Banding highly contorted. Two white quartz stringers with some chalcopyrite.													

DRILLED BY _____

4312 not received

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van 15

PROPERTY VANGORDA

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE			SLUDGE SAMPLE			
90-95	2.0	Dark gray sericite schist. Banding regular at 15° to core. One white quartz stringer. Minor amounts brown pyrrhotite.											
95-100	0.6	Sericite schist. Brecciated & carbonatised.											
100-105	1.8	Same as above											
105-110	1.7	Sericite schist with white quartz stringers. Not so highly altered											
110-113	1.4	Light grey sericite schist. Shearing regular at 35° to core.											
113-116	2.6	Same as above.											
116-117	0.4	Same as above											
117-120	1.4	Sericite schist. Banding contorted Quartz stringers.											
120-122	1.4	Sericite schist. Quartz stringers. Shearing regular at 45° to core.											
122-126	2.6	Sericite schist. Contorted & injected with quartz stringers. Minor chalcopyrite in the quartz.											
126-133	3.5	Sericite schist. Some regular banding chips at 40° to core.											
133-135	0.8	Sericite schist. Banding at 35° to core.											
135-141.9	4.0	Sericite schist - Bedding ranges from 27° to 34° to core.											
141.9-142	0.1	M Sulphides replacing quartz. Mineral comprises 50% of volume. Sphalerite-galena & pyrite.											

DIAMOND DRILL RECORD,

HOLE NO. Van # 5.

PROPERTY VANGORDA

SHEET NUMBER 5. SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE				SLUDGE SAMPLE			
				FROM	TO	WIDTH	Ag %	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn
142-150	8.0	Massive sulphides 142.0-146.8 Quartzite 60% replaced by pyrite, galena & sphalerite. Pyrite most abundant. Galena & sphalerite occur in a very fine grained state making an estimate of amount almost impossible.	4235	4.1 142.9-146	4314	140-145	1.92 65.8	0.28	1.68	7.97	0.92	0.10	0.52	3.15
	M	146.8-147.8 Massive dark brown sphalerite with pyrite and some galena. White quartz eyes throughout.	4236	5 146-147	4315	145-148	2.78 95.3	0.30	3.19	9.06	2.44	0.10	2.09	6.67
150-154	4.0	147.8-150.0 Same as section 142.0 to 146.8. Massive sulphides 150.0-152.0 Quartzite 50% replaced by pyrite, sphalerite & galena. Lead and zinc abundant - 15% combined. Some masses resinous sphalerite in white carbonate veinlets. 152.0-154.0 Quartzite 60% replaced by pyrite with some sphalerite and galena. Latter two very fine grained and difficult to estimate.	4237	5 150-154	4316	150-151	2.04 69.9	0.30	3.74	7.18	2.40	0.13	2.85	7.47

DIAMOND DRILL RECORD,

HOLE NO. 1015

PROPERTY VANGORDA

SHEET NUMBER 6. SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage		CORE SAMPLE				SLUDGE SAMPLE				
				FROM	TO	Ag g/t	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn	
154-159	5.0	Massive sulphides 154.0-156.0 Quartzite 75% replaced by massive pyrite, Galena and sphalerite together & separately in streaks throughout the pyrite. Fine galena & sphalerite in unreplaced sections of quartz.		4317	155-160						2.16	0.15	4.50	6.77
		156.0-157.2 Pyrite, galena & sphalerite. Latter two 15% to 20% combined. Two colors sphalerite present, light resinous & deep brown. Mineral generally very fine grained although some coarse galena & sphalerite present.	4238		156.0-157.2	5	3.08	0.28	5.00	10.78				
		157.2-159.0 Quartzite 50% replaced by sulphides - pyrite, galena and sphalerite. Combined lead-zinc 10% approximately.												
159.0-160	1.0	Massive sulphides - same as Section 157.2-159.0 above	4239		157.2-162.0		2.30	0.30	3.96	8.28				
160-162	1.4	Massive sulphides - quartzite 50% replaced by sulphides		4318	160-165						1.84	0.18	3.51	5.50
162-169	7.0	Massive sulphides - massive pyrite with galena & sphalerite concentrated in streaks & stringers with white carbonate. Lead-zinc estimated to be 20% combined. Some chalcopyrite. Some of the galena quite coarse grained.	4240		162-167		2.16	0.28	3.52	6.48	2.36	0.13	6.70	5.41
			4241		167.0-170.5		2.40	0.28	7.58	5.88				

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van 2.5

PROPERTY PANGORDA

SHEET NUMBER 7 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Slings TO Sample	Footage WIDTH	CORE SAMPLE			SLUDGE SAMPLE				
							Ag %	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn
169-174.5	5.0	Massive sulphides <u>167.0-170.5</u> Same as section 162.0-169.0 above.												
		<u>170.5-173.0</u> Pyrite & white quartz carbonate replacing quartz. Some galena & sphalerite.	4242	170.5-174.5	4320	170-175 0.76 26.0	0.43	0.88	1.20	0.90	0.23	1.38	1.87	
		<u>173.0-174.5</u> Quartzite 70% replaced by pyrite. Galena & sphalerite recognisable locally.												
174.5-182.0	7.5	Massive sulphides <u>174.5-182.0</u> Quartzite 80% replaced by massive brassy pyrite. No recognisable galena or sphalerite. A few scattered grains chalcopyrite	4243	174.5-179.5	4321	175-180 0.84 28.8	0.50	0.50	0.90	0.86	0.30	0.80	1.08	
			4244	179.5-184.5	4322	180-185 0.50 17.1	0.68	0.28	0.60	0.70	0.40	0.50	0.49	
182-192	10.0	Massive sulphides <u>182.0-185.0</u> Same as section 174.5 to 182.0, above. A few scattered grains galena & sphalerite	4245	184.5-189.0	4323	185-190 0.64 21.9	0.70	0.47	0.90	0.80	0.48	0.75	0.79	
		<u>185.0-191.9</u> streaky pyrite - the streaks due to quartz with galena & sphalerite. Some chalcopyrite	4246	189.0-191.9		0.54	0.53	0.75	1.10					
		<u>191.9-192.0</u> Same as section 192.0 to 192.8 below.				18.5								

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 7-15

PROPERTY TANGORDA

SHEET NUMBER 8 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core Sample NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
							Ag g/t	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn
192-199	6.0	192.0-192.8 Massive pyrite with stringers - massive sphalerite & galena. Lead-zinc estimated to be 20% combined. About 1% chalcopyrite. 192.8-193.7 White quartz vein with a few grains of chalcopyrite and pyrrhotite.	4247	191.9-192.8	4324	190-195	1.92 6.8	0.48	2.90	4.48	1.26	0.25	1.87	1.97
		193.7-199.0 Sericite schist. Highly contorted. Some banding regular at 45° to core.			4325	195-200				0.34	0.20	0.22	0.49	
199.0-205.0	4.0	Sericite schist - banding highly contorted. Injected with quartz in blebs & veinlets. Brown pyrrhotite present in scattered grains, blebs and veinlets.												
205-211	4.0	Sericite schist - locally silicified Section 205.9 to 206.5 contains resinous sphalerite with some galena associated with brown pyrrhotite in stringers & blebs. Lead-zinc estimated to be 2% combined. Pyrrhotite common throughout the entire section.	4462 n.a.	205.9-206.5										
211-212	0.6	Sericite schist with a 1" section brown pyrrhotite.												
212-216	4.0	Sericite schist altered to light colored talc. Locally silicified and cut by white quartz stringers. Pyrrhotite scattered throughout forming approximately 1% of the volume.	4463 n.a.	212-213.2										

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Yan 1-5

PROPERTY YANGORDA

SHEET NUMBER 9 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag %	Cu %	Pb %	Zn %		
		<u>212.0-213.2</u> Fine galena throughout the core. Some sphalerite, chalcopryrite & pyrite. Mineral less than 3% of volume.										
216-220	3.5	Sericite schist, locally altered to talc. Banding locally regular at 45° to core. Barren white quartz vein from 216.0 - 217.2. Some pyrrhotite in blebs and small stringers.										
220-221	1.0	Talc-sericite schist with small quartz stringers.										
221-226	3.5	Sericite schist altered to talc. Original banding still present & highly drag-folded. Section 10% to 15% sulphides, mostly pyrrhotite but approximately 2% galena with some sphalerite and chalcopryrite. Mineral in blebs & stringers in highly folded sections. Some pyrrhotite stringers are drag-folded.	4268			0.16	0.38	0.41	0.10			
						55						
226-231	3.7	<u>226.0-228.0</u> Same as section 221.0 to 226.0 above. <u>228.0-231.0</u> Sericite schist altered to talc. 5% sulphides consisting of brown pyrrhotite with some sphalerite and pyrite.	4269	226-228		0.44	0.30	0.91	0.30			
			4464 M.C.	228-231		15.1						

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van 7-5

PROPERTY YANGORDA

SHEET NUMBER 10 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage		CORE SAMPLE				SLUDGE SAMPLE				
				FROM	TO	WIDTH								
231-236	4.0	Sericite schist altered to talc. Banding contorted & drag folded. generally but some regular at 54° to core. Section 7% sulphides in stringers & blebs. Some stringers are drag folded. Mineral pyrrhotite with minor amounts of chalcopyrite.	4465 N.A.	231-232										
236-241	4.5	Sericite schist altered to talc. Contorted & silicified. Approximately 5% sulphides throughout - pyrrhotite with some chalcopyrite except for section 231.5-231.8. This section 20% sulphides in schist & white quartz. - pyrrhotite, galena, sphalerite & chalcopyrite. Lead-zinc approximately 8% combined.	4466 N.A.	236.5-237.5										
241-249	7.0	Sericite schist altered to talc. 241.0-246.0 Silicified & 15% replaced by brown pyrrhotite in stringers & blebs. Mineral & banding both drag-folded. 246.0-249.0 Same as above except for white quartz stringers and the presence of 2% to 3% galena & sphalerite. The latter are concentrated in and near the quartz. Some chalcopyrite.	4467 N.A.	246-249										

DIAMOND DRILL RECORD,

HOLE NO. 10-15

PROPERTY VANGORDA

SHEET NUMBER 11 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE		SLUDGE SAMPLE	
				FROM	TO	WIDTH				
260-254	5.0	Talc-sericite schist. Bedding drag-folded & contorted but locally regular at 60° to core. Section 10% replaced by brown pyrrhotite in stringers with quartz. Five white quartz stringers with scattered small grains galena. Some galena in the pyrrhotite - some chalcopyrite	4468 n.a.	249-254						
254-260	5.0	Talc sericite schist - some sections pure talc. Bedding regular at 45° to core. Section 7% replaced by brown pyrrhotite. Some chalcopyrite.								
260-264	2.1	Talc - 10% replaced by brown pyrrhotite. Pyrite & minor amounts galena.	4469 n.a.	260-264						
264-270	4.0	Schist altered to talc, silicified & replaced by white quartz carbonate. A fracture & the carbonate stringer lie along the core. 5% mineral made up of 3% pyrrhotite, 1% galena, 1% sphalerite, pyrite & chalcopyrite.								
270-274	2.0	Same as above except mineral all pyrrhotite.								
274.0-280.0	4.6	Talc sericite schist. Bedding regular at 55° to core. Quartz carbonate stringers along core. Pyrrhotite - some sphalerite and galena related to the stringer.	4470 n.c.	274-276.3						

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Vm 45

PROPERTY VANGORDA

SHEET NUMBER 12 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
269-297	15.0	Talc sericite schist. Banding generally regular at 50° to 55° to core. Scattered quartz stringers. 7% sulphides composed of pyrrhotite with 3% chalcopyrite.												
297-305	6.7	Same as above												

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 7m 5

PROPERTY VANCOEDA

SHEET NUMBER 1 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag/yt	Cu%	Ni%	Pb	zn	
305-311	5.5	Sericite schist partially altered to talc. Bedding contorted. Deep brown pyrrhotite in stringers and blebs										
311-316	4.6	Talc with remnants of unaltered schist. Banding locally regular at 50° to core.										
316-320	4.0	Same as above. Minor amounts brown pyrrhotite.										
320-323	3.0	Same as above.										
323-330	7.0	Sericite schist highly altered to talc. Banding highly contorted & drag folded. 2% brown pyrrhotite in stringers and blebs. Many of these stringers drag-folded with the banding.										
330-334	4.0	Sericite schist altered to talc. Banding regular at 70° to 80° to core. White quartz stringers comprise 10% of section. Brown pyrrhotite in stringers and blebs along banding.										
334-344	9.8	334-335.8 Sericite schist partially altered to talc. Banding regular at 70° to core. Section 35% replaced by brown pyrrhotite in stringers along and across banding. 1% to 2% chalcopryrite. 335.8-339.4 Talc sericite schist. Banding regular at 70° to core. One quartz stringer with brown pyrrhotite.	4250 Ni/Cu	334.0 335.8	4326 n.c.	335 340			Cu Ni 0.8 nil			

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 5

PROPERTY YANGORDA

SHEET NUMBER 14 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE				SLUDGE SAMPLE				
				FROM	TO	WIDTH	Ag	Cu	Ni	Pb	Zn				
		<u>339.6-346.0</u> Section altered to talc & carbonate. Cut by white carbonate stringers. Sulphides comprise 35% of section and increase in amount towards the end. Consist of pyrite & pyrrhotite with 1% chalcopyrite. Reddish sphalerite noted in first few inches. Pyrrhotite common at beginning of section gives way to pyrite at end.	4351	339.6	4327	340-345	0.62	0.23	-	0.83	1.47				
							21.3								
346-347	3.0	25% sulphides replacing quartzite or silicified schist. Banding in schist obscure. Sulphides 15% pyrite, 7% sphalerite, 2% chalcopyrite, 1% galena.	4352 +Ni		4328	345-350	0.66	0.20	-	0.66	0.20				
							22.6								
347-358	10.6	347.0-349.6 25% sulphides replacing silicified schist. Sulphides 17% pyrite, 5% sphalerite, 2% chalcopyrite, 1% galena.	4353 +Ni	347-349.9	4329	350-355	0.14	0.23	Ni	0.41	0.59				
							4.8								
		349.4-349.9 Schist altered to talc. 3% pyrrhotite.													
		349.9-353.0 40% sulphides replacing silicified talc. Sulphides 15% pyrrhotite, 12% pyrite, 7% sphalerite, 4% galena, 2% chalcopyrite. Galena may be more abundant but occurs in such small grains it is difficult to estimate.	4354 +Ni	349.9-353			1.04	0.28	Ni	1.96	4.59				
							35.6								

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 5

PROPERTY VANCOEVA

SHEET NUMBER 15 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
		<u>353.0-358.0 Schist altered to talc.</u>								
		<u>Banding regular at 70° to core flattening to 55° at end of section. Several quartz stringers 2% brown pyrrhotite.</u>								
<u>358-364</u>	<u>6.0</u>	<u>Sericite schist partially altered to talc. 6" quartz vein at beginning of section. 4% pyrrhotite with some chalcopyrite. Banding regular at 55° to core.</u>								
<u>364-372</u>	<u>8.0</u>	<u>Sericite schist. Banding locally regular at 55° to core locally contorted. White quartz in stringers & blebs comprises 5% of section. 1% to 2% brown pyrrhotite in blebs and narrow stringers.</u>								
<u>372.0</u>		<u>END OF HOLE.</u>								

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 6

PROPERTY VANGORDA

SHEET NUMBER 1

SECTION FROM _____ TO _____

STARTED Oct 16, 1953

LATITUDE 30.088.73 9171.04N

DATUM _____

COMPLETED Oct 19, 1953

DEPARTURE 29.951.06 9129.08E

BEARING _____

ULTIMATE DEPTH 1171.0 (35.66)
1234.97 m

ELEVATION 4021.29 1304.63 m

DIP -00°

PROPOSED DEPTH _____

Checked
Hayes
April 15, 1957

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
							Ag %	Cu %	Pb %	Zn %	Ag	Cu	Pb	Zn
0.0-44.5	134 ^m	Casing - large granite boulders in overburden			4176	41-44					0.88	0.28	2.64	0.50
44-46	1.8	Massive sulphides with 15% quartz. Pyrite with galena & sphalerite in streaks and blebs. Galena & sphalerite approximately 7% combined	4230		4177	44-46	2.84	0.35	6.48	6.38				
46-47	1.0	Massive sulphides - pyrite with galena & sphalerite in sections of unreplaced quartzite. Some relatively coarse galena	4231		n.a.		3.40	0.28	9.56	8.37				
47-49	1.1	Massive sulphides. Pyrite, galena and sphalerite. Lead and zinc estimated to be 15% combined. Some quartzite partially replaced by pyrite at end of section	4232		4178	46-49	3.16	0.43	8.02	13.55				
49-51	0.9	Blue white quartzite shattered and replaced by 10% to 15% brassy pyrite. Scattered blebs of sphalerite & galena	4233		4179	49-51	0.80	0.43	2.27	3.09	0.56	0.43	1.21	2.29
51-60		Ground core - sludge grey sericite schist.			4180	51-53					0.16	0.23	0.06	0.70
					4181	53-55					Tr.	0.15	Tr.	1.01
					4182	55-57					0.16	0.10	Nil	Trace
					4183	57-60					0.18	0.20	Nil	Nil
60-62	0.4	Chips recovered are all white quartz chips with some brassy pyrite. Sludge indicates the presence of schist.			4184	60-62					0.30	0.18	Tr.	Nil
		Quartz is the vein type rather than quartzite.												

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 6

PROPERTY VANGORDA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
							Ag/gt	Cu%	Pb%	Zn%	Ag	Cu	Pb	Zn
62-64	0.4 chips	Same as above	4234	60-64	4185	62-64	0.22	0.18	0.11	nil	0.26	0.20	Nil	0.10
64-68	1.3	Black graphite schist with quartz at the beginning of the section - Banding highly contorted			4186 4187	64-66 66-68					0.28 0.22	0.18 0.23	Nil Nil	0.50 0.20
68-76		Ground core - sludge dark schist			4188 4189	68-73 73-78					0.34 0.04	0.30 0.25	Nil Nil	0.30 0.10
76-78	1.3	Graphitic schist - extremely contorted and drag folded. Brecciated. Some brassy pyrite in carbonate veinlets & blebs.												
78-81	0.5 chips	Same as above			4190	78-83					0.18	0.15	Nil	Nil
81-83	0.1 chips	Same as above												
83-85	0.1 chips	Two chips core recovered are white quartz with a few grains pyrite. Sludge largely schist.												
85-87		Ground core - sludge schist.												
87-89	0.1 chips	Graphitic schist												
89-93	0.4 chips	Same as above												
93-95	0.4 chips	Graphitic schist - shearing contorted. Some pyrite in quartz-carbonate veinlets.												
95-98	0.4 chips	Graphitic schist. pyrite in a quartz stringer.												
98-100	0.1 chips	Graphitic schist												
100-109	0.2 chips	Graphitic schist & quartz												
119-111	0.5 chips	Same as above												

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 6

PROPERTY VANGORDA

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
111-113	0.8	Graphitic schist. Banding regular but core too broken up to determine angle.													
113-116	1.0	Graphitic schist. Some white quartz. Banding regular at 40° to core.													
116-117	0.7	Graphitic schist. Banding regular at 47° to core. Schist harder and more slaty in appearance. Tiny quartz veinlets along the bedding containing pyrite and what may be a few grains of sphalerite.													
117.0		END OF HOLE.													

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 7

PROPERTY YANCOODA

SHEET NUMBER 1 SECTION FROM _____ TO _____ STARTED Oct 22, 1953
 LATITUDE 29936.80 9124.73N DATUM _____ COMPLETED _____
 DEPARTURE 30081.97 9168.98E BEARING _____ ULTIMATE DEPTH _____
 ELEVATION 3991.86 +259' 1295.6 DIP 99° PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE					
							Ag	Cu	Pb	Zn	Ag	Cu	Pb	Zn		
0.0-53.0	16.1'	<u>Quartzite</u>														
53-55		<u>GROUND CORE - Sludge pyrite in quartzite</u>			4331	53-55						0.70	0.10	0.17	Nil	
55-57	0.2 chips	<u>Chips recovered one quartzite 50% replaced by pyrite.</u>			4332	55-57						0.62	0.10	0.17	Nil	
57-59	0.9	<u>Quartzite 40% to 50% replaced by brassy pyrite. Mineral tends to occur in massive beds.</u>	4355 ✓		4353	57-59	0.76	0.13	0.25	Nil		0.98	0.10	0.39	Nil	
59-63	0.9	<u>Same as above</u>	4356 ✓		4334	59-61						0.50	0.13	0.19	Nil	
63-65	0.8	<u>Quartzite replaced by sphalerite, galena & pyrite. Lead-zinc possibly 8% to 10% combined</u>	4357 ✓	2	4335	61-63	0.68	0.13	0.55	Nil		0.90	0.15	1.16	Nil	
65-66	0.6	<u>Quartzite 30% replaced by pyrite with some galena & sphalerite. Lead-zinc possibly 5% combined.</u>	4358 ✓	1	4336	63-65	8.36	0.55	7.57	10.49		1.98	0.40	3.73	4.42	
66-67	0.5	<u>Quartzite 30% replaced by pyrite with some galena & sphalerite. Lead-zinc possibly 5% combined</u>	4359 ✓	1		65-67	1.66	0.45	3.30	4.00		1.28	0.35	2.42	3.24	
67-70	0.9	<u>Same as above</u>	4360 ✓	3			1.18	0.15	2.19	3.43						
70-71	1.0	<u>Quartzite 35% replaced by pyrite - some in massive stringers. Scattered grains galena & sphalerite.</u>	4361 ✓	1	4338	67-69	0.76	0.50	1.38	1.67		1.40	0.43	2.31	2.75	
71-72	0.5	<u>Quartzite 30% replaced by pyrite</u>	4362 ✓	2	4339	69-71	0.80	0.53	1.22	2.35		0.94	0.58	1.44	2.16	
72-73	0.5	<u>Quartzite 30% replaced by pyrite with some galena & sphalerite. Lead-zinc possibly 5% combined.</u>			4340	71-73	1.20	1.46	1.32	5.38		1.06	0.75	1.21	3.24	

DIAMOND DRILL RECORD,

HOLE NO. Ym # 7

PROPERTY YANGORDA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE				SLUDGE SAMPLE			
				FROM	TO	WIDTH	Ag	Cu	Pb	Zn	Ag	Cu	Pb	Zn
73-76	1.7	Quartzite 30% replaced by pyrite, sphalerite & galena. Lead-zinc 5% to 7% combined.	4363 ✓	3 ⁹⁰	4341	73-75	1.10	0.53	1.66	3.62	1.06	0.80	1.30	2.65
					4342	75-77					1.06	0.78	1.57	2.95
76-81	2.5	Quartzite 30% replaced by pyrite, - galena & sphalerite common in last foot of section.	4364 ✓	5	4343	77-81	0.60	0.70	0.80	1.96	0.96	1.00	0.83	1.78
81-84	2.0	Quartzite with black schist beds 20% replaced by pyrite, galena & sphalerite. Lead-zinc possibly 5% combined. Bedding 10° to core.	4365 ✓	3	4344	81-84	0.96	0.30	2.02	3.23	0.74	0.58	1.35	3.05
84-85	0.3	Quartzite & black schist 20% replaced by pyrite, galena & sphalerite. Lead-zinc 10% to 12% combined. Bedding contorted but at a high angle to core	4366 ✓	2	4345	84-86	1.90	0.23	4.72	5.28	1.36	0.40	3.06	3.93
85-86	1.0	Same as above												
86-90	1.2	Same as above	4367 ✓	4	4346	86-90	1.66	0.30	4.61	3.52	3.26	0.48	3.26	4.03
90-92	1.2	Quartzite & schist 20% replaced by sphalerite, galena & pyrite. Lead-zinc possibly 10% combined.	4368 ✓	2	4347	90-92	1.16	0.13	3.08	4.90	1.56	0.78	2.95	5.50
92-94	0.6	Quartzite & schist 20% replaced by galena, sphalerite & pyrite. Lead-zinc possibly 7% combined.	4369 ✓	-	4348	92-94	1.34	0.15	3.84	4.12	1.36	0.13	3.15	5.50
94-96	0.9	Quartzite 75% replaced by pyrite with some galena & sphalerite. Lead-zinc 8% combined	4370 ✓	2	4349	94-96	1.92	0.20	4.06	6.66	2.04	0.28	2.43	6.39
96-99	1.0	Massive pyrite & galena. Galena occurs in very fine grains throughout making an estimate impossible. Possibly 15%.	4371 ✓	3	4350	96-99	1.96	0.38	6.38	9.80	3.44	0.35	10.10	10.62

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 7

PROPERTY YANGORDA

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE				SLUDGE SAMPLE			
				FROM	TO	WIDTH	Ag	Cu	Pb	Zn	Ag	Cu	Pb	Zn
99-101	2.0	Same as above - a few grains of sphalerite.	4372 ✓	4401	99-101	3.64	0.38	5.71	9.30	2.52	0.18	2.68	5.01	
101-103	1.3	Same as above. Sphalerite a little more abundant.	4373 ✓	4402	101-103	2.84	0.30	4.61	7.65	1.98	0.48	2.76	4.33	
103-104	0.9	Quartzite & schist 10% replaced by pyrite, galena and sphalerite.	4374 ✓	4403	103-105	1.76	0.05	0.76	0.49	1.34	0.28	1.82	2.65	
104-108	3.2	Interbedded quartz & schist 5% to 10% replaced by pyrite. No recognizable galena or sphalerite. Bedding extremely contorted.	4375 ✓	4404	105-107	0.38	0.15	0.83	nil	1.16	0.35	1.82	1.38	
108-111	1.4	Section same as above except for last 6 inches which is massive pyrite.	4376 ✓	4406	109-111	0.90	0.10	1.88	2.36	1.54	0.33	1.52	1.38	
111-112	0.3	Massive pyrite & galena	4377 ✓	4407	111-113	1.90	0.23	3.51	6.10	1.40	0.30	2.21	2.65	
112-113	0.9	Massive pyrite and galena. Estimate of galena content impossible to fine-grained occurrence but is abundant. Possibly 15%												
113-116	3.0	Quartzite 80% replaced by brassy pyrite. 8% galena & 5% sphalerite in streaks through the pyrite.	4383 ✓	4408	113-115	3.70	0.48	5.22	9.04	1.96	0.33	3.12	3.83	
116-117	0.3	Massive pyrite with some galena chips		4409	115-117					2.62	0.43	3.92	6.49	
117-119	1.4	Massive pyrite with 20% quartz, 10% to 12% galena & sphalerite combined.	4384 ✓	4410	117-119	2.92	0.38	4.08	8.75	2.58	0.40	3.64	5.70	
119-127		GROUND CORE - Sludge white talc schist. Some sulphides due to caving.		4411	119-127					1.56	0.98	4.20	4.78	

DIAMOND DRILL RECORD,

HOLE NO. Van # 7

PROPERTY VANGUARD

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
							Ag	Cu	Pb	Zn				
127-130	0.5 chips	Quartz stringers in talc sericite schist.		3 ³⁴										
130-133		GROUND CORE - Sludge white sericite schist.		3										
133-138	1.8	Section 60% sulphides and 40% white quartz. Sulphides 40% pyrite & 20% combined galena & sphalerite	4385 ✓	5			3.18	0.15	5.69	10.81				
138-142	4.0	80% sulphides & 20% quartz. Sulphides 70% pyrite & 10% galena & sphalerite combined. Latter confined generally to sections containing quartz and narrow quartz carbonate veinlets cutting the pyrite.	4386 ✓	4			1.44	0.15	2.96	7.67				
142-147	4.7	80% massive sulphides & 20% quartz. Sulphides 60% pyrite, 12% galena, 8% sphalerite.	4387 ✓	5			1.46	0.50	2.98	3.24				
147-152	1.7	147.0-148.0 Massive sulphides same as section 142.0-147.0 above. 148.0-152.0 Talc sericite schist.	4388 ✓	147 148	0 0		1.56	0.15	3.48	6.49				149'
152-157	3.8	Contorted sericite schist altered to talc. Cut by scattered white quartz stringers. Two narrow stringers contain galena, sphalerite, pyrite and pyrrhotite.		5	0 0				Nic	Nic				
157-161	3.5	Massive sulphides with 10% quartz. The first foot of the section is pyrrhotite with some pyrite, galena, & sphalerite - the remainder massive pyrite with some galena & sphalerite. Galena & sphalerite 5% combined.	4389 ✓	4			1.14	0.40	2.51	2.85				

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Ym # 7

PROPERTY YANGORRA

SHEET NUMBER 5 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE				SLUDGE SAMPLE						
				FROM	TO	WIDTH	Ag	Cu	Pb	Zn							
161-165	0.9	The chips recovered are white quartz and brassy pyrite. Sludge is largely tale-sericite schist.															
165-167	1.2	Sericite schist altered to tale and 50% replaced by white quartz veins.															
167-168	0.8	Highly contorted sericite schist altered to tale and white carbonate.															
168-172	0.3 chips	Chips recovered are white quartz mineralized with small amounts of pyrite, galena & sphalerite. Sludge is tale sericite schist.															
172-176	2.0	First half of section is white tale sericite schist; - the last half dark sericite schist. All 10% replaced by brown pyrrhotite in small stringers & blebs. 1% chalcocopyrite.															
176-181	4.3	Medium-grey sericite schist highly altered to tale. Banding obscure, 1% replaced by brown pyrrhotite. 1% to 2% chalcocopyrite. Relatively coarse galena & sphalerite in schist at 180.0. Some galena in a white quartz stringer at end of the section.	4390 ✓				0.18	0.15	0.25	0.39							
181-187	5.6	181.0-181.5 White quartz vein mineralized with pyrite, pyrrhotite & chalcocopyrite. 181.5-187.0 Highly altered dark grey green schist. 20% replaced by brown pyrrhotite. 1% to 2% chalcocopyrite. Local occurrences of galena & sphalerite.	4391 ✓				Tr.	0.15	0.14	0.39							

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 7.

PROPERTY YANGORBA

SHEET NUMBER 6 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage		CORE SAMPLE				SLUDGE SAMPLE		
				FROM	TO	WIDTH	Ag	Cu	Pb	Zn		
187-192	4.7	Sericite schist highly altered and silicified. 20% replaced by pyrrhotite. 1% to 2% chalcopyrite. One splash sphalerite.										
192-197	4.8	192.0-192.5 Talc sericite schist. 192.5-193.0 White quartz. Coarse galena in fractures. 193.0-194.1 Quartz 30% replaced by sphalerite, galena and some pyrite. 194.1-195.8 Massive pyrite with 5% to 8% combined galena & sphalerite. Heavy chalcopyrite in one remnant of quartz and talc. 195.8-197.0 Section 60% replaced by sphalerite, pyrite & galena. Galena-sphalerite possibly 30% combined.	4392 ✓	192.5-197.5	1.58	0.23	0.25	4.23				
197-202		197.0-197.5 Schist 30% replaced by massive sphalerite & pyrite. Some galena. 197.5-202.0 Highly contorted black graphitic schist. Injected with white carbonate veinlets. Some pyrrhotite. Schist altered to talc. A 1" stringer of massive sphalerite, pyrite & galena at 202.3.										
202-204	1.1											
204-208	2.2	Sericite schist altered to talc.										
208-210	1.7	Talc sericite schist with white quartz stringers.										

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van 1-7

PROPERTY YAKORBA

SHEET NUMBER 7 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag	Cu	Pb	Zn		
210-213	0.5	Same as above										
213-216	1.5	Talc sericite schist. 1 st sulphides at end of section										
216-221	5.0	Massive sulphides with 10% to 15% quartz. Sulphides 15% sphalerite, 10% galena, the remainder pyrite. Banding in sulphides evident.	4456 ✓			5	1.70	0.28	4.48	5.60		
221-227	5.0	Sphalerite rich sections prominent. 221.0-221.8 Sulphides 60% replaced by white carbonate. 221.8-224.8 Massive pyrite with 15% combined sphalerite & galena. 224.8-226.5 Talc sericite schist	4457 ✓	221.0-224.8		3.8	2.48	0.25	3.59	7.57		
227-232	3.0	226.5-227.0 Quartz 70% replaced by pyrite, sphalerite & galena. 227.0-227.2 Sulphides same as section 226.5-227.0 above. 227.2-232.0 Schist altered to talc. Locally silicified and cut by quartz stringers. Scattered brassy pyrite.	4458 ✓	227.0-227.3		2.5	1.36	0.20	2.32	3.93		
232-236	1.5	Talc sericite schist.										
236-238	2.0	Talc sericite schist.										
238-243	1.7	Talc sericite schist. Scattered pyrite										
243-246	2.2	Talc sericite schist. 3% combined pyrite & pyrrhotite.										
246-251	4.2	Talc sericite schist. Shearing contorted but generally at a low angle to the core. Scattered quartz stringers with brassy pyrite.										

DIAMOND DRILL RECORD,

HOLE NO. 7

PROPERTY VANGORDA

SHEET NUMBER 8 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	CORE SAMPLE NO.	Footage		CORE SAMPLE				SLUDGE SAMPLE	
				FROM	TO Sample	Ag	Cu	Pb	Zn		
252-254	1.2	Same as section above.									
254-259	5.0	254-254.3 Sericite schist 254.3-254.6 Quartz stringer with galena, chalcopyrite, sphalerite and pyrite. 254.4-254.9 Quartz & schist, some interbedded. Replaced by 15% pyrite. A few grains of sphalerite. 254.9-259.0 Section 70% sulphides 30% quartz and white carbonate. Sulphides pyrite with 5% pyrrhotite and 5% combined galena & sphalerite. A few grains of chalcopyrite.	4459 ✓	254.3-259	0.88	0.43	1.24	1.57			
259-263	4.0	Interbedded quartz & schist. Thin quartz veinlets cut the schist bands indicating silicification. Bedding regular at 20° to core. Section 15% replaced by brassy pyrite which selectively replaces the quartz. A few grains sphalerite.	4460 ✓		0.46	0.15	0.88	Tr.			
263-268	3.1	263.0-264.6 Quartz & white carbonate 60% replaced by brassy pyrite. 4% sphalerite and galena. 264.6-265.4 Interbedded schist and quartz 10% replaced by pyrite. 265.4-268.0 Black graphitic schist.	4461 ✓	263.0-264.6	0.76	0.20	1.27	1.97			
268-273	1.3	Black graphitic schist. Minor amounts of pyrite.									

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 7

PROPERTY VANGORBA

SHEET NUMBER 9 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage			CORE SAMPLE				SLUDGE SAMPLE						
				FROM	TO	WIDTH											
273-276	0.9	Same as above. Shearing contorted.															
276-277	0.4	Black graphitic schist.															
277-281	1.8	Same. Highly contorted. Some pyrite.															
281-286	0.1	Black graphitic schist.															
286-288	0.1	" " "															
288-291	0.1	" " "															
291-296	1.4	Highly contorted black graphitic schist.															
296-300	3.9	Same. Scattered pyrite.															
300		<u>END OF HOLE</u>															

DIAMOND DRILL RECORD,

HOLE NO. Van # 8

PROPERTY YANGORDA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core Sample NO.	Footage FROM	Sludge Sample TO	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag	Cu	Pb	Zn		
111-113	2.0	Same as above. 35% sulphides as follows - 15% pyrite, 8% sphalerite, 8% galena, 4% chalcopyrite.	4379	111-113	2'	2	1.36 46.6	0.20	3.70	4.13		
113-124	10.5	113.0-117.5 Schist and quartz 35% replaced by sulphides - 15% sphalerite, 8% pyrite, 5% galena, 4% pyrrhotite, 3% chalcopyrite. 117.5-124.0 Sericite schist altered to talc. The contact between the quartz-schist section and the sericite schist is abrupt but not sharp. Shearing at 40° to 50° to core.	4380	113.0-117.5	4.5'	4.5	1.18 40.4	0.13	2.54	5.21		
124-136	10.0	Talc sericite schist. Shearing regular but varies from 40° to 50° to core. Scattered quartz stringers. Two stringers pyrrhotite.										
136-145	7.3	Sericite schist. Shearing 45° to core. 142.5-145.0 Sulphides present as small replacement stringers and blebs along bedding. Sphalerite, pyrrhotite and galena.	4381	142.5-145.0			0.16 5.5	0.10	0.52	4.62		
145-153	3.5	145.0-146.0 Sphalerite, galena, pyrrhotite & chalcopyrite comprise 20% of section. 146.0-153.0 Sericite schist.										
153-159	5.5	Sericite schist. Shearing 20° to core 10% pyrrhotite in massive stringers and finely disseminated grains. 1% chalcopyrite.										

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. Ym 18

PROPERTY VANGORDA

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage Sludge Footage			CORE SAMPLE		SLUDGE SAMPLE	
				FROM	TO	WIDTH				
159-167	7.0	Talc sericite schist. Shearing regular but changes from 70° to core at beginning to 50° at the end. 10% pyrrhotite in massive stringers. Sphalerite & pyrite with the pyrrhotite at 166.0.								
167-175	8.0	Sericite schist altered to talc in sections. Shearing contorted but locally regular at 55° to core. 5% pyrrhotite in stringers. Pyrite with 1% to 2% sphalerite from 166.0-169.0.								
175-180	5.0	Sericite schist. Shearing 50° to 55° to core. Section 177.0 to 178.5 20% replaced by pyrrhotite with minor amounts of sphalerite and chalcopyrite.								
180-189	9.0	Sericite schist. Shearing 45° to 50° to core. 15% replaced by pyrite & pyrrhotite in stringers. Scattered sphalerite & galena comprising less than 1% of the section.								
189-200	11.0	Talc sericite schist. Shearing regular at 50° to 55° to core. Section 5% replaced by pyrite & pyrrhotite. A few tiny veinlets of sphalerite.								
200-203	2.3	Sericite schist.								

DIAMOND DRILL RECORD,

HOLE NO. Van 1 6

PROPERTY VANGORDA

SHEET NUMBER 6 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage FROM	Sludge TO Sample	Footage WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag	Cu	Pb	Zn		
278-285	7.3	278.0-280.8 Schist & quartz highly contorted and 10% replaced by pyrite. A few scattered grains chalcopyrite & sphalerite.	4397 ✓	278.0-280.8		2.8	0.32	0.23	0.53	0.20	4C	
		280.8-285.0 Quartz and white carbonate 75% replaced by brassy pyrite. 1% to 2% combined galena & sphalerite. Some chalcopyrite. Remnants of talc schist in the section.	4398 ✓	280.8-285.4		4.6	0.92	0.25	1.27	0.96	4EC	
285-295	5.3	285.0-285.4 Same as section 280.8-285.0 above.	4399 ✓	285.4-290.0								
		285.4-295.0 Graphitic schist and quartz in thin beds. Quartz partially replaced by pyrite and some sphalerite	4400 ✓	290.0-295.0								
295-297	1.5	Black graphitic sericite schist with white quartz veins.										
297-304	3.9	First half of section well sheared talc sericite schist - last half graphitic schist. Shearing at 30° to core. Sparsely mineralized with pyrite.										
304-312	6.1	Black graphitic schist changing to interbedded schist and quartz from 307.0 to 310.0. Quartz partially replaced by brassy pyrite and pyrrhotite. A few grains of sphalerite in one section.										

DRILLED BY _____

4399/4400 held back

quartz. Less than 1% & sphalerite.

DRILLED BY _____

DIAMOND DRILL RECORD,

HOLE NO. Van # 8

PROPERTY VANGORDA

SHEET NUMBER 8 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Core Recovery	FORMATION	Core SAMPLE NO.	Footage			Sludge Footage				CORE SAMPLE				SLUDGE SAMPLE					
				FROM	TO	WIDTH	Ag	Co	Pb	Zn										
332-344	7.8	332.0-336.5 Black schist 90% replaced by equal amounts of quartz & pyrite. A few scattered grains of galena & sphalerite.	1154 ✓	332.0			0.64	0.20	1.58	1.77										
		336.5-338.0 Black graphitic schist with 10% pyrite	1155 ✓	336.0			21.9													
		338.0-339.0 Massive pyrite same as section 332.0 to 336.5 above.		340.0			0.40	0.18	0.33	0.98										
		339.0-344.0 Black graphitic schist. First foot of section mineralized with pyrite.					13.7													
344-353	4.5	Contorted and altered black graphitic schist. Minor amounts of pyrite.																		
353-360	2.5	Contorted black graphitic schist. Some pyrite in small veinlets.																		
360.0		END OF HOLE.																		

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Yangoria

SHEET NUMBER 2

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
						Ag.	Cu	Pb	Zn		
	<i>Recovery</i>										
73-78	5.0 80% massive fine pyrite replacing sericite schist. Between 5 and 10% fine Pbs & Zns.	4473	73	78	5.0	76.1 2.22	0.15	3.96	6.19		
78-81	2.1 85% massive pyrite. 5 to 10% fine Pbs & Zns.	4474	78	81	3.0	76.8 2.24	0.20	5.94	4.82		
81-85	2.5 70-75% massive pyrite. About 5% fine Pbs & Zns	4475	81	85	4.0	1.96 67.2	0.18	3.41	7.08		
85-88	2.2 70% massive pyrite. A few threads of fine sphalerite and galena.	4476	85	88	3.0	0.96 12.9	0.33	1.32	2.06		
88-93	4.0 70 to 80% massive pyrite. Less than 10% Pbs & Zns.	4477	88	93	5.0	1.54 52.8	0.28	1.57	2.36		
93-94	0.9 As above	4478	93	94	1.0	0.76	0.25	0.77	1.67		
94-97	1.7 Sharp change at 94 to 20% pyrite, 30% Zns and 5% Pbs. Congue consists of graphitic schist mottled with greyish-white chalky alteration (not carbonate)	4479	94	97	3	2.14	0.15	4.66	9.44		
97-100.5	2.6 Same mineralization as above	4480	97	100.5	3.5	1.50	0.18	2.70	5.80		
100.5-104	3.0 Same as above: slightly less white chalky mottling. Schist less graphitic - dark grey colour, contorted.	4481	100.5	104	3.5	2.18	0.18	4.73	12.68		
104-111	2.9 Greenish-grey sericite schist Graphitic (0.5) at 105.										

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Yangorda

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	RECOVERY	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag	Cu	Pb	Zn		
111-115	2.4	Pale grey sericite schist containing irregular streaks of pale green material. Schistosity contorted.										
		114-115 0 1/4 massive pyrite fragments from 114.6 to 115.	4482	114	115	1	5.5 0.16	0.18	0.17	0.69		
115-120	1.2	0 1/4 containing 30% pyrite in sericite schist. Sparse sphalerite	4483	115	120	5	-	0.15	-	-		
120-125	1.5	Greenish-grey sericite schist containing about 30% white Qtz. replacement.										
125-128	1.1	Fragments of greenish-grey sericite schist.										
128-131	1.3	Medium grey sericite schist containing about 30% vuggy qtz. replacement										
131-134	1.4	As above. About 20% white qtz. fragments.										
134-136	1.1	As above, grading to light grey sericite schist at 136.										
136-140	3.3	Sharp contact at 136, to 25% pyrite and 25% Pbs and about 5% fine Zns.	4484	136	140	4	52.8 1.54	0.18	2.98	7.27		
140-144	3.6	50% py. 40% Pbs & Zns. (latter in minority) Massive appearance	4485	140	144	4	83.6 2.66	0.10	3.52	8.75		
144-147	2.9	80% massive pyrite, about 5% Pbs, sparse Zns. Gangue - threads of qtz. (bluish)	4486	144	147	3	2.68 85.0	0.28	6.82	4.03		

11-11.17
 8'-3.9

DRILLED BY Boyles Bros. Ltd.

SIGNED D. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
						Ag	Cu	Pb	Zn		
	<i>Recovery</i>										
147-155	6.3 Massive pyrite as above, estimate of about 3 to 5% Pbs possibly low due to its fine association with the pyrite. Gangue- fine qtz. threads. 147-151 Described above	4487	147	151	4	30.8	0.90	0.28	1.66	2.16	
155-156	1.0 Massive pyrite as in previous section Terminates sharply at 156.	4488 4489	151 155	155 156	4 1	1.22 0.78	0.23	0.35	1.90	2.26	1.28
156-162	3.2 Light grey sericite schist. 10% white qtz. Shearing platy and contorted. Less than 1% pyrite threads.										
162-166	2.6 As above, with 30% white qtz. replacement. Two small patches of pyrrhotite are surrounded by pyrite, suggestive of a sequence in introduction.										
166-170	2.9 Light grey sericite schist grading to medium grey at 169. Shearing at 80° to core- wavy and fissile. A few threads of pyrrhotite following the shearing - less than 5%.										
170-174 174-179	3.0 3.0 As above. Medium grey sericite schist changing to graphitic schist at 178. 10% pyrrhotite and pyrite containing 3 to 5% fine Pbs and Ens.	4490*	174	179	5						

DRILLED BY Boyles Bros. Ltd.

x not sent for assay

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 5

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	RECOVERY	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Ag	Cu	Pb	Zn		
179-183	2.4	Medium grey sericite schist replaced by 50% pyrrhotite and several rice-sized patches of chalco pyrite. Shearing contorted.	4491	179	183	4						
183-189	4.4	Medium grey sericite schist streaked with graphitic schist. Wavy fissile shearing at 50° to core.										
189-192	2.0	As above. Several threads of pyrrhotite following the shearing - less than 1%.										
192-196	2.0	Medium grey (pearly lustre) sericite schist. Fissile shearing at 60° to core. 10% white qtz.										
196-204	5.4	Medium grey sericite schist containing streaks of graphitic material parallel to shearing at 60° to core. A few specks of pyrrhotite at 203.										
204-208	1.8	As above. Sparse pyrrhotite.										
208-218	8.0	Dark grey sericite schist and graphitic schist (interbanded) at 70° to core. Knotted in part. Sparse pyrrhotite.										
218-223	5.0	60% pyrite and 15% combined and finely associated Pbs and Zns. Several irregular small patches of chalco-pyrite edged by and penetrated by pyrrhotite.	4492	218	223	5	32.9 0.96	0.25	1.68	3.15		
223-227	3.2	As above. the gangue consists of bluish-grey quartz introduction.	4493	223	227	4	30.2 0.88	0.25	2.15	3.24		

DRILLED BY Boyles Bros. Ltd.

X not sent for assay

MINED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Victoria

SHEET NUMBER 6 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	RECOVERY	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
							Ag.	Ca	Pb	Zn				
227-233	6	Same mineralization as above.					56.2							
		227-230 Mineral as described above.	4494	227	230	3	1.64	0.25	1.30	1.87				
		230-233	4495	230	233	3	0.36	0.28	0.22	0.10				
233-245	11.8	Graphitic schist with about 30% greyish-white cherty qtz. in nodular patterns in the shearing. Bluish cast to qtz- when wet. Shearing at 45°, but contorted in part.					12.3							
		233-239 Between 10 and 20% fine pyrite. No Pbs or Zns noted.	4496 ^x	233	239	6								
		239-245 About 5% fine pyrite in graphitic schist described above.	4497 ^x	239	245	6								
245-251	5.6	Graphitic schist containing greyish-white quartz mottling in irregular rice-size patches; the quartz has a bluish cast when wet. Shearing at 55° to core. Sparse pyrite from 245 to 249.												
		249 to 251 - 10 to 15% fine pyrite. A few small threads of pyrrhotite and chalcovrite.	4498 ^x	249	251	2								
251-254	3.0	Graphitic schist 0. '2 of 10% pyrite at 251.2												
254-260	5.0	Graphitic schist containing 2 to 3% fine pyrite. Shearing at 70° to core												
260-266	6.0	Graphitic schist as above. 2 to 3% fine pyrite; sparse pyrrhotite. Partially silicified.												

DRILLED BY Boyles Bros. Ltd.

x not sent for assay

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 7 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag	Cu	Pb	Zn				
	<u>Recovery</u>												
266-272	6.0 As above with 3 to 5% fine pyrite in the shearing.												
	<u>266-269</u>	4499 X	266	269	3								
	<u>269-272</u>	4500 X	269	272	3								
272-277	5.0 Graphitic schist with 5 to 10% pyrite												
277-282	5.0 Qtz. present in the shear planes. Graphitic schist containing 20% pyrite. 30% qtz. with the pyrite. Shearing at 50° to core; local contortions indicative of drag folding. No visible Pbs or Zns.												
	<u>285-290</u>	4501 X	272	277	5								
	<u>290-295</u>	4502 X	277	282	5								
282-285	2.6 As above. 20-25% pyrite.	4503 X	282	285	3								
285-295	9.0 As above- sharply contorted shearing suggestive of small drag-folds.												
	<u>285-290</u> 40% pyrite, 30% associated qtz., balance schist.	4504	285	290	5	4.8	0.14	0.28	0.14	0.20			
	<u>290-295</u> 25% pyrite, 20% qtz., balance schist.	4505	290	295	5	0.90	0.20	0.14	0.10				
295-305	10.0 Graphitic schist changing gradually to medium grey sericite schist at 302.					30.8							
	<u>295-300</u> : 10% fine pyrite. Shearing knotted and contorted.	4506 X	295	300	5								
	<u>300-305</u> : 10-15% pyrite and some pyrrhotite.	4507 X	300	305	5								

DRILLED BY Boyles Bros. Ltd.

X) not sent for assay

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 8 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag	Cu	Pb	Zn				
	<u>Recovery</u>												
305-311	6.0												
	Medium grey sericite schist inter-banded with minor graphitic schist. About 50% irregular white quartz replacement. A few threads of pyrrhotite.												
311-341	22.5												
	Light grey to medium grey sericite schist. Shearing at 60° to core but contorted in part. A few irregular small patches of pyrrhotite totalling 2 to 3%. The schist has a talcy feel, over the normal sericite smoothness.												
341-350	8.0												
	Graphitic schist containing 30% inter-banded silica replacement. (grayish-white)												
	341-344: 20% pyrite, some pyrrhotite.	4508 X	341	344	3								
	344-347: 30% pyrite, 5% pyrrhotite.	4509 X	344	347	3								
350-352	1.9												
	70% massive pyrite; gangue grayish-white qtz. which has a bluish cast when wet. Remnants of graphitic schist present 50° to core.	4510 X	350	352	2								
352-355	3.0												
	50% pyrite, 10% pyrrhotite. Gangue-grayish-white silicification.	4511 X	352	355	3								
355-360	5.0												
	Graphitic schist. About 30% grayish-white silicification in the shear planes. 5% scattered pyrite grains.												
360-369	6.5												
	Graphitic schist with 40% silicification. Shearing 80° to core.												

DRILLED BY Boyles Bros. Ltd.

X not sent for assay

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 9 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag	Cu	Pb	Zn				
	Recovery												
	360-364: 20% pyrite. A little pyrrhotite and chalcopyrite.	4512 ^x	360	364	4								
369-376	2.5 Graphitic schist. Core badly broken. Fragments of white quartz at 369.3												
376-378	1.3 Graphitic schist. 10% pyrite.	4513 ^x	376	378	2			0.62	1.27				
378-381	2.7 As above, 10-15% pyrite.	4514 ^x	378	381	3			2.14	1.27				
381-383	2.0 As above, 10% pyrite, 10% pyrrhotite sparse chalcopyrite.	4515 ^x	381	383	2			1.16	1.87				
383-385	2.0 30% pyrite, 5% Pbs. sparse pyrrhotite and chalcopyrite. Silicified matrix has a reddish (granitic) cast.	4516	383	385	2	1.12		3.30	5.11				
385-389	4.3 Graphitic schist. 20% pyrite, sparse chalcopyrite. Shearing at 45° ab core.	4517 ^x	385	389.3	4.3	38.4							
389.3-401	1.4 Greyish-white sericite schist replaced by 30 to 40% white qtz. No sulphide present. This rock is highly fractured and caved badly in the hole. Clay odour when wet.							0.87	1.38				
	END OF HOLE AT 401												
	RECOVERY: 66%.												

7' - 4.73

DRILLED BY Boyles Bros. Ltd.

not sent for assay

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SLUDGE RECORD

SHEET NUMBER 1, Sludges SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE		
21-25	Scattered fine schist flakes and ovbd. sand. No evidence of sulphide.										Pb	Zn
25-30	As above.											
30-35	Aggregate of dark schist flakes.											
35-40	As above. Some sand. No sulphide.											
40-45	As above.											
45-50	Medium grey schist flakes. No sulphide.											
50-55	As above; some admixed sand.											
55-60	As above.											
60-65	Mixture of light to dark grey schist flakes. No sulphide.											
65-70	As above but with sparse pyrite in small grain.	4445	65	70	5							
70-75	Fine pyrite disseminated in dark grey composite.	4446	70	75	5						2.10	3.74
75-80	Heavy concentration of fine pyrite in dark grey composite.	4447	75	80	5						4.20	4.92
80-85	As above.	4448	80	85	5						1.13	1.77
85-90	As above.	4449	85	90	5						1.35	1.47
90-95	As above.	4450	90	95	5						1.96	2.65
95-100	As above.	4551	95	100	5						3.07	6.38
100-105	Same as above; higher sericite content.	4552	100	105	5						0.64	1.57
105-110	Sparse visible pyrite grains	4553	105	110	5							
110-115	Light grey sericite flakes; no visible sulphide grains.	4554	110	115	5							
115-120	Medium grey, sericitic flakes. No visible sulphide.	4555	115	120	5							
120-125	Sericite flakes and admixed sand											

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 2, Sludges SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
125-130	Aggregate of creamy grey sericite flakes.								
130-135	As above with some admixed sand.								
135-140	As above; no visible sulphide	4556	135	140	5				
140-145	Medium grey; pyrite grains present.	4557	140	145	5				
145-150	Dark brownish grey; numerous pyrite grains present.	4558	145	150	5				
150-155	As above.	4559	150	155	5				
155-160	Less pyrite evident.	4560	155	160	5				
160-165	Light brownish-grey. No visible sulphide.								
165-170	Cream-coloured sericite. No visible sulphide.								
170-175	As above.								
175-180	Brownish-grey aggregate. No visible sulphide.	4561	175	180	5				
180-185	Medium grey, sericitic.	4562	180	185	5				
185-190	Dark grey; minute flakes of sericite. No evidence of sulphide.								
190-195	As above.								
195-200	Medium grey; numerous sericite flakes.								
200-205	As above; no evidence of sulphide.								
205-210	Brownish cast; no evidence of sulphide.								
210-215	Dark grey flakey aggregate.								
215-220	Brown, flakey aggregate. No visible sulphide.	4563	215	220	5				
220-225	Brownish-black. Numerous pyrite grains	4564	220	225	5				
225-230	As above.	4565	225	230	5				
230-235	Considerable fine pyrite.	4566	230	235	5				
235-240	Minute grains of pyrite present	4567	235	240	5				
240-245	As above, in dark grey aggregate.	4568	240	245	5				
245-250	As above; fine pyrite grains.	4569	245	250	5				

Boyles Bros. Ltd.

DRILLED BY _____

SIGNED B. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda

SHEET NUMBER 3, Sludges SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE			SLUDGE SAMPLE		
250-255	Brownish-black; sparse pyrite.	4570	250	255	5						
255-260	As above; sparse pyrite.										
260-265	As above; no visible sulphide.										
265-270	As above; a few pyrite grains	4571	265	270	5						
270-275	Sparse pyrite grains	4572	270	275	5						
275-280	Brownish-black graphitic. Sparse pyrite.	4573	275	280	5						
280-285	As above.	4574	280	285	5						
285-290	As above; fine pyrite grains	4575	285	290	5						
290-295	As above.	4576	290	295	5						
295-300	Scattered pyrite grains.	4577	295	300	5						
300-305	No visible sulphide- in graph. schist.										
305-310	Brownish-grey. No evidence of sulphide										
310-315	As above.										
315-320	As above; sparse pyrite grains.										
320-325	As above Medium grey; fine sericite flakes										
325-330	As above; brownish cast.										
330-335	Same as above.										
335-340	Dark grey; no visible sulphide.										
340-345	Fine grains of pyrite.	4578	340	345	5						
345-350	Graphitic; considerable fine pyrite	4579	345	350	5						
350-355	High pyrite content in black graphitic composite.	4580	350	355	5						
355-360	As above.	4581	355	360	5						
360-365	Some fine pyrite grains	4582	360	365	5						
365-370	Dark grey, sparse pyrite										
370-375	Graphitic., Considerable fine pyrite	4583	370	375	5						
375-380	As above.	4584	375	380	5						

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY Vangorda.

SHEET NUMBER 4, Sludges

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
380-385	As above. Heavy pyrite content	4585	380	385	5								
385-390	Graphitic schist as above- with some fine pyrite grains.	4586	385	390	5								
390-395	Light grey. Sent for assay because of low core recovery.	4587	390	395	5								
395-400	Same as above.	4588	395	400	5								

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD

HOLE NO. 20

PROPERTY Yamaguchi

SHEET NUMBER 1

SECTION FROM 4 TO 5

STARTED Aug. 10, 1953

LATITUDE 29, 71.81

9077.5N

DATUM ADP

COMPLETED Aug. 20, 1953

DEPARTURE 30, 330.36

9214.15E

BEARING 130.276M

ULTIMATE DEPTH 500

ELEVATION 4,015.17 + 25'

1302.76M

DIP

PROPOSED DEPTH To test mineralization on Projected dip.

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
						Ag	Gr	PB	2u		
0-87 ²⁶⁵	Casing (in overburden)										
87-118	0 Sludge only (see sludge record at end of log).										
118-120	1.5 60% pyrite, 10% Fbs and 10% Zns. Gangue - grayish white quartz (AX core)	4601	118	120	2	5.04		6.16	11.60		
120-121	0.7 Massive sulphide - 80% pyrite, 20% combined Fbs and Zns. (AX core)	4602	120	121	1	3.30		6.16	7.08		
121-132	0.2 Two fragments of gray sericite schist (AX core) No sulphide mineralization.										
132-134	1.1 Fragments of medium gray sericite schist. No sulphides. (AX core from 132 on).										
134-139	1.3 Medium gray, platy sericite schist. Shearing at 80° to core.										
139-141	0 (See sludge record at end of log.)										
141-144	1.7 Medium gray sericite schist. A pea-sized patch of pyrrhotite at 143.5. Shearing contorted.										
144-149	1.0 Buttons of medium gray sericite schist. No sulphide.										
149-155	0 See sludge record.										
155-160	1.0 Light gray sericite schist fragments. One fragment of white quartz.										
160-165	1.1 Fragments of light gray sericite schist. Talcy keel on shear planes.										
165-170	1.2 As above. Shearing 80° core.										
170-171	0 See sludge record.										

DIAMOND DRILL RECORD,

HOLE NO. 10

PROPERTY Yanagoda

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	Recovery	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
							Hg	Cu	Pb	Zn		
171-182	3.8	White quartz. No sulphide mineralization.										
182-182.5	0.5	Medium grey sericite schist. Shearing 60° to core.										
182.5-	1.8	White quartz. Vestiges of sericite schist.										
184.3-184.3-187	2.7	Massive sulphide consisting of 7% pyrite and 25% combined Pb and Zn. (Former appears to be in greater amount.)	4603	184.3	187	2.7	4.30		10.53	10.33		
187-192	5.0	Massive sulphide consisting of an estimated 15% to 25% combined Pb and Zn; balance fine pyrite.	4604	187	192	5	2.90		3.52	7.86		
192-196	4.0	As above; possibly a higher Pb-Zn content.	4605	192	196	4	1.24		1.54	3.24		
196-201	5.0	As above. Sparse quartz gangue in small threads and patches.	4606	196	201	5	3.00		7.70	6.39		
201-202.8	1.8	Massive pyrite, Pb and Zn as in above sections.	4607	201	202.8	1.8	2.86		8.14	8.95		1634
202.8-223	8.6	Light grey talc sericite schist. Shearing contorted. Streaks of chloritic material.		202.8	223	20.2			0.0	0.0		
223-226	3.0	Massive sulphide; pyrite with about 15% combined Pb and Zn. Some quartz gangue from 225.2 to 226.	4608	223	226	3	0.80		1.76	2.36		
226-231	4.2	60% pyrite, 25% combined Pb and Zn. Balance greyish white silicification.	4609	226	231	5	0.84		1.76	3.05		

DIAMOND DRILL RECORD,

HOLE NO. 30

PROPERTY Vanguard

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION		SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
							Ag	ln	Pb	Zn					
231-236	5.0	As in previous section; gangue slightly greater from 235 to 236.	4610	231	236	5	0.94		2.09	2.46					
236-241	4.6	50% pyrite, 20% combined Pbs and Zns.	4611	236	241	5	0.36		0.44	0.49					
241-246	5.0.	Balance siliceous gangue. 40% Pbs from 241 to 243.5 Balance in ratio as in previous sample.	4612	241	246	5	2.06		4.62	4.33					
Please note:		The hole lost water at 245'. No sludges past this point. Casing tight and not able to advance it further.													
246-251	4.2	70% massive sulphide consisting of pyrite and 10 to 15% associated Pbs and Zns. Gangue - silicified sericite schist. 0.1 pyrrhotite.	4613	246	251	5.0	1.24		2.75	3.05					
251-259	4.8	Light grey sericite schist. Shearing at 40° to core. No sulphide present.		251	259	8			0	0					
259-262	3.0)	0.1 pyrrhotite at 259. Massive sulphide containing estimated 15 to 20% combined Pbs and Zns, 70% pyrite and balance quartz gangue.	4614	259	262	3	1.12		2.42	3.05					
262-267	4.5	Massive sulphide as above. Possibly a slightly higher Pbs-Zns content.	4615	262	267	5	2.12		2.42	5.02					
267-270.1	1.5	Light grey sericite schist. Talay, shearing contorted.		267	270.1	3.8	∅		0	0					
270.1-274	3.9	Massive pyrite containing about 20% Pbs + Zns in close association.	4616	270.1	274	3.9	2.70		4.84	6.60					

DIAMOND DRILL RECORD,

HOLE NO. 10

PROPERTY Vanguard

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery					<i>Ag</i>	<i>Pb</i>	<i>Zn</i>	
<i>274-278</i>	3.3 Massive sulphide as in previous section	4617	<i>274</i>	<i>278</i>	<i>4</i>	<i>1.70</i>	<i>4.28</i>	<i>5.41</i>	
<i>278-279.6</i>	1.6 Light grey sericite schist containing specks of chloritic material. Shearing contorted.		<i>278</i>	<i>279.6</i>	<i>1.6</i>		<i>0</i>	<i>0</i>	
<i>279.6-284</i>	4.4 Massive sulphide containing estimated 70% pyrite, 25% combined Pbs and Zns. balance gangue silica.	4618	<i>279.6</i>	<i>284</i>	<i>4.4</i>	<i>1.50</i>	<i>3.85</i>	<i>3.93</i>	
<i>284-286</i>	2.0 Massive pyrite. Sparse Pbs and Zns content indicated from visual examination.	4619	<i>284</i>	<i>286</i>	<i>2</i>	<i>0.60</i>	<i>0.99</i>	<i>0.89</i>	
<i>286-291</i>	5.0 70% pyrite, 20% combined Pbs and Zns, former in greater amount.	4620	<i>286</i>	<i>291</i>	<i>5</i>	<i>0.94</i>	<i>3.08</i>	<i>1.68</i>	
<i>291-296</i>	5.0 50% pyrite, 15% combined Pbs and Zns. Sparse pyrrhotite and chalcocopyrite.	4621	<i>291</i>	<i>296</i>	<i>5</i>	<i>0.56</i>	<i>1.10</i>	<i>.79</i>	<input checked="" type="checkbox"/>
<i>296-301</i>	Balance - white quartz. As above. Pyrite more massive from 299 to 301. Pbs and Zns less prominent.	4622	<i>296</i>	<i>301</i>	<i>5</i>	<i>0.61</i>	<i>1.54</i>	<i>.79</i>	
<i>301-306</i>	5.0 Massive pyrite with an estimated 20% Pbs and Zns. About 10% siliceous gangue.	4623	<i>301</i>	<i>306</i>	<i>5</i>	<i>1.40</i>	<i>3.63</i>	<i>3.34</i> <i>2.75</i>	
<i>306-311</i>	5.0 As above but with increase of siliceous gangue and sericite schist remnant from 308.5 to 311. Trace of shearing at 40° to core.	4624	<i>306</i>	<i>311</i>	<i>5</i>	<i>2.30</i>	<i>2.09</i>	<i>6.00</i>	
<i>311-316</i>	5.0 Massive pyrite containing an estimated 20% Pbs and Zns. About 10% silica in the sulphide interstices.	4625	<i>311</i>	<i>316</i>	<i>5</i>	<i>2.08</i>	<i>6.92</i>	<i>4.43</i>	

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 20

PROPERTY _____

SHEET NUMBER 5 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
						Ag	Cu	Pb	Zn		
	Recovery										
316-321	5.0 Massive sulphide as in previous section grading to partially replaced silicified schist from 320 to 321. Fine speckles of Pbs throughout.	4626	316	321	5 ²¹	1.76		2.42	5.22		
321-326	5.0 50% pyrite, 15% combined Pbs and Zn. Balance- siliceous (grey) gangue.	4627	321	326	5	1.92		2.86	4.92		
326-331	5.0 As in previous section. Trace of replaced shearing at 50° to core.	4628	326	331	5	1.76		2.53	4.62		
331-336	4.5 At 326 there is a change from pyrite to pyrrhotite (massive) over a distance of about 0'.3. The pyrrhotite contains some pyrite patches, scattered threads of reddish-brown Zn and a little Pbs.	4629	331	336	5	1.40		2.20	3.34		
336-338.	2.2 As above. Scattered threads of bluish grey quartz containing Pbs and Zn.	4630	336	338.2	2.2	1.50		1.98	3.34		
338.2-341	2.0 Medium grey sericite schist. No sulphide. Shearing at 40° to core.										
341-345	3.5 Dark grey sericite schist containing graphitic laminations. Shearing at 50° to core. No sulphide.										
345-349	3.9 As above. 10% irregular quartz threads.										
349-354	4.8 Light greenish-grey sericite schist. About 3% pyrrhotite in irregular patches up to 1/4"; sparse chalcopyrite.										
354-358	As above, but more talcy. Sparse pyrrhotite. About 10% irregular white quartz patches.										

DIAMOND DRILL RECORD,

HOLE NO. 10

PROPERTY Vangorda

SHEET NUMBER 6 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery								
358-363	3.4 Medium grey sericite schist. Several small patches of pyrrhotite, less than 3%.								
363-368	4.0 Same as previous section. Shearing 60° to core.								
368-373	4.7 As above.								
373-380	6.5 As above. About 10% irregular white quartz replacement.								
380-384	4.0 The medium grey sericite schist has a greenish cast in part. Shearing at 60° to core.								
384-389	5.0 As above. Several 1/2" patches of pyrrhotite. Not worthy of assay.								
389-395	5.0 Light greenish grey sericite schist. A little pyrrhotite. Shearing at 60° to core.								
395-396	1.0 0'.7 of massive fine grained pyrite from 395.1 to 395.8. No visible Pbs or Zns.	4631 ^x	395	396	1				
396-399	3.0 Graphitic schist. Shearing at contorted angles to core.								
399-404	4.0 Same as previous section. 399-400: 0'.3 fine massive pyrite. No visible Pbs or Zns.	4632 ^x	399	400	1				
404-411	4.9 Interbanded graphitic and sericite schist. Some greyish-white silicification.								

DRILLED BY _____

Start out for assay

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 30

PROPERTY Vanguard

SHEET NUMBER 7 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery								
411-419	5.6 Same as previous section. About 1% white quartz.								
419-428	5.2 Medium gray sericite schist. 40% white quartz. A little pyrrhotite - not worthy of assay.								
428-439	5.0 Medium gray sericite containing a little interbedded graphitic material. Shearing at 50° to core.								
433-443	7.0 Medium gray, finely bedded sericite schist.								
443-473	20.0 Medium gray sericite schist as above. Shearing at 60°.								
473-474	1.5 About 20% pyrrhotite, 20% white quartz in medium gray sericite schist.	4633 ✓	473	474.5	1.5				
474.5-489	11.5 Medium gray, fissile sericite schist. Shearing at 60° to core. Sparse pyrrhotite in threads. 1% white quartz patches.								
489-555	37.6 Medium gray sericite schist. Shearing at 80° to core. White quartz from 526.5 to 527 and from 535 to 536.6.								
	END OF HOLE AT 555								
	RECOVERY: 65%								

DRILLED BY Boyles Bros. Ltd.

Just sent for assay

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 10

PROPERTY Yamgoda

SLUDGE RECORD

SHEET NUMBER 1 SLUDGES SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
87-92	No core recovered. Sludge is light grey sericite. No evidence of sulphide.	4591	87	92	5				
92-97	As above.	4589	92	97	5				
97-102	Same as above two sections.	4590	97	102	5				
102-107	Light grey sericite flakes.	4592	102	107	5				
107-112	As above.	4593	107	112	5				
112-117	As above.	4594	112	117	5				
117-120	Sericite flakes as before.	4595	117	120	3				
120-122	Darker than previous sections.	4596	120	122	2				
122-124	As above.	4597	122	124	2				
124-126	Light grey sericite.	4598	124	126	2				
126-131	As above. No evidence of sulphide.	4599	126	131	5				
131-136	Medium grey sericite aggregate.	4600	131	136	5				
136-141	Light grey sericite. No evidence of sulphide.	4651	136	141	5				
141-146	As above.	4652	141	146	5				
146-151	Same type of sericite aggregate.	4653	146	151	5				
151-155	As above. Slight brownish cast.	4654	151	155	4				
155-160	Brownish tone. Possibly sand contamination.	4655	155	160	5				
160-165	Pale brown. Possibly oxidation during drying of sample.	4656	160	165	5				
165-170	Grayish brown. No evidence of sulphide.	4657	165	170	5				
170-175	Grey with brown speckles.	4658	170	175	5				
175-180	Pale grey. Some sericite flakes.	4659	175	180	5				
180-185	As above.	4660	180	185	5				
185-187	Core recovery good.								
187-189	Complete core recovery.								
189-191	As above.								

Boyles Bros. Ltd.

DRILLED BY _____

SIGNED E. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 10

PROPERTY Vanguard

SLUDGE RECORD

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
191-196	No core loss.												
196-201	No core loss.												
201-206	Water loss started here. A little sludge.	4661	201	206	5								
206-211	No water return.												
211-216	A little water return. Light grey sludge.	4662	211	216	5								
216-221	Water return poor. A little grey sludge.	4663	216	221	5								
221-224	As above.	4664	221	224	3								
224-226	Complete core return; very little sludge.												
226-230	As above.												
230-235	Complete core recovery; small amount of sludge.												
235-240	Good core recovery; very little sludge return.												
240-245	Complete core recovery; fair sludge return. Return water lost at												
245-250	Slight sludge return. Return water lost at 245. No further sludges from hole - good core recovery in the mineralized sections from here on.												

Boyles Bros. Ltd.

DRILLED BY _____

SIGNED _____

R. W. Baker

	3'	1 3'	POINT Pb+2N	POINT 48.76	→ 16.252/ 118-121' / 3'
184.3	3.7		20.86	56.42	
↓	5		11.38	56.90	
	4		4.78	19.12	
	5		14.09	70.45	
202.8	1.8		17.09	30.75	
1	18.5'			233.64	→ 12.61/18.5' 184.3 → 202.8'
222-223	20.2		0.00	0.00	

223	3'		4.12	12.36	
	5		4.81	24.05	
	5		4.55	22.75	
	5		9.3	4.65	
	5		8.95	+ 4.75	
251	5/28		3.80	24.00 = 137.56	→ 4.97/28' 223 → 251'
	8		0.00	0.00 X	

257	3		5.47	16.41	
267	5/8'		7.44	37.20 → 53.61	→ 7% 6.7/8' 257 → 267'
70.1	2.9		0.00	0.00 X	

271			139.42	5.46	
10.9			143.42	5.85	
36.2			44.71	5.24	
36			188.17		

270.1	3.9		11.74	44.62	
↓	4		9.69	38.76	
	1.6		0.00	0.00 X	
	4.4		7.78	34.23	48.76 — 3'
	2		1.88	3.76	233.64 18.5
	5		4.76	23.80	137.56 28'
	5		1.89	9.45	53.61 8'
	5		2.33	11.65	450.56 68.1
	5		6.97	34.85	924.13 135.6
	5		8.09	40.45	6.81 % PbZ 7.37%
	5		11.35	56.75	
	5		7.64	38.20	
	5		7.78	38.90	
	5		7.15	35.75	
	5		5.54	27.70	

338.2	2.2		5.32	11.69	
68.1			450.56	6.62	
4.6				6.78%	
68.5					

6.6% / 68.1'
270.1 → 338'

DIAMOND DRILL RECORD,

HOLE NO. 11

PROPERTY Yamgora

SHEET NUMBER 1

SECTION FROM _____ TO _____

STARTED Nov. 20, 1953.

LATITUDE 28,629.24

DATUM 4000'

COMPLETED Nov. 26, 1953.

DEPARTURE 31.257.25

BEARING 47° 11' ea.

ULTIMATE DEPTH 445'

ELEVATION 4043.25

DIP 41° at collar; 50° at 445'

PROPOSED DEPTH to cross-section mineralisation

DEPTH FEET	RECOVERY	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE						
							Ag	Cu	Pb	Zn							
0-23	75	Casing (overburden)															
23-40	1.5	White quartz fragments.															
40-55	2.0	Mixed fragments of white quartz and dark grey sericite schist. (Possibly graphitic interbands).															
55-70.5	3.5	80% white quartz containing vestiges of dark grey schist.		(450)?													
70.5-87.5	8.4	Massive pyrite with some fine Pb and Zn which shows an increase toward the end of the section at 87.5.															
		70.5-75: Mineralization described above.	4518	70.5	75	6.5	24.7 0.72	0.15	0.88	0.79							1.5
		75-79: Mineralization as described on previous page. <i>above</i>	4519	75	79	4	42.5 1.24	0.23	1.10	0.98							2.0
		79-84: As above. Pb and Zn begins to show an increment in this section.	4520	79	84	5	220.1 6.42	0.18	3.41	6.39							2.5
		84-87.5: Mineralization as above. This section shows greatest Pb-Zn content of above four samples—possibly 20% combined.	4521	84	87.5	3.5	53.5 1.96	0.35	2.75	4.72							2.4
																	8.4
87.5-91	1.0	Graphitic schist containing 40% grayish-white silicification and 20% Zn. A little Pb visible.	4522	87.5	91	3.5	34.3 1.00	0.25	3.30	9.34							
91-101	5.0	Grayish-white quartz containing remnants of graphitic schist. Some sericite in close association with the quartz.				10'											

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 11

PROPERTY Vangoria

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE	
						Ag	Cu	Pb	Zn		
	Recovery										
91-101	5.0 Greyish-white quartz containing remnants of graphitic schist. Some sericite in close association with the quartz.										
101-106	3.5 Massive pyrite containing some finely associated Pbs and Zns. Amount difficult to estimate.	4523	101	106	5	74.0 2.16	0.13	1.93	7.86		
106-111	4.7 Same as above.	4524	106	111	5	64.4 1.88	0.20	4.39	6.35		
111-115	4.0 Same as above. The fine Pbs occurs in streaks through the pyrite.	4525	111	115	4	20.4 1.10	0.23	3.62	5.70		
115-120	5.0 About 30% greyish-white siliceous gangue. Balance, pyrite with a little visible Pbs. No Zns noted, although it may be closely associated with Pbs.	4526	115	120	5	6.9 0.60	0.20	0.77	1.67		
120-124	4.0 As above, but with 10% siliceous gangue.	4527	120	124	4	0.20	0.45	1.55	1.77		
124-130	6.0 Greyish-white sericite schist containing about 5% pyrrhotite. Shearing at 55° to 70° to core.										
130-157	22.2 As above. Shearing at 70° to core. Pyrrhotite content not worthy of assay.										
157-177	15.5 Medium grey sericite schist. Some irregular white quartz replacement. Shearing at 70° to core.										
177-179	2.0 As above. About 10% pyrrhotite in irregular patches.	4528	177	179	2	5.1 0.15	0.10	0.14	0.20		

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 11

PROPERTY Vanguard

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
						Ag	Cu	Pb	Zn					
	Recovery													
	318-321: 20% pyrite and pyrrhotite. Some fine galena present.	4532 ^x	318	321	3.0									
323-333	8.4 Graphitic schist laminated with light grey sections which may be the result of silicification or original quartzite interbanding.													
333-336	13.5 Same as above with slightly higher graphitic schist content.													
	334.6-335.6: 40% pyrrhotite. Sparse muscovite.	4533 ^x	334.6	335.6	1.0									
356-361	3.1 Graphitic schist highly silicified with only vestiges of graphitic sections remaining. 20% fine pyrite; some Pb present.	4534 ^x	356	361	5.0									
361-366	4.4 Grayish-white shaly silicification containing 30% pyrite and a little Pb. Streaks of graphitic schist at 365.5	4535 ^x	361	366	5.0									
366-369	2.1 Graphitic schist with 40% grayish-white silica interbanding. 10% pyrite.	4536 ^x	366	369	3									
369-374	5.0 Graphitic schist with about 20% grayish white silica laminations. 10% white quartz threads. 10% pyrite; sparse visible Pb.	4537 ^x	369	374	5									
374-395.1	14.2 Graphitic schist as above. Sluicing at 30° to core.													
	374-379.1 15% fine pyrite, sparse Zn noted.	4538 ^x	374	379	5									

DRILLED BY _____

x) not sent for assay

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 11

PROPERTY Vungorda

SHEET NUMBER 5 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag	Cu	Pb	Zn				
	Recovery												
	379-384: Same mineralization as previous section.	4539 ^x	379	384	5								
	384-389: About 20% fine pyrite with a little pyrrhotite. Sparse Zn visible.	4540 ^x	384	389	5								
	389-395.2: 15% pyrrhotite with threads of chalcopyrite. About 5% pyrite. Occasional thread of Zn visible.	4541 ^x	389	395.2	6.2								
395.2-429	30.4 Light gray to medium gray sericite schist. Shearing at 50° to core. Some irregular sections of chlorite associated with white quartz.												
	418-421: About 15% irregular pyrrhotite threads. A little chalcopyrite and Zn visible. In the balance of the section from 395.2 to 429 there is a little pyrrhotite, less than 5%.	4542 ^x	418	421	3								
429-	3.8 White quartz. About 20% chlorite patches. For Ag only.	4543 ^x	429	434.2	5.2								
434.2-	1.1 Medium greenish-gray sericite schist.												
436													
436-445	5.0 Medium gray sericite schist containing about 10% irregular white quartz patches. Shearing at 55° to core.												
	END OF HOLE AT 445.												
	RECOVERY: 66.												

DRILLED BY _____

x list sent for assay

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 21

PROPERTY Vangorda

SLUDGE RECORD

SHEET NUMBER 1 - SLUDGES SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO	FROM	TO	WIDTH	CORE SAMPLE			SLUDGE SAMPLE	
									Pb	Zn
65-70	A few pyrite grains visible in a grayish-brown aggregate.	4665	65	70	5				0.15	0.54
70-75	Numerous fine pyrite grains.	4666	70	75	5				0.81	1.57
75-80	As above.	4667	75	80	5				0.50	1.47
80-85	Pyrite grains in brownish-grey aggregate. Some galena recognizable.	4668	80	85	5				0.95	3.84
85-90	Considerable pyrite in brownish-black aggregate.	4669	85	90	5				1.20	3.14
90-95	Light brownish grey. Some pyrite grains visible.	4670	90	95	5				0.37	0.93
315-320	Black, graphitic sludge containing fine grains of pyrite.	4671 ^x	315	320	5					
355-360	Dark grey, to medium grey. Scattered pyrite grains.	4672 ^x	355	360	5					

DRILLED BY Boyles Bros. Ltd.

X not sent for assay

SIGNED R. W. Baker.

DIAMOND DRILL RECORD,

HOLE NO. 12

PROPERTY Vangorda

SHEET NUMBER 1

SECTION FROM _____ TO _____

STARTED Nov. 27, 1953.

LATITUDE 28,627,20

DATUM 4000'

COMPLETED Nov. 28, 1953.

DEPARTURE 31,155.35

BEARING 47° 11' az.

ULTIMATE DEPTH 162'

ELEVATION 4043.3

DIP 68° at collar.

PROPOSED DEPTH To establish dip of mineralization.

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery						Pb	Zn	
0-23.1	Casing (over burden)								
23-66.7	17.6 Light grey sericite schist containing about 15% irregular white quartz patches. 3 to 5% fine pyrite with a little Pbs and Zns, insufficient for assay.								
66.7-71.5	3.7 Light grey sericite schist with 10% pyrite. Some Pbs and Zns with the pyrite.	4544	66.7	71.7	5		4.06	9.14	
71.7-74.2	1.2 Greyish-white sericite schist. Shearing contorted.		71.7	74.2	2.5				
74.2-76	1.8 Massive fine pyrite containing 15% Pbs and 5% Zns. A little siliceous gangue.	4545	74.2	76	1.8		1.98	6.68	
76-81	4.2 As above with an estimated 20% combined Pbs and Zns.	4546	76	81	5		2.86	6.28	
81-83.6	2.6 Same as previous section. Relict shearing at 80° to core - laminated with sulphide.	4547	81	83.6	2.6		2.20	2.84	
83.6-91.5	2.3 G Graphitic schist with 20% greyish-white silicification. Sparse pyrite.				4.1				
91.7-96	3.1 Massive pyrite with considerable fine galena - grey colour to core. Some fine siliceous gangue, about 10%. Estimated Pbs - Zns about 25%.	4548	91.7	96	4.3		3.95	10.50	
96-100	4.0 Same as previous section. Fine siliceous gangue in the interstices of the sulphide.	4549	96	100	4		2.86	6.77	

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 12

PROPERTY Vangorda

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE			SLUDGE SAMPLE	
	Recovery						4b	2n		
100-105	5.0 Same as previous section, but with some pyrrhotite present in irregular patches. About 15% siliceous gangue.									
105-108.2	3.2 Pyrite with sections of massive pyrrhotite. About 10% closely associated Pbs and Zns. Siliceous gangue at irregular intervals.	4550	100	105	5		2.20	2.20	3.04	
108.2-162	26.2 Light grey to medium greenish-grey sericite schist, grading to the latter at about 123. Shearing at 70° to core. Sparse pyrite and pyrrhotite. 10% white quartz.	4701	105	108.2	3.2		1.24	1.38		
	END OF HOLE AT 162.									
	RECOVERY: 68%.									

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker.

DIAMOND DRILL RECORD,

HOLE NO. 13

PROPERTY Vancouver

SHEET NUMBER 1

SECTION FROM 0 TO 120

STARTED Nov. 27, 1953

LATITUDE 29,807.70 9085.39

DATUM 4000

COMPLETED Dec. 3, 1953

DEPARTURE 30,003.57 9145.09

BEARING _____

ULTIMATE DEPTH 305

ELEVATION 3986.18 1293.93

DIP Vertical

PROPOSED DEPTH indicated strike.
To test mineralization on

DEPTH FEET	RECOVERY	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
0-69	21	Casing (over burden)								
69-85	0	Sludge only: (See sludge record at end of log).								
85-92	5.0	Massive sulphide consisting of 60% Pyrite, 30% combined Pbs and Zns and balance gangue.								
		85-87: As described.	4634	85	87	2				
		87-92: As described.	4635	87	92	5		11.06	4.42	
92-101	0.4	Medium grey sericite schist fragments.								
101-120	1.1	Medium to light grey sericite schist (buttons) Shearing 80° to core.								
120-125	0	Lost water temporarily in this section.								
125-127	1.5	White sericite schist containing 30% pyrite and 15% Pbs and Zns. Shearing at 80° to core.	4636	125	127	2		69	2.26	
127-131	1.2	Interbedded graphitic and sericite schist. Fragments only.								
131-135	1.1	As above. Shearing at 80° to core.								
135-146	8.1	Light grey, fissile tale sericite schist.								
146-151	1.2	Graphitic schist fragments								
151-155	3.2	Grayish-white sericite schist. Shearing at 80° to core. Two small patches of pyrite, not worthy of assay.								
155-159	3.2	Sericite schist as above.								
		156.2 to 157.6: Massive sulphide and quartz fragments. Pbs and Zns present in the pyrite.	4637	156.2	157.6	1.6				

DRILLED BY Boyles Bros. Ltd.

1/2nd sheet for assay

SIGNED R. W. Egan

DIAMOND DRILL RECORD,

HOLE NO. 13

PROPERTY Yungoda

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery						Pb	Zn	
159-165.4	5.8 Greyish-white sericite schist. 10% white quartz. Shearing at 80°-90° to core.								
165.4-169.5	2.7 Massive Pyrite with an estimated 10% Pbs and Zns combined.	4638	165.4	169.5	4.1		1.87	3.74	
169.5-171	1.5 As above but with a higher Pbs and Zns content indicated, from grey streaks paralleling relief shearing. Change from Ax to Bx core at 171.	4639	169.5	171	1.5		2.02	3.24	
171-175	2.0 As above with an increase of white quartz toward end of section	4640	171	175	4		1.41	2.06	
175-180	3.2 40% irregular pyrite replacing grey sericite schist. Pbs and Zns visible in the pyrite.	4641	175	180	5		0.99	2.16	
180-186	2.6 80% pyrite and pyrrhotite mottled with remnants of sericite schist and grey siliceous gangue. Pbs and Zns visible in the pyrite	4642	180	186	6		2.02	1.86	
186-191	2.5 Medium grey sericite schist containing streaks of chlorite. Shearing at 60° to core.								
191-196	0.7 Light grey sericite schist. 20% white quartz.								
196-201	1.7 Graphitic schist. Shearing contorted. About 15% pyrite and pyrrhotite, and about 10% combined Pbs and Zns.	4643	196	201	5		0.75	1.77	
201-206	1.3 Graphitic schist. Shearing contorted. One 1/2" patch of pyrite; not worthy of assay.								

Boyles Bros. Ltd.

DRILLED BY _____

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 13

PROPERTY Vangorda

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	<u>Recovery</u>						<u>96</u>	<u>2m</u>	
206-211	0.7 Interbanded graphitic and medium gray sericite schist.								
211-224.3	2.7 Graphitic schist.								
224.3-229	4.7 Graphitic schist. Scattered threads of pyrrhotite. A little chalcocopyrite, pyrite; sparse Pbs and Zns.	4644	224.3	229	4.7		0.33	0.39	
229-238	7.1 Graphitic schist containing irregular patches of pyrrhotite. Sparse pyrite, chalcocopyrite, Pbs and Zns. Not worthy of assay. Shearing at 30° to core.								
238-244.2	4.2 Graphitic schist. 10 to 15% pyrrhotite sparse chalcocopyrite, Shearing at 50° to core.								
244.2-247	2.8 Graphitic schist gradational to grayish-white siliceous rock at 247. appears to be silicification accompanying about 30% pyrrhotite and a little chalcocopyrite.	4645	244.2	247	2.8				
247-252	3.1 As above but with about 5% pyrrhotite.								
252-299	39.0 Medium greenish-gray sericite schist. Shearing at 30° to core. 10' section of pyrrhotite at 257.5, and a 0.1' section of pyrrhotite at 291.6 with some Pbs; insufficient for assay.								

DRILLED BY _____

? Not recovered.

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 13

PROPERTY Vergorda

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE			SLUDGE SAMPLE		
	<u>Recovery</u>										
333- <u>369.6</u>	<u>19.6</u>										
369.6 <u>385</u>	<u>5.2</u>										
	<u>Medium grey to dark grey sericite schist. Sparse pyrrhotite at 364. Graphitic schist. Shearing contorted; average 30° to core.</u>										
	<u>END OF HOLE AT 385.</u>										
	<u>CORE RECOVERY; 50%.</u>										

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 13

PROPERTY Vangorda

SLUDGE RECORD

SHEET NUMBER 1 SLUDGES SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO	FROM	TO	WIDTH	CORE SAMPLE			SLUDGE SAMPLE			
70-75	Light grey sericite sludge. No evidence of sulphide.											
75-80	As above.											
80-85	As above; no evidence of sulphide.											
85-87	Brownish-grey; some grains of pyrite visible.	4673	85	87	2							
87-89	Greyish-black; fine sulphide grains.	4674	87	89	2							
89-91	Sparse visible sulphide.	4675	89	91	2							
91-93	Heavy concentration of pyrite, Zns and Fbs grains.	4676	91	93	2							
93-95	Strong sulphide concentration as above.	4677	93	95	2							
95-100	Sericite flakes; no visible sulphide.											
100-123	Sludges are pale grey and consist chiefly of sericite flakes and silica grains. No visible sulphide.											
123-125	Brownish-grey; sparse pyrite.	4678	123	125	2							
125-129	No evidence of sulphide in sludge.											
129-131	Black and grey grains admixed. Sparse pyrite.	4679	129	131	2							
131-165	No evidence of sulphide; sludges not worthy of assay.											
165-170	Some fine sulphide grains.	4680	165	170	5							
170-175	Fine sulphide; pyrrhotite picked up by magnet.	4681	170	175	5							
175-178	Sulphide grains in dark grey sludge.	4682	175	178	3							
178-180	Light grey; sulphide grains present.	4683	178	180	2							
180-182	Mineralized section of low core recovery.	4684	180	182	2							
182-184	As above.	4685	182	184	2							
184-186	As above.	4686	184	186	2							
186-196	No evidence of sulphide.											

DIAMOND DRILL RECORD,

HOLE NO. 13

PROPERTY Vangorda

SLUDGE RECORD

SHEET NUMBER 2 SLUDGES SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
196-201	Some fine sulphide grains.	4687	196	201	5								
201-226	Core recovery good in mineralised portions; sludges also good.												
226-246	Core recovery good.												
	WATER RETURN LOST AT 246.												
	No further sludges.												

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 14

PROPERTY Vangorda

SHEET NUMBER 1 SECTION FROM 0 TO 65 STARTED Nov. 30, 1953
 LATITUDE 28,493.66 DATUM ADN COMPLETED Dec. 6, 1953
 DEPARTURE 31,009.11 BEARING 48° 20' 21" north ULTIMATE DEPTH 77
 ELEVATION 4043.91 DIP 43° 57' at collar; 48° 30' at 350' PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE			
						Ag	Cu	Pb	Zn				
Recovery													
0-8 (2.4)	0												
8-30	0.2												
30-46	0.1												
46-51	0.1												
51-57	0.8												
57-62	1.1												
62-65	1.5	4702	62	65	3			1.69	4.23				
65-70	1.6												
70-73	1.8	4703	70	73	3			1.77	4.33				
73-77	3.9	4704	73	75.3	2.3			3.55	7.27				
77-82	2.9		76.3	82.0	5.7								
82-86	2.1	4705	82	86	4			3.44	6.68				

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 14

PROPERTY Vangorda

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE				SLUDGE SAMPLE				
						P ₄	Cu	Pb	Zn					
	Recovery													
86-91	3.1 Medium grey sericite schist grading to graphitic schist at midpoint of run.													
91-140	13.5 Graphitic schist. Very carbonaceous and brittle. Shearing contorted, varying from 30° to 5° to core.													
140-174	13.4 Graphitic schist. Wavy schistosity at 10° to 30° to core. Core brittle and badly broken. Change to KI core, at 170'.													
174-263.5	51.7 Medium grey sericite schist. Some chloritic sections associated with irregular patches of white quartz. Shearing at 50° to core. Scattered threads of pyrrhotite and reddish-brown sphalerite; sparse chalcopyrite. Not more than 3% sulphide. Not worthy of assay.													
263.5-275	4.5 Graphitic schist mottled with rice-sized patches of grey quartz in the plane of shearing. - at 50° to 60° to core. About 10% fine pyrite; no visible Pbs or Zns. Not worthy of assay.													
275-280	5.0 Graphitic schist silicified with about 60% grey quartz. 20% fine pyrite; sparse chalcopyrite. No visible Pbs or Zns.	4706	275	280	5			0.17	0.39					

DIAMOND DRILL RECORD,

HOLE NO. 14

PROPERTY Tangorda

SHEET NUMBER 3

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	FORMATION		SAMPLE NO	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery									
280-284	1.8	80% massive pyrite. Balance gray siliceous gangue. No visible Pbs or Zns.	4707	280	284	4				
284-288	3.1	As above. Several streaks of fine Pbs at 285.	4708	284	288	4				
288-292	4.0	Massive pyrite with about 5% Pbs; no visible Zns.	4709	288	292	4				
292-294.5	2.5	As above; possibly a little more Pbs	4710	292	294.5	2.5				
294.5-329	15.8	Graphitic schist with about 20% inter-banded gray silica. Shearing clearly defined at 80° to core.								
329-343	12.0	Graphitic schist. Shearing at 45° to core.								
343-348	5.0	Graphitic schist replaced by about 60% pyrite, 10% pyrrhotite, 5 to 10% Pbs and reddish-brown Zns. Balance -gray siliceous gangue and graphitic remnants.	4711	343	348	5				
348-350.7	2.7	As in previous section but with slightly higher pyrite and Pbs-Zns content.	4712	348	350.7	2.7				
350.7-371	14.0	Graphitic schist containing about 30% gray silica following the shear planes. Shearing at 80° to core.								
		END OF HOLE AT 371								
		CORE RECOVERY; 46%								

Boyles Bros. Ltd.

R. W. Baber

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 14

PROPERTY Vanguard

SLUDGE RECORD

SHEET NUMBER 1 SLUDGES SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
8-20	Numerous sericite flakes in light brown matrix of oxidised material.	4688	8	20	12				
20-25	As above.	4689	20	25	5				
25-30	Pale brown; highly oxidised.	4690	25	30	5				
30-60	Dark grey indicative of graphitic schist. No evidence of sulphide.								
60-65	Core contains fragments of pyrite.	4691	60	65	5				
65-70	Patch of pyrite in the core.	4692	65	70	5				
70-75	Pyrite fragments in core; low recovery.	4693	70	75	5				
75-80	Mineralised section of good core recovery.								
80-85	Mineralised section of low core recovery.	4694	80	85	5				
85-175	Greyish-black, indicative of graphitic schist.								
175-235	Medium grey sludge indicative of sericite schist.								
235-240	Brownish cast to sludge due possibly to sulphide.	4695	235	240	5				
240-265	Medium grey to dark grey sludge. No evidence of sulphide.								
265-275	Dark grey, indicative of graphitic material.								
275-280	Mineralised section of good core recovery.								
280-285	Mineralised section of low core recovery.	4696	280	285	5				
285-295	Mineralised section of good core recovery.								
	SLUDGE return lost at 300'.								

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 1

SECTION FROM _____ TO _____

STARTED Dec. 8, 1953

LATITUDE 30,006.25 9145.90

DATUM 4000

COMPLETED Dec. 13, 1953

DEPARTURE 30,293.65 9233.50

BEARING _____

ULTIMATE DEPTH 379

ELEVATION 4006.84 +2.95 1360.2 m

DIP Vertical

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE		
	Recovery						Pb	Zn		
0-10 ³	Casing (overburden)									
10-60	14.0 Fragments of medium greenish-grey sericite schist. Some chlorite in association with about 20% white quartz. Shearing at 60° to core.									
60-65	2.1 Graphitic schist fragments up to 0.1. One 1/16" thread of Zns (reddish-brown) at 62.2. Sparse pyrite. Not worthy of assay.									
65-69	1.6 Graphitic schist with about 50% grey silicification. About 10% pyrite. Several threads of reddish-brown Zns and a little associated Pbs.	4646	65	69	4		2.70	5.80		
69-80	3.9 Graphitic schist silicified as above. Sparse pyrite.									
80-84	2.9 Graphitic schist, silicified. 20% pyrite. Sparse visible Pbs and Zns.	4647	80	84	4		0.90	0.69		
84-88	4.0 Graphitic schist with 40% greyish-white silicification. 25% pyrite. A little chalco pyrite and pyrrhotite. No visible Pbs or Zns.	4648	84	88	4		0.37	0.19		
88-92	4.0 Graphitic schist, silicified as in previous section.	4649	88	92	4		0.11	0.39		
92-96.5	4.2 As above with about 20% pyrite. Sparse Pbs and Zns.	4650	9.2	96.5	4.5		0.39	0.59		
96.5-101	4.5 Graphitic schist highly silicified with grey quartz. 30% pyrite. No visible Pbs or Zns.	4713	96.5	101	4.5		0.39	0.59		

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE		
	Recovery						Pb	Zn		
101-105	3.2 Graphitic schist as above. Shearing contorted. 15% pyrite.	4714	101	105	4		0.42	0.98		
105-107	2.0 Same as above.	4715	105	107	2		0.51	0.59		
107-112	5.0 20% pyrite in silicified graphitic schist.	4716	107	112	5		2.26	2.36		
112-128	10.5 Graphitic schist, containing about 30% silicification. Less than 5% pyrite.		112	128	16					
128-130	1.1 0.1/4 of massive pyrite. Several threads of reddish brown Zns in graphitic schist.	4717	128	130	2		3.98	6.68		
130-134	2.0 Graphitic schist replaced by 30% white quartz.				4		0	0		
134-137	2.8 Graphitic schist; 30% silicification; 20% pyrite. Shearing at 60° to core.	4718	134	137	3		1.89	2.26		
137-141	3.4 Graphitic schist changing to white sericite schist at 139'-0. 30% pyrite some Pbs and Zns.	4719	137	141	4		2.03	2.26		
141-144.6	3.6 Massive pyrite and pyrrhotite containing a little finely associated Pbs and Zns.	4720	141	144.6	3.6		7.96	5.80		
144.6-154	8.2 Greyish-white sericite schist. 30% white quartz. Shearing at 40° to core.				7.9		0	0		
	152.5-154 0.1/4 of pyrite, Pbs and Zns in grey gangue at 153.1	4721	152.5	154	1.5		3.27	7.37		
154-164.9	9.4 Greyish-white sericite schist. Soft and fissile. Shearing at 80° to core.		154	164.9	10.9		0	0		

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE		
	Recovery						Pb	Zn		
164.9-166	1.1 Massive pyrite. Some closely associated Pbs and Zns.	4722	164.9	166	1.1	} 5.0				
166-169.5	3.9 Massive with about 20% to 30% combined Pbs and Zns. Some fine siliceous gangue. Relict banding at 50° to core.	4723	166	169.9	3.9		3.08	5.90		
169.9-171	1.1 Greyish-white sericite schist. Sparse pyrite.				1.1		0	0		
171-176	5.0 Massive pyrite with 20% fine Pbs and Zns. Some closely associated siliceous gangue.	4724	171	176	5		1.49	5.51		
176-179	2.8 40% pyrite, 25% combined Pbs and Zns. Balance siliceous gangue.	4725	176	179	3		3.61	7.68		
179-184	5.0 55% pyrite, 25% combined Pbs and Zns, a little chalcopryrite and pyrrhotite. Balance siliceous gangue, with vestiges of sericite schist.	4726	179	184	5		3.78	5.61		
184-187	2.9 Massive sulphide consisting of 60% pyrite, 30% combined Pbs and Zns. Balance light grey silica.	4727	184	187	3		4.05	6.88		
187-192.5	5.0 Same type of mineralization as in previous section. Ratio of Zns to Pbs about 3 to 2.	4728	187	192.5	5.5		3.41	5.12		
192.5-198	5.2 Massive sulphide consisting of 70% pyrite, 20% combined Pbs and Zns, 10% siliceous gangue. Some sections magnetic, indicating presence of a little pyrrhotite.	4729	192.5	198	5.5		1.07	1.08		

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 4 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery								
198-203	5.0 As on previous page, but showing an increase of Pbs and Zns over the pyrite possibly 25% combined.	4730	198	203	5		3.28	3.44	
203-208	4.7 As in previous section but with massive pyrrhotite from 205 to 206.5, lessening overall percentage of Pbs and Zns.	4731	203	208	5		5.67	9.15	
208-213	4.8 Massive sulphide consisting of pyrite pyrrhotite in minor amount and about 25% combined Pbs and Zns. Matrix of about 15% silica. Vestigial banding at 50° to core.	4732	208	213	5		3.13	6.78	
213-217	4.0 Massive sulphide as above. Slight increase in silica gangue content.	4733	213	217	4		1.92	6.88	
217-222	5.0 Massive sulphide as in previous section but with vestiges of white sericite schist from 218 to 221.	4734	217	222	5		2.17	4.23	
222-227	5.0 Massive mineralization consisting of pyrite, minor pyrrhotite, 20 to 30% combined Pbs and Zns and about 20% siliceous gangue and achist remnants.	4735	222	227	5		3.44	5.61	
227-231	4.0 As in previous section but with about 15% siliceous gangue.	4736	227	231	4		2.82	7.18	
231-236	5.0 Massive sulphide as in previous sections but with several irregular white quartz patches from 231 to 233. Sulphide assemblage fine grained.	4737	231	236	5		1.75	5.71	

71.7

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 5 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE		
	Recovery						Pb	Zn		
236-241	5.0 Massive sulphide containing about 15 to 20% combined Zns and Pbs. Increase of pyrrhotite relative to the pyrite indicated magnetically.	4738	236	241	71-1 5		3.41	2.66		
241-246	5.0 Pyrite, pyrrhotite and about 15% Pbs and Zns. A little chalc pyrite. One 0.1 section of relatively coarse Pbs. About 30% siliceous gangue.	4739	241	246	5		2.56	1.08		
246-251	5.0 60% sulphide consisting of pyrite, Pbs and Zns (15%). Balance grey siliceous gangue.	4740	246	251	5		0.45	1.67		
251-256	5.0 Massive sulphide consisting of 75% pyrite and pyrrhotite, 20% combined Pbs and Zns and balance siliceous gangue.	4741	251	256	5		0.48	0.89		
256-261	5.0 Massive pyrite with about 30% combined Pbs and Zns.	4742	256	261	5		9.51	1.57		
261-266	5.0 Massive sulphide as above. Pyrrhotite present with the pyrite.	4743	261	266	5		0.37	0.59		
266-271	5.0 Increase of siliceous gangue to about 30% sulphide ratios as in previous sections.	4744	266	271	5		0.54	1.38		
271-276	5.0 50% pyrite, 15% pyrrhotite, sparre chalcopyrite, 15% combined Pbs and Zns. Balance siliceous gangue.	4745	271	276	5		0.71	1.18		
276-281	5.0 As above. Zns present in greater amount than Pbs. Brecciated and healed with pyrite at 277, along with greyish-black soft material. (Gouge)	4746	276	281	5		0.62	0.69		

115-1

DRILLED BY _____

SIGNED _____

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 6 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
	Recovery						Pb 27		
281-286	5.0 30% pyrite and pyrrhotite, 15% Zns and 5% Pbs. Balance rey siliceous gangue. Sericite schist at 285.5.	4747	281	286	115-1 5		3.96 5.41		
286-325	Medium greenish-grey sericite schist containing about 30% irregular white quartz patches. Shearin at 60° to 80° to core. Some chlorite in association with the quartz. Sparse pyrrhotite, and chalcovrite.								
325-335	7.0 Medium rey sericite schist, platy and soft. Shearing at 70° to 80° to core. A little pyrrhotite, not worthy of assay.								
335-344	5.4 As in previous section but with more pyrrhotite present.								
	338.3 to 340.7 About 40% pyrrhotite in irregular patches up to 0.13. Sample taken for Ni only.	4699	338.3	340.7	2.4				
344-352	3.5 Medium grey sericite schist. A little pyrrhotite.								
352-357	2.6 White quartz containin remnants of sericite schist.								
357-379	18.3 Medium grey sericite schist. Quite fissile at 80° to core. 10% white quartz.								

120.1

318.17 494.39

DRILLED BY _____

SIGNED _____

32.65
412
COMBINED OLD 120.1

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SHEET NUMBER 7

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO	FROM	TO	WIDTH	CORE SAMPLE			SLUDGE SAMPLE			
	The last section of sericite schist is similar to that obtained below the mineralized horizon in Hole 10.											
	END OF HOLE AT 379											
	RECOVERY : 67%											

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 15

PROPERTY Vangorda

SLUDGE RECORD

SHEET NUMBER 1 Sludges SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	FORMATION	SAMPLE NO.	FROM	TO	WIDTH	CORE SAMPLE		SLUDGE SAMPLE	
10-60	Light grey. Numerous sericite flakes. No evidence of sulphide.								
60-62	Graphitic. Low core recovery in section containing a little Zns.	4697	60	62	2				
62-67	As above.	4698	62	67	5				
67-141	Graphitic schist indicated from sludges. Core recovery good in mineralized portions. Some sludges lighter in colour due to silicification								
141-193	Light grey sludges. Core recovery good in this section. Water return lost at 193'. Core recovery good over this section.								

DRILLED BY Boyles Bros. Ltd.

SIGNED R. W. Baker

DIAMOND DRILL RECORD,

HOLE NO. 16

PROPERTY VANGUARD CREEK Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 161.6 STARTED March 28, 1954
 LATITUDE 30,073.41 9166.37 DATUM 4000 COMPLETED April 3, 1954
 DEPARTURE 30,557.37 9313.88 BEARING _____ ULTIMATE DEPTH 416.0
 ELEVATION 4039.94 ^{+2.9'} 1310.31 DIP Vertical hole PROPOSED DEPTH To cross-section mineralized "A X" core

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
12.0	36	Casing (overburden)												
0-86.2	48.1	Medium grey to dark grey sericite schist. Pearly lustre on cleavage faces. Shearing at 70° to 80° to core. About 20% white quartz replacement, in irregular patches.												
0-102.0	12.6	Graphitic schist containing fine laminations of grey silica. Shearing at 70° to core. About 30% pyrite some reddish brown sphalerite visible in the laminations.												
		86.3 to 91.3 Noted above	4751	5.0			1.91	3.78						
		91.3 to 96.3 " "	4752	5.0			1.46	3.18						
		96.3 to 102.0 " "	4753	5.7			1.83	3.08						
2.0-1.6	46.2	Medium grey sericite schist. Greenish cast in part. Several small patches of pyrrhotite and pyrite. Fissile shearing at 80° to core.												
1.6-188.6	27.0	Medium grey to dark sericite schist. Greenish cast indicative of some associated chlorite. Pearly translucent lustre on shear planes. Shearing (fissile) at 80° to core. A few irregular threads and small patches of pyrrhotite and pyrite.			water return lost at 180'									
8.6-91.0	2.7	Graphitic schist. Gradational to a high carbon content at 191.0. Several irregular white quartz threads and some interbanded grey silica. A little pyrrhotite and pyrite.												
1.0-93.0	1.9	Graphitic schist containing grey interbanded silica 15% pyrite and a little sphalerite.	4754	2.0			1.21	2.79						

DIAMOND DRILL RECORD,

HOLE NO. L 16

PROPERTY VANGONDA

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
3.0- 96.0	2.5	Graphitic schist as above. 20% pyrite with a little sphalerite.	4755	3.0			1.80	3.78	✓					
96.0- 98.0	2.0	Graphitic schist. Shearing at 80° to core. Wavy banding containing grey silica at right angles to the shearing. Scattered fine pyrite; a little associated sphalerite.	4756	2.0			0.82	2.39	✓					
98.0- 03.0	5.0	Graphitic schist replaced by irregular pyrrhotite, sphalerite and pyrite. Sparse visible galena and chalcopyrite. Mineralization changes to a massive silicified type at 201.5 leaving no vestiges of original rock.	4757	5.0			6.46	5.97						
03.0- 08.0	5.0	About 60 to 70% fine grained sulphide consisting of pyrite, pyrrhotite, sphalerite and galena in a silica matrix. Sparse chalcopyrite.	4758	5.0			6.27	6.47	✓					
08.0- 13.0	5.0	Heavy sulphide mineralization similar to previous section. Appears to be less silica and a greater amount of pyrite and pyrrhotite present.	4759	5.0			3.51	2.79	✓					
13.0- 18.0	5.0	Finely associated pyrrhotite, pyrite, galena, sphalerite and chalcopyrite, with greater silica content and less pyrite than in previous section.	4760	5.0			3.65	3.78						
18.0- 23.0	5.0	Similar mineralization to above. Strong pyrrhotite content indicated from Alnico magnet.	4761	5.0			2.84	1.79	✓					
23.0- 28.0	5.0	Heavy fine sulphide and silica gangue as above	4762	5.0			1.77	1.39						

DIAMOND DRILL RECORD,

HOLE NO. 16'

PROPERTY VALECORDA

SHEET NUMBER 3 SECTION FROM TO STARTED
 LATITUDE DATUM COMPLETED
 DEPARTURE BEARING ULTIMATE DEPTH
 ELEVATION DIP PROPOSED DEPTH

EPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
28.0- 33.0	5.0	Greater silica gangue content in sulphide mineralisation similar to previous sections.	4763	5.0			1.85	2.69						
33.0- 38.0	5.0	Finely associated pyrrhotite, pyrite, galena and sphalerite in a grey silica gangue as in previous sections. Wisp of chlorite probably vestigial from replaced rock.	4764	5.0			1.07	1.49	✓					
38.0- 41.0	3.0	Light yellow fine pyrite, and fine silica matrix. A little visible sphalerite and galena in bands at 80° to core- probably guided by shearing, now replaced.	4765	3.0			2.92	1.19	✓					
41.0 - 46.0	5.0	Fine pyrite and silica in a ratio of about 60:40	4766	5.0			0.45	0.10						
46.0 - 51.0	5.0	Mineralisation similar to above. Sphalerite or galena not noted.	4767	5.0			0.42	2.79						
51.0 - 56.3	5.0	Mineralisation as above. Pyrite massive in some sections.	4768	5.0			0.28	0.90						
56.3- 63.5	2.8	Medium grey silicified schist. Compact, non-fissile due to silicification. Has a slight greenish cast and numerous white qtz. threads.												
63.5- 66.0	2.5	Fine pyrite in a siliceous matrix, in proportion of about 40:60. Sphalerite or galena not visible.	4769	2.5			0.42	0.99						
66.0 - 70.0	4.0	Same as previous mineralisation. Pyrite massive in some sections.	4770	4.0			0.34	2.19						

DIAMOND DRILL RECORD,

HOLE NO. 16

PROPERTY VANGORDA

SHEET NUMBER 4

SECTION FROM _____ TO _____

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOVER	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-4.0	4.0	Same as previous section with an increase of silica over the pyrite.	4771	4.0			0.37	0.99						
4.0-8.5	4.5	Fine pyrite admixed with gray silica. The pyrite is massive in part. Galena or sphalerite not seen.	4772	4.5			0.04	0.30						
8.5 - 1.9	3.4	Medium gray silicified schist. Relief shearing at 80° to core.												
1.9 - 5.3	2.8	Fine pyrrhotite and pyrite in a gray silica matrix. The sulphides total about 50% and are massive in parts of the section.	4773	3.4			0.03	0.10						
5.3-5.6	2.8	Same as previous section. Alnico magnet shows presence of pyrrhotite.	4774	3.3			0.22	0.20						
33.6-33.0	4.4	Heavy fine pyrite mineralization, about 60% of the sample. Other sulphides not seen.	4775	4.4			Tr	0.10						
33.0-38.0	5.0	Same as previous section. Section of graphitic schist from 296.5 to 297.5 with considerable gray silica.	4776	5.0			0.22	N11						
38.0-33.0	5.0	Heavy fine pyrite mineralization. Vestiges of graphitic schist between 301.5 and 302.5; Shearing at 70° to core.	4777	5.0			0.05	N11						
33.0-38.2	5.2	About 60% fine pyrite, chiefly in massive sections. Balance, vestiges of graphitic schist and inter-banded silica.	4778	5.0			0.28	0.99						

DIAMOND DRILL RECORD,

HOLE NO. 18

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 123.0

STARTED April 13, 1954

LATITUDE 29,646.34

DATUM 4000

COMPLETED April 18, 1954

DEPARTURE 30,846.62

BEARING _____

ULTIMATE DEPTH 319.0

ELEVATION 4,082.45

DIP Vertical hole

PROPOSED DEPTH to Test mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-79.0		Casing (overburden)												
79.0-98.0	11.5	Medium grey to dark grey sericite schist. About 20% white quartz. Shearing at 80° to core.												
98.0-103.0	2.4	Graphitic schist containing scattered fine pyrite and about 10% reddish-brown sphalerite. Some inter-banded grey silica.	4781	5.0			0.84	2.19						
103.0-108.0	5.0	Same as previous section, changing to massive fine pyrite at 104.0. Some fine galena present between 103.0 and 108.0	4782	5.0			7.20	6.47						
108.0-113.0	5.0	Massive fine pyrite. Presence of pyrrhotite indicated by magnet. Fine PBs present	4783	5.0			5.36	3.68						
113.0-118.0	5.0	Same as previous section. Some associated fine pyrrhotite and silica.	4784	5.0			0.48	1.59						
118.0-123.0	5.0	Interbanded pyrite, pyrrhotite and silica. Sulphide totals about 60%. Relict banding at 50° to core.	4785	5.0			0.03	0.50						
123.0-128.0	5.0	Same as previous section. No galena or sphalerite visible in the core. Water return lost at 123.0	4786	5.0			Tr	0.10						
128.0-133.0	5.0	Sections of massive and disseminated pyrite totalling about 60%, in a grey silica matrix.	4787	5.0			nil	nil						
133.0-138.0	5.0	Same mineralization as above.	4788	5.0			Trace	nil						

DIAMOND DRILL RECORD,

HOLE NO. 18

PROPERTY VANGORDA CREEK

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
38.0-43.0	5.0	Dominantly fine pyrite in a silica matrix. Some associated pyrrhotite.	4789	5.0		10	0.48	nil						
43.0-48.0	5.0	Same as previous section.	4790	5.0			0.05	nil						
48.0-53.0	5.0	Slight increase in silica relative to the pyrite.	4791	5.0			Trace	nil						
53.0-58.0	5.0	As above. Strong indications of pyrrhotite with the magnet.	4792	5.0			0.03	0.10						
58.0-63.0	5.0	Increase in pyrrhotite relative to the pyrite. Some chalcopyrite and fine galena and sphalerite present.	4793	5.0			1.35	4.48 ^{4.39}						
63.0-68.0	5.0	Dominantly pyrrhotite and silica with a little fine galena, sphalerite and chalcopyrite.	4794	5.0			3.94	4.48						
68.0-73.0	5.0	As above, but with a stronger galena and sphalerite content.	4795	5.0			2.81	2.79						
73.0-78.0	5.0	Same as previous section. Greenish gray schist and quartz from 173.0 to 174.0	4796	5.0			2.11	1.89						
78.0-83.0	5.0	About 40% siliceous matrix containing pyrrhotite, pyrite, sphalerite and galena. Sparse chalcopyrite.	4797	5.0			1.97	3.28						
83.0-88.0	5.0	Patches and threads of pyrrhotite and chalcopyrite in a silica matrix. Some associated pyrite. Sparse Pb and Zn.	4798	5.0			0.05	0.40						

DIAMOND DRILL RECORD,

HOLE NO. 18

PROPERTY VANGORDA CREEK

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
88.0-93.0	5.0	Same as previous section.	4799	5.0			0.03	0.10						
93.0-98.0	5.0	Slightly more pyrite than above. Relict banding at 60° to core.	4800	5.0			0.03	0.40						
98.0-103.0	5.0	Considerable fine reddish-brown sphalerite and galena between 200.0 and 202.0. Relict banding at 50°.	4801	5.0			2.78	2.59						
103.0-108.0	5.0	About 90% pyrite and pyrrhotite in a silica matrix.	4802	5.0			0.03	nil						
108.0-113.0	5.0	Irregular patches of pyrrhotite and pyrite.	4803	5.0			0.02	nil						
113.0-118.0	5.0	Same as above.	4804	5.0			Tr	nil						
118.0-123.0	5.0	Pyrite and pyrrhotite as above. Relict banding at 70° to core.	4805	5.0			0.03	nil						
123.0-128.0	5.0	Pyrite and pyrrhotite as in previous section. Remnant of carbonatized and silicified graphitic schist from 225.5 to 226.0.	4806	5.0			Trace	nil						
128.0-133.0	5.0	As above. Relict banding at 70° to 80° to core, followed by the pyrrhotite in some cases.	4807	5.0			0.17	0.50						
133.0-138.0	5.0	As above. Sparse chalcopyrite.	4808	5.0			0.28	0.50						

DIAMOND DRILL RECORD,

HOLE NO. 20

PROPERTY VANQORDA CREEK

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
204.0-208.8	3.0	Greyish-white sericite schist containing about 20% white quartz. This type has a bleached, calery-like colour. One 1" patch of pyrite, Pbs and Zns at 205.0	4828	4.8			1.46	4.18						
208.8-210.5	1.7	About 50% finely associated pyrite, galena and sphalerite. Balance sericite schist and quartz.	4813	1.7			12.15	10.83						
210.5-234.0	19.4	Medium grey to light grey sericite schist. Shearing at 70°. Clay-like and cut easily with knife. From 220.6 to 221.0 a patch of fine pyrite, Pbs and Zns; a similar section from 229.0 to 229.5. These sections in themselves are not worthy of assay.	4886 4887 4888 4889 4890	210.5-215 4.5' 215-220 5.0' 220-225 5.0' 225-230 5.0' 230-234 4.0'			tr	0.49						
234.0-239.0	2.3	One foot of massive fine pyrite, galena and sphalerite walled on both sides by fragments of sericite schist, quartz and carbonate. Core loss has taken place in the schist.	4814	5.0 ✓										
239.0-246.0	7.0	Pale grey sericite schist. About 20% white quartz in four irregular patches. Shearing at 50° to core.	4891 4892	239-242 3.0' 242-246 4.0'			nil	nil						
246.0-250.0	4.0	Fine, compact massive sulphide consisting of pyrite pyrrhotite and silica. Galena and sphalerite not recognized in the section.	4815	4.0										
250.0-253.4	3.4	Massive fine pyrrhotite and pyrite containing fine threads of reddish-brown sphalerite.	4816	3.4										
253.4-260.5	7.1	Light grey sericite schist. Shearing at 70° to core. Graphitic interbanding from 257.2 to 258.0	4893	7.1			nil	0.54						

DIAMOND DRILL RECORD,

HOLE NO. 20

PROPERTY VANGORDA CREEK

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
260.5- 261.9	1.4	Fine galena and sphalerite in a pyrite-pyrrhotite-silica gangue.	4817	1.4			3.23	6.55						
261.9- 274.5	10.8	Graphitic schist containing short irregular sections of sericite schist. Shearing at 70° to core.												
274.5- 303.0	27.1	Light grey to medium grey sericite schist. Shearing at 80° to core. The rock has a clay-like feel. About 2% fine pyrrhotite.												
303.0- 321.8	15.3	Medium grey to light grey sericite schist. Some chloritic bands and about 5% irregular pyrrhotite, in small patches and also following the banding.												
321.8- 327.0	3.8	Graphitic schist. A little interbanded sericite schist. Sparse pyrrhotite and Zns.												
327.0- 332.0	5.0	Graphitic schist with about 30% gray silica. Average shearing to core about 80°. Pyrite, pyrrhotite and reddish-brown sphalerite totals about 30%.	4818	5.0			1.56	3.97						
332.0- 334.1	2.1	Graphitic schist containing about 15% pyrite and some reddish-brown sphalerite.	4819	2.1			1.14	3.97						
334.1- 338.7	4.6	Light grey sericite schist. Shearing at 80° to core. Two small patches of pyrite with a little galena and sphalerite. Not worthy of assay.		4.6										
338.7- 43.7	5.0	Three patches of pyrite and pyrrhotite containing visible galena and sphalerite in sericite schist. Each section about 1 ft. long.	4820	4 5.0			2.79	4.97						

DIAMOND DRILL RECORD,

HOLE NO. 21

PROPERTY VANGORDA CREEK, Y.T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 203.5

STARTED April 27, 1954

LATITUDE 29,449.97 8976.35

DATUM 400'

COMPLETED May 3, 1954

DEPARTURE 29,970.96 9135.15

BEARING _____

ULTIMATE DEPTH 367.0

ELEVATION 4,049.71⁺²⁵⁹ 1313.29

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
0.0- 118.0		Overburden (casing) Large granite boulder drilled through immediately before bedrock.												
118.0- 151.0	7.5	Graphitic schist. Mainly fragments containing patches of white quartz from 136.0 to 151.0. No evidence of sulphide mineralization. Shearing at 10° to core, changing 50° to core toward 151.0.												
151.0- 198.5	13.8	Sericite schist. Light grey to greenish grey. Talcose from 186.0 to 198.5.												
198.5- 203.5	5.0	From 198.5 to 200.0, graphitic containing pyrite - silica banding at 40° to core. At 200.0 mineralization changes to massive pyrite-pyrrhotite and silica. Sample taken from 198.5 to 203.5	4822	5.0	water return lost at 200'		3.25	2.99						
203.5p 208.5	5.0	Heavy pyrite and pyrrhotite in a silica matrix. Relict banding at 40° to core.	4823	5.0			2.99	2.79						
208.5- 213.5	5.0	As in the previous section. Some fine galena and sphalerite visible. Sparse chalcopyrite.	4824	5.0			3.57	3.68						
213.5- 227.2	1.8	Graphitic schist, in fragments. Shearing at 50° to core.												
227.2- 231.0	1.6	Fragments of sericite schist containing pyrrhotite and pyrite. A little associated graphitic schist.	4825	3.8			2.04	2.09						
231.0- 236.0	5.0	Mainly pyrrhotite and pyrite in a silica matrix. Some galena and sphalerite present.	4826	5.0			2.16	2.69						

DIAMOND DRILL RECORD,

HOLE NO. 26

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 117.0
 LATITUDE 30,141.85 9187.23 DATUM 4000
 DEPARTURE 30,171.52 9196.28 BEARING _____
 ELEVATION 4039.33 ⁺²¹⁹ 1310.13m DIP Vertical hole

STARTED May 15, 1954
 COMPLETED May 18, 1954
 ULTIMATE DEPTH 351.0
 PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-23.0	?	Casing (overburden)												
23.0-01.0	24.0	Medium to dark grey sericite schist. Shearing at 80° to core. Considerable white quartz from 46 to 57.0. Core badly buttoned.												
01.0-07.0	3.1	Graphitic schist. Several small threads reddish-brown sphalerite. A little pyrite. Not worthy of assay. Shearing at 80° to core.												
07.0-12.0	5.0	Graphitic schist containing some reddish-brown sphalerite associated with fine pyrite and silica.	4829	5.0			1.46	3.88						
12.0-17.0	5.0	Graphitic schist with fine pyrite. A little visible sphalerite.	4830	5.0			0.52	1.09						
17.0-22.0	5.0	Graphitic schist containing considerable fine pyrite and grey interbanded silica. Sphalerite and galena present. Banding variable, average 80° to core.	4831	5.0			4.66	8.56						
22.0-27.0	5.0	Same as previous section with less galena and sphalerite.	4832	5.0			1.39	3.28						
27.0-32.0	5.0	Graphitic schist with about 50% irregular silica and pyrite.	4833	5.0			0.12	0.60						
32.0-37.0	5.0	Same as previous section. Irregular wavy banding.	4834	5.0			0.27	0.50						
37.0-42.0	5.0	Same as previous section with an increase of pyrite relative to the silica.	4835	5.0			0.07	0.10						

DIAMOND DRILL RECORD,

HOLE NO. 26

PROPERTY VANGORDA CREEK, T. T.

SHEET NUMBER 2 SECTION FROM _____ TO _____ STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

191.2 - 20
 203.7 - 20

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
142.0-185.4	38.6	Graphitic schist with about 40% interbanded silica. A little pyrite and pyrrhotite. Shearing at 80° to core.												
185.4-191.2	5.8	Massive sulphides consisting of pyrite pyrrhotite galena and sphalerite. A little silica in the matrix.	4836	5.8			5.12	8.46						
191.2-203.7	8.0	Graphitic schist with interbanded silica. Shearing at 80° to core.	5089 5090 5091 5092	191.2-195 195-200 200-205 205-210	3.8 5.0 5.0 5.0		0.05 0.02 tr tr	0.69 0.10 ml ml						
203.7-215.6	7.4	Medium to light grey sericite schist. Numerous quartz-carbonate threads. Shearing obscured.	5093	210-215	5.6		tr	ml						
215.6-220.0	3.0	Massive sulphides consisting of pyrite, galena and sphalerite.	4837	5.0			4.64	5.07						
220.0-225.0	3.7	Same as previous section. A little increase in silica matrix.	4838	5.0			3.92	6.37						
225.0-230.0	5.0	Same as previous section with some pyrrhotite in last 0.5.	4839	5.0			3.25	4.28						
230.0-235.0	5.0	Increasing pyrrhotite relative to the pyrite. Considerable sphalerite and galena.	4840	5.0			4.82	5.97						
235.0-240.0	5.0	Less pyrrhotite relative to the pyrite in the massive sulphides. Some silica in the matrix. Several small patches milky quartz.	4841	5.0			2.26	6.37						
240.0-245.0	5.0	Massive sulphide as above containing 10 to 15% white quartz-carbonate. Sparse pyrrhotite.	4842	5.0			1.14	2.59						

DIAMOND DRILL RECORD,

HOLE NO. : 26

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 3 SECTION FROM _____ TO _____ STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
45.0-250.0	5.0	Same as previous section, Vestigial banding at 70° to core. Remnant of carbonatized schist from 247.0 to 248.5.	4843	5.0			1.96	6.07						
250.0-255.0	5.0	Same as previous section with increased amount of silica. Pyrrhotite present in this and previous section.	4844	5.0			2.68	4.88						
255.0-260.0	5.0	Massive sulphides consisting of pyrite, galena and sphalerite. Several irregular patches of white quartz-carbonate.	4845	5.0			0.82	1.89						
260.0-265.0	5.0	Massive sulphides as in previous section. Increased quartz-carbonate content. About 0.5 carbonatized schist at 261.5.	4846	5.0			0.79	1.19						
265.0-270.0	5.0	Prominent galena and sphalerite associated with pyrite. Horne of carbonatized and silicified schist from 267.7 to 269.0.	4847	5.0			2.28	3.98						
270.0-275.0	5.0	About 30% irregular quartz and carbonate in pyrite, sphalerite and galena.	4848	5.0			1.02	0.98						
275.0-280.0	5.0	Remnants of sericite schist containing pyrite, quartz-carbonate and some galena and sphalerite.	4849	5.0			0.51	0.10						
280.0-285.0	5.0	Irregular pyrite, pyrrhotite with some galena, sphalerite. Highly silicified and carbonatized.	4850	5.0			0.22	0.39						
285.0-290.0	5.0	Medium grey silicified sericite schist replaced by about 30% irregular pyrrhotite. A little galena and sphalerite present.	4851	5.0			0.04	0.39						

DIAMOND DRILL RECORD,

HOLE NO. 27

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 240.0
 LATITUDE 30,272.25 9226.98N DATUM 4000
 DEPARTURE 30,322.28 9242.23E BEARING _____
 ELEVATION 4,060.25 4259' 1316.51m DIP Vertical Hole

STARTED May 18, 1954
 COMPLETED May 22, 1954
 ULTIMATE DEPTH 448.0
 PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-20.0		Casing (overburden)												
20.0-121.7	51.7	Light grey to medium grey sericite schist. About 15% irregular white quartz patches. Shearing at 50° to core. Between 105.0 and 116.0 there is a little interbanded graphitic material containing sparse reddish-brown sphalerite - not worth of assay.												
121.7-174.0	45.4	Medium grey to dark grey sericite schist. Some interbanded graphitic sections at 159.0, 170.0 and 173.5												
174.0-220.5	42.5	Light greenish-grey sericite schist. Shearing at 40° to core. Sparse pyrrhotite between 209.0 and 210.0.												
220.5-225.0	5.0	Graphitic schist with about 40% interbanded grey silica. Shearing at 60° to core but distorted in part. Some reddish-brown sphalerite, pyrite and pyrrhotite.	4852	5.0			2.26	4.43						
225.0-230.0	5.0	Graphitic schist with mineralization similar to previous section, with no pyrrhotite noted.	4853	5.0			2.11	4.33						
230.0-235.0	4.0	Graphitic schist containing irregular patches of pyrite, sphalerite and galena.	4854	5.0			2.64	4.43						
235.0-240.0	5.0	Graphitic schist with considerable interbanded grey silica and pyrite. Shearing at 70° to core. Sparse sphalerite.	4855	5.0			0.91	1.87						

DIAMOND DRILL RECORD,

HOLE NO. 27

PROPERTY VANGORDA CREEK, I. T.

SHEET NUMBER 2 SECTION FROM 240.0 TO 283.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
240.0-245.0	5.0	Sericite schist containing several patches of fine pyrite, sphalerite and galena.	4856	5.0			2.46	2.36						
245.0-250.0	5.0	Massive sulphide consisting of pyrite pyrrhotite galena and sphalerite in a silica matrix. Fragments of grey sericite schist from 245.0 to 246.0 and vestiges from 247.0 to 248.5.	4857	5.0			3.98	6.10						
250.0-255.0	5.0	Massive sulphide as above with remnant of sericite schist from 252.5 to 254.5. Shearing at 80° to core.	4858	5.0			2.97	6.98						
255.0-260.0	5.0	Massive fine pyrite pyrrhotite galena and sphalerite in a silica matrix.	4859	5.0			3.04	8.27						
260.0-263.0	3.0	As in previous section with some vestiges of sericite. Relict banding at varying steep angles to core.	4860	3.0			3.36	8.07						
263.0-266.7	3.7	Strong massive sulphides. Relict banding at 10° to 20° to core.	4861	3.7			5.84	9.55						
266.7-273.0	6.3	Light grey sericite schist. Variable shearing. Some quartz carbonate stringers. One 1" patch of pyrite galena and sphalerite at 269.2 and at 271.6, these not worthy of assay.	5094	6.3			0.02	nil						
273.0-278.0	5.0	Graphitic schist containing rice-sized grains of interbanded silica. Shearing at 60° to core. Pyrite galena and sphalerite present.	4862	5.0			1.23	2.46						
278.0-283.0	5.0	Graphitic schist containing silica grains as in previous section. Considerable fine pyrite and a little visible sphalerite.	4863	5.0			1.58	4.04						

DIAMOND DRILL RECORD,

HOLE NO. 29

PROPERTY VANGUARD CREEK, I. T.

SHEET NUMBER 1
 LATITUDE 30,550.81 93° 11.88' N
 DEPARTURE 30,347.94 9250.05 E
 ELEVATION 4,106.41 1330.57

SECTION FROM 0.0 TO 424.0
 DATUM 4000
 BEARING _____
 DIP Vertical hole

STARTED May 23, 1954
 COMPLETED May 28, 1954
 ULTIMATE DEPTH 502.0 (153) 1177.57 m
 PROPOSED DEPTH To test extent of mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
¹⁵⁻⁸⁷ 0.0-52.0		Casing (overburden)												
52.0-210.0	105.0	Medium gray to dark gray sericite schist. About 10% white quartz in irregular patches. A few small threads and patches of pyrrhotite. Shearing at an average of 70° to core, but contorted in part.												
210.0-249.0	39.0	Light gray sericite schist. 3 to 5% pyrrhotite in irregular patches and threads. Shearing at 60° to core.												
249.0-370.0	99.2	Medium gray to dark gray sericite schist. Shearing at 70° to core. Some interbedded graphitic material from 274.8 to 277.0. A little pyrrhotite.												
370.0-401.5	24.2	Light gray to medium gray sericite schist increasing chlorite and talc toward 401.5. Shearing at 70° to 80° to core. 3 to 5% fine pyrrhotite.												
401.5-403.0	1.5	Irregular galena and reddish-brown sphalerite in talc-sericite schist. Some finely associated magnetite and numerous small subhedral garnets. Some green chlorite.	4894	1.5			0.41	0.98						
403.0-419.5	9.5	Light gray talc-sericite schist. Some threads and small patches of pyrrhotite.												
419.5-424.0	4.5	Medium gray talc-sericite schist containing a few seams of reddish-brown sphalerite, galena and pyrrhotite. Numerous small garnets in proximity to the sulphide.	4895	4.5			0.02	0.49						

DIAMOND DRILL RECORD,

HOLE NO. 30

PROPERTY YANCOODA CREEK, Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 198.0
 LATITUDE 30,423.73 9273.15N DATUM IND
 DEPARTURE 30,196.42 9203.84E BEARING _____
 ELEVATION 4,075.76 1321.2 DIP Vertical hole

STARTED May 25, 1954
 COMPLETED May 30, 1954
 ULTIMATE DEPTH 423.0 (128.93) 1192.27m
 PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-52.0		Casing (overburden)												
52.0-125.7	49.1	Medium to dark gray sericite schist. Shearing at 70° to core. Several small white quartz patches. Minor interbanded chlorite and pyrrhotite.												
125.7-142.0	15.9	Light gray sericite schist. A few stringers of pyrrhotite and several patches of white quartz. Shearing at 50° to core.												
142.0-147.0	5.0	Graphitic schist, highly fractured and healed by white quartz and carbonate. Some disseminated pyrite sphalerite and galena.	4897	5.0			1.99	4.03		9.95	20.15			
147.0-178.0	25.2	Medium gray sericite schist. Shearing at 60° but contorted in part. Several threads of white quartz carbonate.												
178.0-183.0	5.0	Graphitic schist, fractured, in part, and healed with a little fine pyrite. Sparse galena and sphalerite.	4898	5.0			1.14	4.26		5.7	5.7			
183.0-188.0	2.5	Graphitic schist with some interbanded gray silica. Pyrite galena and sphalerite present.	4899	5.0			2.28	4.38		11.4	21.65			
188.0-193.0	2.5	Same as previous section - graphitic host.	4900	5.0			2.34	4.33		11.7	21.66			
193.0-198.0	3.5	Graphitic schist, highly fractured and healed by white quartz threads, galena and sphalerite. White quartz later than the sulphide, following fractures.	4901	5.0			2.46	4.82		12.3	24.10			

DIAMOND DRILL RECORD,

HOLE NO. 30

PROPERTY VANGORDA CREEK, I. T.

SHEET NUMBER 2 SECTION FROM 198.0 TO 242.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS				
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.	
198.0-203.0	4.5	Graphitic schist containing mineralization similar to previous section but slightly increased in amount, relative to the schist.	4902	5.0			3.50	5.02		17.5	25.1				
203.0-206.0	3.0	Graphitic schist containing mineralization similar to previous section up to 205.0; schist is almost wholly replaced by massive pyrite with galena and sphalerite from 205.0 to 206.0.	4903	3.0			5.15	7.18		25.75	35.9				
206.0-209.4	3.4	Massive fine pyrite with galena and sphalerite containing vestiges of graphitic with some white quartz stringers.	4904	3.4 31.4			4.57	6.39		22.85 107.2	31.95 166.06				
209.4-222.6	9.4	Light gray sericite schist. Shearing at 50° to zone. One 0.4' section of pyrite, galena and sphalerite at 219.6, not worthy of assay.		13								3.41	5.29		31.4
222.6-227.0	4.4	Massive sulphides consisting of pyrite, galena and sphalerite. Some fine associated silica. Vestiges of sericite schist from 222.6 to 225.0.	4905	4.4			3.74	4.62		18.70	23.1				
227.0-232.0	5.0	Massive fine sulphides as above. Horae of carbonatized greenish-grey schist from 229.4 to 230.4.	4906	5.0			2.72	6.49		13.6	32.45				
232.0-237.0	5.0	Massive fine pyrite, galena and sphalerite in a silica matrix. Vestiges of sericite schist present. Halict banding at 70° to core.	4907	5.0			1.32	7.08		6.6	35.4				
237.0-242.0	5.0	Same as previous section with a little pyrrhotite present.	4908	5.0			2.55	3.84		12.75	19.20				

DIAMOND DRILL RECORD,

HOLE NO. 30

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 3 SECTION FROM 242.0 TO 423.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
242.0- 247.0	5.0	Irregular patches of fine pyrite in a matrix of silica and carbonate. Some vestiges of graphitic schist.	4909	5.0			0.82	3.84		4.1	19.2	2.28	5.30	
									55.75	129.35		24.4		
247.0- 248.8	1.8	Same as previous section. Galena or sphalerite not seen.	4910	1.8			0.91	1.67						
248.8- 253.0	4.2	Graphitic schist. Shearing contorted. Brittle and badly broken. A little white quartz-carbonate.												
253.0- 263.0	5.2	Pale gray sericite schist. Bleached, silicified appearance. A little associated graphitic schist at 254.0. Shearing obscured by alteration.												
263.0- 280.0	7.7	Graphitic schist. Shearing contorted. Brittle and core badly broken. Some fine stringers of quartz-carbonate.												
280.0- 293.0	4.0	Graphitic schist. Core badly broken. Considerable white quartz and some associated carbonate.												
293.0- 423.0	77.8	Graphitic schist. Shearing at 40° to core. Some interbanded grey silica. A few quartz-carbonate threads and small patches.												
		<u>End of hole at 423.0</u>												
		Recovery: 67.15												
		Size of core: A X (1 3/16") to 293.0												
		E X (7/8") to 423.0												
		Recovery in mineralized sections: 88.8%												

DIAMOND DRILL RECORD,

HOLE NO. 1

PROPERTY VAN CORDA CREEK, Y. T.

SHEET NUMBER 2

SECTION FROM 146.0 TO 221.0 End

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
146.0-151.0	0.2	One button of quartz and three fragments of grey sericite schist.												
151.0-155.0	0.3	Three fragments greenish-gray sericite schist. From 131.0 to 155.0 the lack of core points to a gouge fault some condition.												
155.0-160.0	1.2	Pale greenish-gray sericite schist. Shearing highly contorted. One 1/2" piece containing some pyrite at 160.0.												
160.0-165.0	5.0	Pale greenish-gray sericite schist. Shearing contorted and crushed and healed by quartz and carbonate. Some fine pyrite and sparse galena and sphalerite present.	4914	5.0			0.26	0.20						
165.0-170.0	3.8	Sericite schist, mineralized as in previous section.	4915	5.0			0.04	0.05						
170.0-182.0	9.0	Pale greenish-gray contorted sericite schist as in previous sections. Some white quartz and carbonate, sparse pyrite.												
182.0-221.0	8.0	Graphitic schist. Badly broken; Shearing, in part at 30° to core. A little fine pyrite and quartz-carbonate. Some interbanded grey silica. Change made from A X core to E X core at 201.0. Hole ends at 221.0, abandoned due to caving. Recovery 40.3%.												
R. W. Baker														

APPENDIX TO D.D. HOLE # 32

DESCRIPTION OF SLUDGES IN SECTIONS OF LOW CORE RECOVERY

- 75 - 80: Black graphitic schist with grains of silica, pyrite and reddish-brown iron oxide.
- 80 - 85: Similar to previous section; less pyrite, more flakey.
- 85 - 95: Same as previous section; a little pyrite.
- 95 - 100: Dark grey flakes, considerable silica, little pyrite, some reddish iron oxide.
- 100 - 110: Dark grey flakes with an increasing amount of pyrite.
- 110 - 120: Dark grey flakes, less pyrite.
- 120 - 146: Dark grey flakes, possibly sericite, a little silica and decreasing pyrite.

V. Papezik

DIAMOND DRILL RECORD,

HOLE NO. 33

PROPERTY VANCOUVER CREEK, T. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 115.0
 LATITUDE 29,833.17 9093.15N DATUM 4000
 DEPARTURE 30,426.64 9274.04E BEARING _____
 ELEVATION 4,020.33 1304.34 DIP Vertical hole

STARTED June 4, 1954
 COMPLETED June 7, 1954
 ULTIMATE DEPTH 407.0
 PROPOSED DEPTH To cross-section of formation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-65.0 (174)		Casing (overburden)												
65.0-75.0	2.6	Graphitic schist containing some intercrossed gray silica. Shearing at 50° to core. Sparse pyrite.												
75.0-80.0	3.0	Graphitic schist. Some fine pyrite and a little galena and sphalerite present.	4923	5.0			3.23	7.77						
80.0-85.0	2.1	Graphitic schist. Shearing at 30° to core. Considerable fine pyrite.	4924	5.0			3.35	5.32						
85.0-90.0	2.1	Graphitic schist with pyrite changing to massive fine pyrite with sphalerite and galena at 85.5. Relict banding at 35° to core.	4925	5.0			2.11	8.96						
90.0-95.0	4.4	Massive fine pyrite, galena and sphalerite in a silica matrix. Horse of silicified sericite schist from 92.0 to 94.0.	4926	5.0			1.62	4.43						
95.0-100.0	2.2	Porous fine pyrite containing some galena and sphalerite. Silica matrix.	4927	5.0			5.77	5.02						
100.0-105.0	3.5	Massive fine pyrite in a silica matrix. Galena and sphalerite present.	4928	5.0			1.76	6.30						
105.0-110.0	4.1	Same as previous section. Stronger galena and sphalerite content.	4929	5.0			4.04	9.05						
110.0-115.0	3.4	Massive fine pyrite, galena and sphalerite. A little white quartz.	4930	5.0			4.49	6.20						

DIAMOND DRILL RECORD,

HOLE NO. 2

PROPERTY VANGORDA CREEK, T. 2.

SHEET NUMBER 2 SECTION FROM 115.0 TO 165.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
115.0-120.0	5.0	previous section. with a little pyrrhotite present.	4931	5.0			3.69	4.72						
120.0-125.0	5.0	Heavy fine pyrite and pyrrhotite. Less sphalerite and galena than previous section.	4932	5.0			0.97	1.57						
125.0-130.0	5.0	Increased galena and sphalerite from 128.0 to 130.0 with heavy pyrite and pyrrhotite.	4933	5.0			1.84	4.03						
130.0-135.0	4.6	Heavy fine pyrite and pyrrhotite with strong galena and sphalerite in a silica matrix. Halict banding at 55° to core.	4934	5.0			1.96	7.77						
135.0-140.0	4.4	Same as previous section, but less pyrrhotite.	4935	5.0			3.46	7.68	✓					
140.0-145.0	5.0	Massive fine pyrite. Horres of silicified and carbonatized graphitic schist 141.0 to 143.2 and 143.6 to 144.5.	4936	5.0			1.70	4.33	✓					
145.0-150.0	5.0	Fine pyrite galena and sphalerite in a silica matrix. Vestige of graphitic schist at 147.0	4937	5.0			2.19	8.17	✓					
150.0-155.0	5.0	Massive sulphide as in previous section. Silicified and carbonatized sericite schist from 51.2 to 52.5	4938	5.0			2.19	4.23	✓					
155.0-160.0	5.0	Massive fine pyrite and prominent pyrrhotite with some galena and sphalerite in a silica matrix.	4939	5.0			3.23	4.13	4					
160.0-165.0	5.0	Massive sulphides as in previous section.	4940	5.0			4.27	6.30	5					

DIAMOND DRILL RECORD,

HOLE NO. 33

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 3

SECTION FROM 165.0 TO 215.0

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
165.0-170.0	5.0	Same as previous section with less pyrrhotite.	4941	5.0			4.22	8.06						
170.0-175.0	5.0	Massive pyrite, sphalerite and galena in a silica matrix.	4942	5.0			4.69	8.65						
175.0-180.0	5.0	Same as above with some associated pyrrhotite.	4943	5.0			2.31	6.39	-					
180.0-185.0	5.0	Massive fine sulphide as above. Helict banding at 60° to core.	4944	5.0			1.79	5.90	-					
185.0-190.0	5.0	Massive sulphides as in previous sections. Horse of silicified and carbonatized graphitic material from 189.4 to 190.0	4945	5.0			2.55	6.49						
190.0-195.0	5.0	Massive fine pyrite, pyrrhotite, galena and sphalerite in a silica matrix. Silicified and carbonatized remnant of sericite schist from 194.3 to 195.0	4946	5.0			4.57	3.54						
195.0-200.0	5.0	Massive fine pyrite. A little galena and sphalerite.	4947	5.0			0.02	0.10						
200.0-205.0	5.0	Increased silica matrix with massive fine pyrite.	4948	5.0			TR	nil						
205.0-210.0	5.0	Same as previous section with pyrrhotite present. Several blebs of sphalerite present.	4949	5.0			nil	nil						
210.0-215.0	5.0	Heavy fine pyrite and pyrrhotite in a silica matrix.	4950	5.0			TR	0.10						

DIAMOND DRILL RECORD,

HOLE NO. 39

PROPERTY VANERDA CREEK V V

SHEET NUMBER 4 SECTION FROM 215.0 TO 265.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
215.0-220.0	5.0	Same as previous section. Several small patches of galena and sphalerite.	4951	5.0			0.02	1.10						
220.0-225.0	5.0	Heavy fine pyrite in a silica matrix. Several small patches of galena and sphalerite.	4952	5.0			0.04	nil						
225.0-230.0	5.0	Massive fine pyrite and pyrrhotite with a little sphalerite and galena. Some interstitial gray silica.	4953	5.0			TR	nil						
230.0-235.0	5.0	Same as previous section. Some irregular white quartz.	4954	5.0			0.29	0.29						
235.0-240.0	5.0	Massive fine pyrite in a silica matrix. A little pyrrhotite.	4955	5.0			0.02	0.20						
240.0-245.0	5.0	Heavy fine pyrite and pyrrhotite. Heavy medium-grained massive galena from 244.5 to 245.0.	4956	5.0			6.30	0.49						
245.0-250.0	5.0	Massive fine pyrite and pyrrhotite with associated silica as a matrix.	4957	5.0			0.65	0.49						
250.0-255.0	5.0	Massive fine pyrite and pyrrhotite in a silica matrix.	4958	5.0			nil	nil						
255.0-260.0	5.0	Same as previous section.	4959	5.0			TR	0.10						
260.0-265.0	5.0	Increased amount of gray silica relative to massive fine pyrite and pyrrhotite.	4960	5.0			0.02	0.49						

DIAMOND DRILL RECORD,

HOLE NO. 33

PROPERTY VANDORA CREEK, T. T.

SHEET NUMBER 5 SECTION FROM 265.0 TO 310.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
265.0-270.0	5.0	A little interbanded reddish-brown sphalerite in vestiges of sericite schist. Heavy fine pyrite and and pyrrhotite present.	4961	5.0			1.58	2.65						
270.0-275.0	5.0	Same as previous section. Irregular patches of grey silica.	4962	5.0			0.76	1.97	✓					
275.0-280.0	5.0	Increased sphalerite content relative to pyrite and pyrrhotite.	4963	5.0			1.61	4.52	✓					
280.0-285.0	5.0	Heavy fine sphalerite and galena with pyrite and fine silica. Vestige of sericite schist from 284.0 to 285.0.	4964	5.0			1.96	7.67						
285.0-290.0	5.0	Considerable reddish-brown sphalerite with fine pyrite. Some associated galena. Vestiges of light grey sericite schist present.	4965	5.0			4.57	9.94	✓					
290.0-295.0	5.0	Same as previous section with about 50% vestigial sericite schist.	4966	5.0			1.61	3.54	✓					
295.0-300.0	5.0	Strong galena and sphalerite in association with pyrite. Vestiges of light grey sericite schist.	4967	5.0			3.49	4.82	✓					
300.0-305.0	5.0	Same as previous section.	4968	5.0			1.44	0.98						
305.0-310.0	5.0	Pyrite and pyrrhotite with vestiges of silicified sericite schist as in previous sections. Some galena and reddish-brown sphalerite present.	4969	5.0			0.56	1.38						

DIAMOND DRILL RECORD,

HOLE NO. 34

PROPERTY VASCONDA CREEK, I. T.

Alexander Y. Po
April, 1977

SHEET NUMBER 1

SECTION FROM 0.0 TO 186.0

STARTED June 5, 1954

LATITUDE 30,833.98 9398.19 N

DATUM 4000

COMPLETED June 8, 1954

DEPARTURE 30,369.09 9256.5 E

BEARING _____

ULTIMATE DEPTH 236.0

ELEVATION 4181.50 1353.46

DIP Vertical hole

PROPOSED DEPTH To test northeast of postulated fault

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS					
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.		
0.0-21.0		Casing (overburden)														
21.0-69.0	34.0	Medium to dark gray sericite schist. Some interbedded graphitic material. Shearing at 75° to core. A little white quartz-carbonate.														
69.0-75.0	3.9	Graphitic schist containing a little interbedded sericite schist.														
75.0-133.0	35.0	Dark gray sericite schist containing several short sections of interbedded graphitic schist. Shearing at 80° to core. Sparse pyrite and pyrrhotite.														
133.0-147.0	14.0	Graphitic schist containing 20% interbedded gray silice. A little interbedded dark gray sericite schist. Several patches of white quartz-carbonate.														
147.0-152.0	5.0	Massive, fine grained pyrite. No visible galena or sphalerite. A horse of badly broken graphitic schist from 150.0 to 151.0.	4976	5.0			.29	5.90								
152.0-157.0	5.0	Massive pyrite as in previous section. Scattered small patches of gray quartz.	4977	5.0			.23	2.16								
157.0-162.0	4.0	Massive pyrite as above.	4978	4.0			.20	.39								
162.0-186.0	6.5	Medium gray sericite schist with some interbedded graphitic schist. Considerable white quartz. No core from 166 to 170 - possible fault zone. Due to caving, change from 1 X to 2 X made at 186.0.														
									4980	160-165						
									4981	165-170						
									4982	170-175						
									4983	175-180						
									4984	180-185						
									4985	185-190						

DIAMOND DRILL RECORD,

HOLE NO. 34

PROPERTY VANDORDA CREEK, Y. T.

SHEET NUMBER 2 SECTION FROM 186.0 TO 256.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
186.0-237.0	22.9	Medium gray to dark gray sericite schist. Shearing at 70' to core. About 30% irregular white quartz-carbonate. Two buttons of white pyrite at 219.0.												
237.0-242.0	0	No core.												
242.0-248.0	1.1	Fragments of fine pyrite in a gray silica matrix. No visible galena or sphalerite.	4979	6.0			.12	nil	4986 4987 4988 4989	235-240 240-245 245-250 250-255				
248.0-256.0	0.1	Three small fragments of graphitic schist containing fine grains of pyrite.												
		Due to caving of the hole at 186.0 and 237.0, it was necessary to abandon it at 256.0. The above two sections possibly represent zones of shearing or faulting related to the postulated NW-SE fault in this section of ground.												
		End of hole at 256.0												
		Recovery for the hole: 56%												
		Recovery in mineralizations: 75.5%												
		R. W. Baker												

SLUDGE DESCRIPTIONS

D.D. HOLE 34

- 160 - 165 (Sample No. 4980) Greyish-black mixture of sericite flakes silica and graphitic material.
- 165 - 170 (Sample No. 4981) Medium grey mixture of sericite flakes and silica.
- 170 - 175 (Sample No. 4982) Same as previous section.
- 175 - 180 (Sample No. 4983) Light grey mixture of sericite flakes and silica.
- 180 - 185 (Sample No. 4984) Same as previous section.
- 185 - 190 (Sample No. 4985) Light greenish-grey, fine grained - possible gouge from fault zone - small sample due to poor water return.
- 235 - 240 (Sample No. 4986) Medium grey mixture of sericite flakes and silica.
- 240 - 245 (Sample No. 4987) Dark grey mixture of sericite flakes and graphitic material.
- 245 - 250 (Sample No. 4988) Medium grey sericite flakes and silica; fine pyrite grains throughout.
- 250 - 255 (Sample No. 4989) Same as previous section.

DIAMOND DRILL RECORD,

HOLE NO. 35

PROPERTY YANCOENA CREEK, T. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 158.0

STARTED June 9, 1934

LATITUDE 29,811.79

DATUM 4000

COMPLETED June 15, 1934

DEPARTURE 30,709.72

BEARING _____

ULTIMATE DEPTH 408.0

ELEVATION 4,058.19

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PS.	ZN.	NO.	FOOTAGE	AG.	CU.	PS.	ZN.
0.0-67.0		Casing (overburden)												
67.0-74.0	6.0	Graphitic schist with about 40% interbanded gray silica. Shearing at 80° to core. Some fine pyrite throughout. Sparse reddish brown sphalerite, not worthy of assay.												
74.0-128.0	3.0	Dark gray sericite schist containing some interbanded graphitic material. Shearing at 80° to 90° to core. A little disseminated pyrite.												
128.0-133.0	5.0	Graphitic schist containing some interbanded gray silica and sericite schist. Some pyrite and reddish-brown sphalerite present.	4991	5.0			0.91	2.06						
133.0-138.0	5.0	Graphitic schist with mineralization same as previous section.	4992	5.0			1.03	2.46						
138.0-143.0	5.0	Graphitic schist with interbanded silica and pyrite changing to massive fine pyrite with a little interbanded sericite schist at 140.0-143.0. A little galena and sphalerite present.	4993	5.0			3.98	4.72						
143.0-148.0	5.0	Massive fine pyrite with a little associated silica, galena and sphalerite.	4994	5.0			1.52	1.67						
148.0-153.0	5.0	Same as previous section with more silica present.	4995	5.0			0.03	0.79						
153.0-158.0	5.0	Massive sulphides consisting of pyrite, pyrrhotite, minor galena and sphalerite. Some silica in association.	4996	5.0			0.15	0.20						

DIAMOND DRILL RECORD,

HOLE NO. 35

PROPERTY VANGUARD CREEK, T. T.

SHEET NUMBER 2 SECTION FROM 198.0 TO 208.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
158.0-163.0	5.0	Pyrite, pyrrhotite in close association in a silica matrix. No visible galena or sphalerite.	4997	5.0			0.02	0.10						
163.0-168.0	5.0	Same as previous section.	4998	5.0			TR	1.10						
168.0-173.0	5.0	Increase of pyrrhotite relative to the pyrite in a silica matrix.	4999	5.0			nil	nil						
173.0-178.0	5.0	Mainly pyrrhotite with some associated pyrite in about 30% silica matrix. 70% SiO ₂ ?	5000	5.0			nil	nil						
178.0-183.0	5.0	Same as previous section.	5001	5.0			nil	nil						
183.0-188.0	5.0	Mainly fine pyrite and silica. A little associated pyrrhotite.	5002	5.0			0.03	nil						
188.0-193.0	5.0	Same as previous section. 20% SiO ₂ ?	5003	5.0			TR	nil						
193.0-198.0	5.0	Fine pyrite in association with about 40% grey silica. Several irregular small seams of magnetite. Sparse pyrrhotite.	5004	5.0			nil	nil						
198.0-203.0	5.0	Same as previous section. Possibly a little more magnetite.	5005	5.0			nil	nil						
203.0-208.0	5.0	Fine pyrite in association with grey silica. A little pyrrhotite and magnetite present.	5006	5.0			nil	nil						

DIAMOND DRILL RECORD,

HOLE NO. 38

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 185.0

STARTED June 16, 1954

LATITUDE 29,942.96

DATUM 4000

COMPLETED June 21, 1954

DEPARTURE 30,876.85

BEARING _____

ULTIMATE DEPTH 455.0

ELEVATION 4,058.75

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-47.0	(19.5)	Casing (overburden)												
47.0-70.5	10.5	Medium grey sericite schist. Shearing at 85° to core. Sparse pyrite. 0.8' of heavy fine pyrite and a little chalcopyrite from 59.1 to 59.4.												
70.5-152.0	58.0	Graphitic schist with about 20% interbanded silica. A little fine pyrite. Sparse pyrrhotite and reddish-brown sphalerite, the latter in fine threads.												
152.0-160.0	7.5	Graphitic schist. Shearing at 80° to core, but distorted in part.												
160.0-165.0	5.0	Graphitic schist containing disseminated pyrrhotite and reddish-brown sphalerite. One patch of pyrite pyrrhotite, galena and sphalerite from 164.4 to 165.0	5020	5.0			0.17	0.20						
165.0-170.0	5.0	Same as previous section. Heavy combined sulphides from 165.0 to 167.0.	5021	5.0			0.53	1.08						
170.0-175.0	5.0	Graphitic schist containing fine pyrite and pyrrhotite. A little reddish-brown sphalerite.	5022	5.0			TR	nil						
175.0-180.0	5.0	Graphitic schist with mineralization similar to previous section. Shearing contorted; some interbanded silica.	5023	5.0			0.03	nil						
180.0-185.0	5.0	Graphitic schist with about 10% pyrrhotite; sparse sphalerite. Shearing at 80° to core.	5024	5.0			TR	nil						

DIAMOND DRILL RECORD,

HOLE NO. 39

PROPERTY VIRGINIA CREEK, Y. T.

Alfred P.
April 15, 1977

SHEET NUMBER 1 SECTION FROM 0.0 TO 217.0
 LATITUDE 30,963.78N 9437.76N DATUM 4000
 DEPARTURE 30,522.41E 9303.23E BEARING _____
 ELEVATION 4,195.15 1357.01 W DIP Vertical hole

STARTED June 19, 1954

COMPLETED June 23, 1954

ULTIMATE DEPTH 334.0 (101.8) 1255.2

PROPOSED DEPTH _____
 To test lateral extent of mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-21.0	(6.4)	Casing (overburden)												
21.0-81.0	21.5	Graphitic schist. Shearing at 85° to core. A little interbanded silica and sericite schist, the latter becoming greater in amount from 76.0 to 81.0. Some white quartz patches.												
51.0-132.0	35.0	Dark grey sericite schist. Shearing at 85° to core. A little interbanded graphitic schist.												
132.0-142.1	4.0	Dominantly graphitic schist. A little interbanded dark grey sericite schist.												
142.1-161.0	15.0	Medium to dark grey sericite schist. Shearing at 90° to core. A little interbanded graphitic schist.												
161.0-191.5	9.6	Graphitic schist interbanded with dark grey sericite schist in about equal amounts. Shearing at 90° to core.												
191.5-196.0	1.4	Dark grey sericite schist. Mainly small fragments.												
	Note:	Change made from AX to XI core at 191.5 due to casing which necessitated use of X casing.												
196.0-201.0	1.9	Dark grey sericite schist as in previous section but with more white quartz present.								5026	200-205			
201.0-217.0	3.4	Fragments of graphitic and sericite schist containing some fine pyrite and a little galena and sphalerite. One sample over section due to low recovery.	5028	16.0			3.25	2.95		5027	205-210			
										5028	210-215			
										5029	215-220			

DIAMOND DRILL RECORD,

HOLE NO. 40

PROPERTY VANDORDA CREEK, I. T.

SHEET NUMBER 3 SECTION FROM 202.0 TO 295.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
202.0-266.0	59.2	Medium gray to dark gray sericite schist slightly talcose. Shearing at 70° to core. About 15% pyrrhotite in irregular patches. Sparse reddish-brown sphalerite.												
265.0-270.0	2.6	Graphitic schist. Shearing at 40° to core. Some interbanded silica and fine pyrite. A little reddish-brown sphalerite in irregular stringers.	5041	5.0			0.03	0.98						
270.0-275.0	5.0	Graphitic schist with an increase of pyrite relative to previous section. Shearing at 40° to core but contorted in part. Pyrrhotite and chalcoc present.	5042	5.0			0.02	nil						
275.0-281.0	3.0	Same as previous section. Sample taken over 6.0' due to low core recovery.	5043	6.0			0.94	0.88						
281.0-285.0	3.3	Massive fine pyrite containing vestiges of graphitic schist and sparse galena and sphalerite. Ballist bending at 70° to core.	5044	4.0			1.16	1.67						
285.0-290.0	3.7	Mineralization same as previous section changing to barren interbanded dark gray sericite schist and graphitic schist from 288.0 to 290.0.	5045	5.0			1.07	0.98						
290.0-295.0	4.0	Dark gray sericite schist changing to massive pyrite from 291.0 to 295.0. Some associated galena, sphalerite and gray silica. Ballist bending showing an anticlinal contortion in the plane of the core at 295.0. Sparse pyrrhotite and chalcocopyrite.	5046	5.0			1.43	1.58						

DIAMOND DRILL RECORD,

HOLE NO. 40

PROPERTY VANDORA CREEK, V. I.

SHEET NUMBER 4 SECTION FROM 295.0 TO 343.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
295.0- 300.0	4.3	Strong pyrite interbanded with grey silica. Relict banding at 50° to core. A little galena and sphalerite.	5047	5.0			4.62	4.82						
300.0- 305.0	2.8	Carbonatized dark grey schist. Some interbanded graphitic schist and a little fine pyrite.	5048	5.0			nil	nil						
305.0- 310.0	3.1	Same as previous section with a slight increase in pyrite content.	5049	5.0			TR	nil						
310.0- 315.0	5.0 M	Massive pyrite containing irregular sections of grey silica. Vestiges of graphitic schist. Relict banding at 50° to core.	5050	5.0			0.69	1.18						
315.0- 319.0	4.0	Interbanded pyrite, silica and graphitic schist. Bands are crenulated and contorted from 317.0- 319.0.	5051	4.0			0.02	0.39						
319.0- 326.0	5.8 5	Medium to dark grey sericite schist. Shearing at 80° to core, contorted in part. Several small patches of white.												
326.0- 331.0	5.0	Graphitic schist containing considerable interbanded grey silica. Scattered grains of pyrite and pyrrhotite. Shearing at 50° to core. Small crenulated fold at 328.5.	5052	5.0			TR	nil						
331.0- 338.0	6.8	Light grey sericite schist. Shearing at 80° to core. Several patches of white quartz containing threads of light brownish-white material. (Schistose)												
338.0- 343.0	3.6	Fine pyrite and pyrrhotite in a siliceous matrix. Sparse chalcopyrite.	5053	5.0			0.91	1.28						

DIAMOND DRILL RECORD,

HOLE NO. 40

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 5 SECTION FROM 343.0 TO 427.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
343.0-349.0	6.0 M	Massive fine pyrite associated with a little pyrrhotite. Relict banding at 60° to core. Vestiges of graphitic schist.	5054	6.0			1.49	2.75						
349.0-358.0	5.7 G	Graphitic schist. Highly carbonatized. Shearing irregular but averaging about 70° to core. Core broken in part.												
358.0-393.0	14.7 G	Graphitic schist. Shearing at 80° to core. A little interbanded silica and sparse pyrite.												
393.0-398.0	4.3 P	Interbanded graphitic schist and grey silica. About 30% fine pyrite. Sparse pyrrhotite and chalcopyrite.	5056	5.0			0.03	0.20						
398.0-403.0	4.7 P	Graphitic schist containing fine pyrite as above. Shearing at 70° to core.	5057	5.0			0.02	nil						
403.0-408.0	2.5 P	Graphitic schist interbanded with about 40% grey silica. Some fine pyrite throughout.	5058	5.0			0.55	0.29						
408.0-413.0	4.2	Graphitic schist with interbanded grey silica. Some disseminated pyrite as in previous sections.	5059	5.0			1.05	0.69						
413.0-418.0	3.9	Same as previous section changing to almost pure silica from 414.7 to 418.0 (quartzitic). About 30% pyrite.	5060	5.0			1.03	0.39						
418.0-423.0	3.8	Graphitic schist and silica in cemented bands. Considerable fine pyrite.	5061	5.0			1.02	0.98						
423.0-427.0	4.0	Same as previous section.	5062	4.0			TR	0.79						

DIAMOND DRILL RECORD,

HOLE NO. 41

April 15, 1977

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 287.0

STARTED June 24, 1954

LATITUDE 31,116.12 9484.19N

DATUM 4000

COMPLETED June 27, 1954

DEPARTURE 30,396.22 9264.76E

BEARING _____

ULTIMATE DEPTH 348.0

ELEVATION 4,212.11

DIP Vertical hole

PROPOSED DEPTH To test extension of mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-90		Casing (overburden)												
9.0-155.5	87.2	Dark gray sericite schist containing some interbanded graphitic schist. Shearing at 85° to core; quite fissile. Several patches of white quartz containing a little cubic pyrite. Core badly buttoned.												
155.5-163.0	2.4	Light greenish-gray, highly carbonatized sericite schist. Vestiges of interbanded graphitic schist.												
163.0-178.0	3.3	Fragments of graphitic schist.												
178.0-182.0	0	No core.												
182.0-233.0	16.0	Light greenish-gray carbonatized sericite schist. Soft, clay-like texture in part. Some intercalated fragments of graphitic schist, some of which is possibly cava. This section is suggestive of a healed fault zone.												
233.0-283.0	32.7	Light gray to medium gray sericite schist. About 10% pyrrhotite in irregular threads and small patches. Shearing contorted. Some patches of green-colored carbonate.												
283.0-287.0	4.0	Medium gray sericite schist containing about 10% pyrrhotite. Sparse reddish-brown sphalerite and chalcopyrite. Shearing at 40° to core.	5055	4.0			0.04	0.49						

DIAMOND DRILL RECORD,

HOLE NO. 42

PROPERTY VANDORDA CREEK, I. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 155.0

STARTED June 29, 1954

LATITUDE 28,812.30

DATUM 4000

COMPLETED July 2, 1954

DEPARTURE 30,765.25

BEARING _____

ULTIMATE DEPTH 326.0

ELEVATION 4,035.48

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-42.0		Casing (overburden)												
42.0-66.0	15.3 S	Light gray to medium gray sericite schist. Slightly talcose. Core broken into buttons in plane of shearing at 80° to core.												
66.0-71.0	1.6 S-T	Light gray sericite schist. Core badly broken. Several pieces of massive pyrite - pyrrhotite. No visible galena or sphalerite.	3065 A	5.0			0.38	0.98						
71.0-115.0	14.7 S	Light gray to medium gray sericite schist. Core badly broken. Sparse pyrrhotite and pyrite.												
115.0-135.0	4.6 G	Graphitic schist interbanded with gray silica. Shearing at 80° to core. A little fine pyrite.												
135.0-140.0	5.0 M	Massive pyrite and pyrrhotite associated with a gray silica matrix. A little reddish-brown sphalerite.	3066	5.0			1.51	1.97						
140.0-145.0	5.0	Same as previous section. Remnants of silicified sericite schist from 141.3 to 142.5.	3067	5.0			1.93	1.18						
145.0-150.0	5.0 P	Several patches of pyrrhotite-pyrite associated with light gray, talcose sericite schist.	3068	5.0			1.10	0.98						
150.0-155.0	1.6 P	Fragments containing pyrite and grayish-white sericite schist.	3069	5.0			0.72	0.59						

DIAMOND DRILL RECORD,

HOLE NO. 43

PROPERTY VANGUARD CREEK, I. T.

SHEET NUMBER 2 SECTION FROM 129.3 TO 255.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
129.3 132.0	2.7 N	Massive fine-grained pyrite. Some galena, sphalerite and silica in close association from 129.3 to 130.2. Sharp contact with the schist at 129.3.	5080	2.7			2.67	4.62						
132.0 - 137.0 137.0 142.0	5.0 M 5.0 M	Massive, fine grained pyrite. A few irregular threads of white quartz-carbonate. Massive fine grained pyrite changing to dominantly pyrrhotite. Some silica and closely associated galena and sphalerite. Mainly pyrrhotite from 140.0 to 142.0	5081 5082	5.0 5.0			0.94	1.18						
142.0 - 200.0	38.2 S	Light gray to medium gray sericite schist. Shearing at 50° to core but contorted in part. About 5% pyrrhotite in irregular patches. Several small patches of white quartz-carbonate. The schist is slightly talcose.												
200.0 205.0	5.0 S	Light gray sericite schist. Talcose in part. Shearing at 60° to core.												
205.0 210.0	5.0 P	Graphitic schist containing about 30% interbedded silica. Shearing contorted and the individual bands eroded. Considerable fine pyrite.	5083	5.0			0.44	0.69						
210.0 - 214.0 214.0 250.5	4.0 P 28.0 S	Graphitic schist containing mineralization similar to previous section. Light gray sericite schist. Strongly talcose and fissile. Shearing at 70° to core but showing local variations. A little interbedded chlorite.	5084	4.0			0.05	0.29						
250.5 - 255.0	4.0 P	Graphitic schist interbedded with gray silica. Some disseminated pyrite. Patches of pyrite pyrrhotite associated with gray silica from 254.0 to 255.0. Several threads of magnetite in the pyrrhotite. Sparse chalcopyrite.	5085	4.5			Tr	nil						

DIAMOND DRILL RECORD,

HOLE NO. 44

PROPERTY VANDORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 119.0

STARTED July 3, 1954

LATITUDE 28,920.46

DATUM 4,000

COMPLETED July 9, 1954

DEPARTURE 31,200.57

BEARING _____

ULTIMATE DEPTH 189.0

ELEVATION _____

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization.

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
1.0-59.0	(M)	Casing (overburden)												
59.0 - 64.0	5.0 M	Massive pyrite, galena and sphalerite associated with a fine silica matrix.	5095	5.0			2.58	7.57						
64.0 - 70.0	5.0 M	Mainly massive pyrite with some bands of pyrrhotite galena and reddish-brown sphalerite. A few small quartz patches.	5096	5.0			1.71	1.18						
70.0 - 77.0	5.0 M	Pyrite, pyrrhotite and silica streaked with minor galena and sphalerite. Some magnetite present.	5097	5.0			3.16	1.57						
77.0 - 89.0	5.0 M	Pyrrhotite and pyrite associated with grey silica and vestiges of sericite schist. A little galena and sphalerite.	5098	5.0			1.16	0.88						
89.0 - 96.0	5.0 M	Same as previous section with only sparse galena and sphalerite.	5099	5.0			0.04	0.29						
96.0 - 102.0	5.0 M	Heavily mineralized with pyrite pyrrhotite and magnetite bands. Some silica present as a matrix	0001	5.0			0.02	0.69						
102.0 - 107.0	5.0 M	estigial banding at 60° to core. Heavy pyrite mineralization in a grey silica matrix	0002	5.0			0.03	0.20						
107.0 - 112.0	5.0 M	Same as previous section. Helict banding at 50° to core.	0003	5.0			0.03	0.10						
112.0 - 107.0	5.0 M	Pyrite and silica similar to previous sections. Sparse Chalcopyrite.	0004	5.0			Tr	0.10						
107.0 - 112.0	5.0 M	Pyrite and pyrrhotite associated with grey silica and vestiges of sericite schist.	0005	5.0			Tr	0.10						
112.0 - 117.0	5.0 M	Fine grained pyrite admixed with silica. No other sulphide visible.	0006	5.0			Tr	Nil						
117.0 - 119.0	5.0 M	Fine pyrite, about 50%, in a silica matrix (silicified sericite schist) Pyrite massive in part.	0007	5.0			Tr	Nil						

DIAMOND DRILL RECORD,

HOLE NO. 44

PROPERTY YANGORDA CREEK, I. T.

SHEET NUMBER 2 SECTION FROM 139.0 TO 276.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
119.0 - 124.0	5.0	Same as previous section.	008	5.0			Nil	Tr						
124.0 - 129.0	5.0	Pyrite associated with grey silica. Vestiges of sericite schist and a little white quartz present.	009	5.0			0.02	0.02						
129.0 - 134.0	5.0	Same as previous section with some interbanded magnetite, sphalerite and galena from 131.5 to 132.0	010	5.0			0.91	2.16						
134.0 - 139.0	5.0	Heavily mineralized with pyrite in a silica matrix. Some interbanded pyrrhotite - magnetite. Sparse galena and sphalerite.	011	5.0			1.60	4.82						
139.0 - 142.0	3.0	Sericite schist containing interbanded pyrite, massive in part, with some associated galena and sphalerite, the latter reddish - brown.	012	3.0			2.72	7.78						
142.0 - 175.3	23.6	Light grey sericite schist containing a little talc and chlorite. Shearing at 80° to core. Sparse pyrite and pyrrhotite.												
175.3 - 180.0	4.7	Light grey sericite schist well mineralized with pyrite and small patches of magnetite.	013	4.7			Nil	0.20						
180.0 - 186.0	6.0	Same as previous section with the pyrite - magnetite massive in part. Some pyrrhotite present.	014	6.0			0.02	0.29						
186.0 - 209.5	20.0	Light grey sericite schist a little talc present. Shearing 70° to core. Sparse pyrite and pyrrhotite.												
209.5 - 273.0	54.0	Medium grey sericite schist. Slight greenish cast due to interbanded chlorite. Scattered pyrrhotite in small patches and threads roughly paralleling the shearing at 60° to core. Several patches of white quartz.												
273.0 - 278.0	5.0	Silicified sericite schist containing about 40% pyrite. Minor associated pyrrhotite, reddish-brown sphalerite and magnetite.	015	5.0			0.36	0.99						

DIAMOND DRILL RECORD,

HOLE NO. 44

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 3

SECTION FROM 278.0 TO 334.0

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
73.0 - 83.0	5.0	Heavy pyrite, pyrrhotite, magnetite and minor sphalerite replacing sericite schist. Relief banding at 80° to core.	016	5.0			0.39	0.49						
83.0 - 88.0	5.0	Interbanded graphitic schist and silica (Quartzite) containing a little disseminated pyrite.	017	5.0			0.11	0.10						
88.0 - 93.0	5.0	Graphitic schist interbanded with gray silica. Some fine pyrite and a little pyrrhotite present. A horse of talcy sericite schist from 228.0 to 239.0.	018	5.0			Tr	0.29						
93.0 - 98.0	5.0	Pyrite interbanded with grey silica. A little associated pyrrhotite and reddish-brown sphalerite. Banding at 70° to core.	019	5.0			0.11	0.29						
98.0 - 103.0	5.0	Same as previous section. Vestiges of graphitic schist present. Carbonated section from 300.0 to 301.0, associated with white quartz.	020	5.0			0.02	0.10						
103.0 - 119.0	11.8	Light gray sericite schist. Slightly talcose. Shearing 70° to core. Sparse sphalerite at 314.5 and 316.5, not worthy of assay.												
119.0 - 124.0	5.0	Graphitic schist containing some dark gray interbanded sericite schist from 319.0 to 322.0. Some disseminated pyrite.	021	5.0			Tr	0.10						
124.0 - 129.0	5.0	Graphitic schist interbanded with gray silica. Some disseminated pyrite and threads of reddish-brown sphalerite.	022	5.0			0.80	1.77						
129.0 - 134.0	5.0	Graphitic schist and interbanded silica. Considerable disseminated pyrite.	023	5.0			0.02	1.18						

DIAMOND DRILL RECORD,

HOLE NO. 45

PROPERTY VANDORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 160.0

STARTED July 5, 1954

LATITUDE 29,528.86N

DATUM 4,000

COMPLETED July 10, 1954

DEPARTURE 30,686.50E

BEARING _____

ULTIMATE DEPTH 367.0

ELEVATION 4,076.80

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization.

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-95.0	^	Casing (overburden)												
95.0 - 100.0	1.4	Fragments of massive sulphide mineralization consisting of pyrite and strong galena and sphalerite in a silica matrix. (Possibly a loose piece of rock at bedrock.)	025	5.0			6.63	7.64						
100.0 - 105.0	4.0	Massive sulphide same as previous section. Good galena - sphalerite content.	026	5.0			3.23	8.33						
105.0 - 110.0	5.0	Sharp increase in pyrite relative to the galena and sphalerite.	027	5.0			6.10	9.13						
110.0 - 115.0	5.0	Massive pyrite, galena and sphalerite in a silica matrix. Relief bending at 50° to core.	028	5.0			2.30	8.73						
115.0 - 120.0	5.0	Strong galena and sphalerite directly associated with pyrite and small patches of magnetite.	029	5.0			2.30	8.04						
120.0 - 125.0	2.0	Same mineralization as previous section, with a little less magnetite.	030	5.0			6.91	9.72						
125.0 - 130.0	2.6	Strong galena and sphalerite with pyrite. Relief bending at 40° to core.	031	5.0			1.38	8.93	✓					
130.0 - 135.0	3.8	Same as previous section to 132.5. Horse of chlorite schist, carbonated from 132.5 to 136.8.	032	5.0			0.02	4.87	✓					
135.0 - 140.0	5.0	Heavy galena and sphalerite associated with pyrite, silica and a little magnetite.	033	5.0			1.84	6.75						
140.0 - 145.0	5.0	Same as previous section with a little pyrrhotite present.	034	5.0			1.50	7.94						
145.0 - 150.0	4.0	Pyrite, silica, galena and sphalerite. One patch of magnetite at 145.2	035	5.0			7.83	7.25						
150.0 - 155.0	3.9	Same as previous section. Several small patches of magnetite at 154.5	036	5.0			4.03	6.16	✓					
155.0 - 160.0	5.0	Same as previous section to 157.0 changing to massive pyrite. A little magnetite and pyrrhotite.	037	5.0			0.92	2.28						

DIAMOND DRILL RECORD,

HOLE NO. 45

PROPERTY VANGUARD CREEK, Y. T.

SHEET NUMBER 2

SECTION FROM 160.0 TO 225.0

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
160.0 - 165.0	5.0	Pyrite, pyrrhotite, galena, sphalerite and magnetite contain vestigial fragments of vestigial sericitic schist. Sphalerite reddish-brown in colour.	038	5.0			2.42	3.08	160					
165.0 - 170.0	5.0	Massive pyrite containing irregular threads and small patches pyrrhotite-magnetite-gray quartz	039	5.0			0.02	0.30						
170.0 - 175.0	5.0	Same as previous section. Sparse galena and sphalerite.	040	5.0			0.58	0.70						
175.0 - 180.0	5.0	Massive fine grained pyrite. Several narrow bands of galena and sphalerite.	041	5.0			1.04	1.29						
180.0 - 185.0	5.0	Massive fine grained pyrite with some interstitial silica.	042	5.0			Tr	Nil						
185.0 - 190.0	5.0	Same as previous section.	043	5.0			Tr	0.30						
190.0 - 195.0	5.0	Same as previous section. A little pyrrhotite present.	044	5.0			Nil	0.10						
195.0 - 200.0	5.0	Several small patches of pyrrhotite in massive pyrite-silica matrix. Sparse chalcopyrite. Remnants of graphitic schist from 195.0 to 200.0	045	5.0			Nil	Nil						
200.0 - 205.0	5.0	Same as previous section. Remnants of graphitic schist from 202.0 to 203.0.	046	5.0			0.02	Nil						
205.0 - 210.0	5.0	Web-like patterns of pyrite and pyrrhotite in a silica matrix. Sparse chalcopyrite.	047	5.0			Tr	0.10						
210.0 - 215.0	5.0	Same as previous section. The pyrite is massive in part	048	5.0			Tr	0.20						
215.0 - 220.0	5.0	Pyrite and silica. Sparse galena and sphalerite.	049	5.0			0.27	0.10						
220.0 - 225.0	5.0	Pyrite and silica containing several small patches of pyrrhotite and magnetite. Sparse galena and sphalerite in direct association with the magnetite.	050	5.0			0.04	0.30						

DIAMOND DRILL RECORD,

HOLE NO. 45

PROPERTY VANDORDA CREEK, Y. T.

SHEET NUMBER 3 SECTION FROM 225.0 TO 280.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
225.0 - 230.0	5.0	Groundmass of pyrite and silica containing some magnetite stringers. A little galena and sphalerite associated with the latter.	051	5.0			0.04	0.29						
230.0 - 235.0	5.0	Web-like patterns of pyrite and silica containing small patches of white quartz, pyrrhotite, magnetite galena and sphalerite. Sparse chalcocopyrite.	052	5.0			0.44	1.38						
235.0 - 240.0	5.0	Same as previous section but showing an increase of magnetite and galena - sphalerite relative to the pyrite.	053	5.0			2.01	4.53						
240.0 - 245.0	5.0	Pyrite and silica heavily mineralized with sphalerite and galena. Magnetite directly associated with sphalerite.	054	5.0			3.14	8.06						
245.0 - 250.0	4.5	Same as previous section	055	5.0			1.90	3.94						
250.0 - 255.0	4.5	Pyrite and silica containing irregular sphalerite galena, magnetite & pyrrhotite.	056	5.0			0.99	2.46						
255.0 - 260.0	5.0	Fine grained closely-associated pyrite, silica, magnetite, pyrrhotite, galena and sphalerite.	057	5.0			2.46	3.25						
260.0 - 265.0	5.0	Same as previous section. Sphalerite has a reddish-brown colour.	058	5.0			1.76	3.64						
265.0 - 270.0	5.0	Assemblage of sulphides as above. Sphalerite prominent in association with magnetite.	059	5.0			3.66	6.69						
270.0 - 275.0	5.0	Same as previous section with an increased amount of silica. A few vestiges of silicified sericite visible.	060	5.0			4.52	4.43						
275.0 - 280.0	5.0	Pyrite and silica containing minor pyrrhotite, galena and sphalerite.	061	5.0			2.67	1.12						

DIAMOND DRILL RECORD,

HOLE NO. 46

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 2.0 TO 232.6

STARTED July 10, 1954

LATITUDE 29,249.10

DATUM L.M.

COMPLETED July 17, 1954

DEPARTURE 30,659.78

BEARING _____

ULTIMATE DEPTH 112.0

ELEVATION 4,092.40

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0 - 27.8	27.8	Casing (overburden) Light grey sericite schist. Shearing at 90° to core Talcose and containing a little interbanded pyrite and pyrrhotite. Core lost from 95.0 to 130.0.												
176.0 - 181.0	4.5	Dominantly pyrite with some associated quartz and a little reddish-brown sphalerite and galena.	065	5.0			2.81	3.32						
181.0 - 195.0	4.9	Light grey sericite schist with inter-banded medium gray sections. Talcose. Core bottomed at 80° to core.		14										
195.0 - 200.0	5.0	Massive pyrite containing some banded sections of magnetite at 50° to core. A little reddish-brown sphalerite.	066	5.0			2.36	2.42						
200.0 - 207.0	1.4	Massive pyrite from 200.0 to 200.8. The remaining 0.6 is fragments of quartz and sericite schist; sample taken over 7.0 rather than 5.0 due to low recovery.	067	7.0			.96	1.31						
207.0 - 212.0	5.0	Massive pyrite from 207.0 to 208.5. Balance of sec- tion is silicified and carbonatized sericite schist.	068	5.0			Tr	Tr						
212.0 - 217.9	5.0	Massive pyrite containing some narrow bands of magnetite. Sparse galena and sphalerite.	069	5.0			1.62	1.31						
217.9 - 222.0	5.0	Massive pyrite with less magnetite than previous section.	070	5.0			1.07	1.01						
222.0 - 227.0	4.5	Massive pyrite containing considerable reddish- brown sphalerite and vestiges of graphitic schist.	071	5.0			2.06	1.32						
227.0 - 232.0	4.5	Massive pyrite with vestiges of graphitic schist.	072	4.5			1.39	1.12						
232.0 - 233.6	1.6	Same as previous section with some reddish-brown galena and sphalerite present.	073	1.6			2.87	1.32						

DIAMOND DRILL RECORD,

HOLE NO. 47

PROPERTY VANCORDA CREEK, Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 195.0
 LATITUDE 29,681.23 DATUM 4,000
 DEPARTURE 30,557.25 BEARING _____
 ELEVATION 4,069.0 DIP Vertical Hole

STARTED July 12, 1954
 COMPLETED July 18, 1954
 ULTIMATE DEPTH 126.0
 PROPOSED DEPTH To cross-section mineralization.

2948

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-90.0		Casing (overburden)												
90.0-110.0	12.2	Graphitic schist interbanded with grey silica. Shearing at 10° to core. A little fine pyrite.												
110.0-115.0	4.0	Graphitic schist and quartzite interbanding as in previous section. Some pyrite and a little associated galena and sphalerite.	077	5.0				1.29	2.52					
115.0-120.0	5.0	Graphitic schist same as previous section with shearing circled in part.	078	5.0				3.71	3.92					
120.0-125.0	4.5	Graphitic schist as in previous section. Several small patches of white quartz.	079	5.0				2.97	2.92					
125.0-130.0	4.0	Graphitic schist as in previous section to 127.0. Massive pyrite from 127.0 to 130.0. Sparse sphalerite.	080	5.0				4.62	5.43					
130.0-135.0	5.0	Massive pyrite containing a little interbanded galena and sphalerite at 70° to core - controlled by original shearing.	081	5.0				4.16	5.06					
135.0-140.0	5.0	Massive pyrite containing intermittent sections high in reddish-brown sphalerite.	082	5.0				5.57	1.01					
140.0-145.0	5.0	Massive pyrite same as previous section. A little pyrrhotite, magnetite, galena and sphalerite. Siliceous breccia from 142.0 to 143.0.	083	5.0				2.28	2.52					
145.0-150.0	5.0	Same as previous section. Strong magnetite content from 147.0 to 148.5. Vestiges of sericite schist at 149.0.	084	5.0				3.71	1.41					
150.0-155.0	5.0	Massive, closely-associated pyrite, galena, sphalerite and silica containing remnants of silicified sericite schist.	085	5.0				2.06	3.23					

DIAMOND DRILL RECORD,

HOLE NO. 47

PROPERTY YANFORDA CREEK, T. T.

SHEET NUMBER 2 SECTION FROM 155.0 TO 230.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
155.0 -	5.0	Same as previous section with a section of white	086	5.0			2.67	6.65						
160.0 -	5.0	Quartz from 150.0 to 160.0 which continues to 160.0.	087	5.0			2.09	7.05						
165.0 -	5.0	Strong galena and sphalerite with pyrite in a silica matrix. Remnants of bleached pericite schist.	088	5.0			0.83	4.23						
170.0 -	5.0	Massive pyrite containing sections of finely associated pyrite, silica galena and sphalerite. A few small patches of white quartz.												
170.0 -	5.0	Fine pyrite, galena sphalerite and silica in close association.	089	5.0			3.60	10.26						
175.0 -	2.5	Mineralization same as previous section, porous and waxy in part.	090	5.0			1.71	5.66						
180.0 -	2.5	Irregular patches of reddish-brown sphalerite and gray silica in massive pyrite.	091	5.0			2.42	3.82						
185.0 -	5.0	Massive pyrite, close-textured containing a little sphalerite and galena	092	5.0			1.27	7.45						
190.0 -	5.0	Closely associated pyrite, silica, galena, sphalerite and magnetite.	093	5.0			2.72	7.25						
195.0 -	5.0	Same as previous section. Ballot banding at 80° to core.	094	5.0			2.92	6.65						
200.0 -	5.0	Sulphide aggregate as in previous two sections.	095	5.0			3.38	7.25						
205.0 -	5.0	Mainly massive pyrite, some galena pyrite and magnetite with silica from 205.0 to 206.0.	096	5.0			1.02	1.81						
210.0 -	5.0	Massive, closely-knit pyrite. A little associated grey quartz.	097	5.0			7.7	3.30						
215.0 -	5.0	Same mineralization as previous section.	098	5.0			8.11	3.10						
220.0 -	5.0	Massive pyrite containing small patches of grey silica.	099	5.0			0.49	7.7						
225.0 -	5.0	Pyrite and silica as in previous section	100	5.0			7.7	8.11						

DIAMOND DRILL RECORD,

HOLE NO. 47

PROPERTY VANGORDA CREEK, L. T.

SHEET NUMBER 3

SECTION FROM 230.0 TO 285.0

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
230.0 - 235.0	5.0	Massive, closely-knit pyrite with small patches of silica as in previous section.	101	5.0			.28	.40						
235.0 - 240.0	5.0	Massive pyrite and silica containing a little sphalerite, galena, white quartz. Sparse pyrrhotite and chalcopyrite.	102	5.0			2.09	.51						
240.0 - 245.0	5.0	Pyrite and silica containing a little relatively coarse galena at 241.0 to 241.3. Sparse pyrrhotite and chalcopyrite.	103	5.0			4.15	.60						
245.0 - 250.0	5.0	Pyrite and silica with heavy galena mineralization from 249.0 to 250.0. Sparse pyrrhotite and chalcopyrite. Sphalerite present with the galena.	104	5.0			11.60	.91						
250.0 - 255.0	5.0	Massive pyrite containing several small patches of magnetite and white quartz.	105	5.0			.11	.60						
255.0 - 260.0	5.0	Banded pyrite and silica at 80° to core. Sparse galena, sphalerite, pyrrhotite and chalcopyrite.	106	5.0			.68	1.11						
260.0 - 265.0	5.0	Strong magnetite patches in pyrite and silica. A little sphalerite with the magnetite.	107	5.0			.33	.50						
265.0 - 270.0	5.0	Heavy streaks of magnetite in pyrite and silica. A little pyrrhotite, sphalerite and chalcopyrite in association with the magnetite. Several patches of white quartz.	108	5.0			.33	1.01						
270.0 - 275.0	5.0	Pyrite and silica containing irregular bands and patches of magnetite, pyrrhotite and some dark brown sphalerite.	109	5.0			1.21	1.41						
275.0 - 280.0	5.0	Same as previous section with sparse greenish mineral suggestive of epidote.	110	5.0			1.50	3.62						
280.0 - 285.0	5.0	Closely associated pyrite, pyrrhotite, galena sphalerite, magnetite and silica.	111	5.0			1.84	8.06						

453.19

DIAMOND DRILL RECORD,

HOLE NO. 48

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 237.0

STARTED July 17, 1954

LATITUDE 29,546.81 9005.87

DATUM L.O.D.

COMPLETED July 25, 1954

DEPARTURE 30,131.22 9183.99

BEARING _____

ULTIMATE DEPTH 381.0

ELEVATION 4,051.04 +257 1313.71

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization.

DEPTH FEET	CORE RECOVER	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-120.0		Casing (overburden)												
120.0 - 169.0	8.0	Medium grey sericite schist. Mainly fragments, but short sections show shearing at 80° to core. Recovery low due to fissile, flakey nature of the schist. White quartz from 168.0 to 169.0. No evidence of sulphide.												
169.0 - 170.5	1.5	Massive pyrrhotite containing a little pyrite, galena and sphalerite in association.	171	1.5			2.92	2.11						
170.5 - 189.3	4.8	Dark grey talc-sericite schist. A few small patches of white quartz-carbonate shearing at 50° to core.												
189.3 - 194.0	4.4	Several patches of magnetite in pyrite. A little associated galena and sphalerite. Vestiges of schist present.	172	4.7			3.66	3.72						
194.0 - 199.0	3.1	Fine pyrite containing small patches and stringers of magnetite. Relict banding at 50° to core. Sparse galena and sphalerite.	173	5.0			3.66	3.62						
199.0 - 204.0	5.0	Same mineralization as previous section. Slight increase in galena-sphalerite content.	174	5.0			3.05	2.72						
204.0 - 212.0	1.7	Banded pyrite, magnetite and sparse sphalerite for 1.2 feet from 204.0. Balance of the run has high core loss in sericite schist. Sample taken over 8.0 feet due to high core loss.	175	8.0			2.72	2.52						
212.0 - 231.4	4.9	Light gray sericite schist. Fissile shearing at 70° to core. Core buttoned.												
231.4 - 237.0	1.9	Light gray kaolinized sericite schist containing pyrite and magnetite bands paralleling the shearing at 80° to core. Sparse galena and sphalerite. Sample taken over 5.6 to block at 237.0 due to low recovery.	176	5.6			1.15	2.32						

DIAMOND DRILL RECORD,

HOLE NO. 49

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 133.0
 LATITUDE 29,226.05 DATUM 4000
 DEPARTURE 30,945.41 BEARING _____
 ELEVATION 4,110.12 DIP Vertical hole

STARTED July 18, 1954

COMPLETED July 23, 1954

ULTIMATE DEPTH 361.0

PROPOSED DEPTH To cross-section depth

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0- 18.0- 18.0- 21.0		Casing (overburden)												
21.0- 121.0	3.0	Massive closely-knit sulphides consisting of pyrite galena and sphalerite in a silica matrix.	125	3.0			3.81	9.06	132	120-125	4'		1.46	3.52
									133	125-130			1.35	1.62
121.0- 125.0	0.4	Three fragments of light gray sericite schist with one button of sulphides similar to previous section. See sludge sample for this section.							134	130-135	2'		1.38	2.52
									135	135-140	5'		.63	1.01
125.0- 130.0	4.5	light gray schist from 125.0 to 125.5. Balance of section is massive pyrite containing several streaks of galena and sphalerite. Some vestiges of sericite schist.	126	5.0			1.65	3.22	136	140-145	5'		.52	.60
130.0- 133.0	2.1	Massive, closely-knit pyrite with a little associated pyrrhotite and silica.	127	3.0					137	145-150	1'		.96	.71
133.0- 141.0	2.1	Fragments of dark gray sericite schist. One button containing streaks of sphalerite. See sludge sample.												
141.0- 146.0	1.2	From 141.0 for 1.2 of the short recovery, dark gray sericite schist with streaks of sphalerite. Balance of sample is massive pyrite with fragments of graphitic schist.	128	3.0			1.29	1.41						
146.0- 151.0	5.0	Massive pyrite containing streaks of galena and sphalerite roughly paralleling vertical bedding at 70° to core. Several patches of gray carbonate etc.	129	5.0			.72	.60						
151.0- 156.0	5.0	As in previous section with increased galena and sphalerite content. A little magnetite.	130	5.0			3.34	5.64						

DIAMOND DRILL RECORD,

HOLE NO. 9

PROPERTY VANDORA CREEK, T. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 139.0

STARTED July 19, 1954

LATITUDE 29,279.06

DATUM 4000

COMPLETED July 24, 1954

DEPARTURE 30,813.34

BEARING _____

ULTIMATE DEPTH 325.0

ELEVATION 4,092.02

DIP Vertical

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-107.0		Casing (overburden)												
107.0-112.0	3.2	Massive pyrite containing some irregular sections of galena, sphalerite and siliceous. Ballist banding at 45° to core.	138	5.0			2.25	4.63						
112.0-117.0	2.5	Increase in galena, sphalerite and silica relative to the pyrite.	139	5.0			11.28	10.86						
117.0-119.0	1.5	Same as previous section. Some remnants of siliceous cream-coloured breccia.	140	2.0			4.26	4.63						
119.0-124.0	3.4	Massive pyrite with irregular threads of magnetite.	141	5.0			7.2	9.1						
124.0-129.0	5.0 15.6	Massive pyrite containing irregular seams of magnetite. Sparse galena and sphalerite. Vestiges of inclined sericite schist.	142	5.0			2.72	4.02						
129.0-133.0	2.8	Massive pyrite containing some porous fractures. Some associated magnetite. Strong siliceous-sphalerite content from 132.0 to 133.0	143	4.0			2.01	4.93						
133.0-136.0	1.5	Massive pyrite. Porous, brecciated appearance due to probable crushing and healing. Low recovery likely due to porosity.	144	3.0			1.46	4.22						
136.0-139.0	0.9	Same type of pyrite as in previous section - low recovery.	145	3.0			1.32	2.11						

DIAMOND DRILL RECORD,

HOLE NO. 3

PROPERTY VANGORDA CREEK, T. T.

SHEET NUMBER 3 SECTION FROM 220.0 TO 279.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
220.0-225.0	2.9	Pyrite, silica, galena and sphalerite in rilliet banding at 60° to core.	155	5.0			3.69	5.23						
225.0-230.0	2.7	Graphitic schist interbanded with gray silica. Pyrite present with sparse sphalerite.	156	5.0			2.08	4.03						
230.0-235.0	1.8	Graphitic schist. Some disseminated pyrite. Shearing contorted.	157	5.0			1.49	2.52	✓					
235.0-246.0	2.7	Medium gray sericite schist fragments. A little white quartz. Sludges sampled over this section.							167 235-240	51C		.80	1.91	
									168 240-245	51C		.34	.80	
246.0-251.0	3.5	Massive pyrite containing bands of magnetite. Some interbanded galena, sphalerite and silica.	158	5.0			5.53	6.53	169 245-250			2.74	4.72	
251.0-255.0	4.0	Massive pyrite with irregular patches of galena, sphalerite and silica.	159	4.0			3.69	4.22	Sludges small in above samples due to weak water return.					
255.0-260.0	5.0	Massive pyrite. Two small patches of combined galena, sphalerite, silica and magnetite.	160	5.0			1.85	3.32						
260.0-265.0	5.0	Massive pyrite containing several bands of galena and sphalerite in rilliet shearing at 50° to core.	161	5.0			1.85	2.82						
265.0-270.0	1.6	Massive pyrite with minor pyrrhotite and magnetite.	162	5.0			2.54	4.12	170 265-270			1.54	2.42	
270.0-275.0	5.0	Massive pyrite with several small patches of galena-sphalerite. A few magnetite threads.	163	5.0			1.62	2.52	Sludge (above) small due to weak water return.					
275.0-279.0	4.0	Massive pyrite combined with pyrrhotite. Dominantly pyrrhotite toward end of sample.	164	4.0			.11	.20						

DIAMOND DRILL RECORD,

HOLE NO. 52

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 140.0

STARTED July 25, 1954

LATITUDE 29,073.78

DATUM 1000

COMPLETED July 31, 1954

DEPARTURE 31,071.61

BEARING _____

ULTIMATE DEPTH 590.0

ELEVATION 4,107.42

DIP Vertical hole

PROPOSED DEPTH To cross section mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
0.0-105.0		Casing (overburden)												
105.0-110.0	4.0 M	Massive pyrite with a few streaks of reddish-brown sphalerite and galena, following relief banding at 70° to core.	194	5.0			6.42	6.53						
110.0-115.0	5.0 M	Same as previous section with increased galena and sphalerite. Fine silica matrix with the sulphides. Quartz-carbonate and vestiges of sericite schist from 118.0 to 119.2.	195	5.0			4.83	8.25						
115.0-120.0	5.0 M	Prominent galena - sphalerite content in association with pyrite and silica. Reddish-brown cast to sphalerite. Relief banding at 40° to core.	196	5.0			4.04	10.15						
120.0-125.0	4.6 M	well mineralised pyrite, galena, sphalerite and silica from 120.0 to 123.5. Semi-porous, granular pyrite from 123.5 to 125.0.	197	5.0			2.71	6.74						
125.0-130.0	MV 2.6	Porous granular pyrite. Recovery low due to this characteristic. Several small patches of white quartz.	198	5.0			1.31	1.01						
130.0-135.0	5.0 ^{MV}	Loosely-knit fine pyrite. One section from 130.5 to 131.5 showing good reddish-brown sphalerite and galena mineralization; silica in the interstices.	199	5.0			1.20	3.02						
135.0-140.0	5.0 M	Massive closely-knit pyrite with a silica matrix. A few rice-sized patches of magnetite. Galena or sphalerite not seen.	200	5.0			1.29	1.30						

DIAMOND DRILL RECORD,

HOLE NO. 2

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 2 SECTION FROM 140.0 TO 187.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
140.0-145.0	5.0	Massive fine pyrite, porous from 142.0 to 143.0. Silica matrix with the pyrite. A little galena and magnetite.	201	5.0			.149	.30						
145.0-150.0	4.2	Considerable magnetite in pyrite from 145.0 to 147.0 Pyrite with galena and sphalerite in silica matrix in remainder of sample. Helict banding at 85° to core	202	5.0			3.02	4.22						
150.0-155.0	M 4.3	Pyrite, galena and sphalerite, the latter two decreasing toward end of sample.	203	5.0			2.45	4.83						
155.0-160.0	M 4.3	Pyrite and silica in a hard, brittle texture.	204	5.0			.54	.20						
160.0-165.0	M 4.8	Same as the previous section with vestiges of graphite schist present.	205	5.0			.14	.40						
165.0-170.0	M-P 5.0	Pyrite, silica and vestiges of graphitic schist in about equal amounts.	206	5.0			.04	nil						
170.0-175.0	5.0	Irregular patches of siliceous matrix in association with fine pyrite. Magnetite or pyrrhotite not visible	207	5.0			.26	.30						
175.0-180.0	T 5.0	Pyrite and silica in closely-associated equigranular patterns. Fracturing at 80° to core.	208	5.0			1.05	1.61						
180.0-185.0	M 5.0	Massive closely-knit pyrite with several small patches of grey silica.	209	5.0			.49	.10						
185.0-187.0	M 2.4	Massive pyrite as in previous section.	210	2.4			.28	nil						

DIAMOND DRILL RECORD,

HOLE NO. 3

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 3 SECTION FROM 187.0 TO 375.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
187.0-253.0	48.2	Light gray to medium gray sericite schist. Talcosse in part. Pearly lustre on shear planes at 70° to core. Some fine interbanded pyrrhotite, and sparse chalcoprite. From 219.0 to 220.5 there is about 40% pyrrhotite with some poikilitic pyrite - not worthy of assay. Several small patches of white quartz.												
253.0-347.0	77.2	Light gray talc-sericite schist. A little interbanded chlorite. Sparse pyrrhotite. One thread of sphalerite at 251.8. Shearing at 80° to core. About 20% pyrrhotite between 298.0 and 306.0.												
347.0-352.0	5.0	Graphitic schist with interbanded silica. Shearing crenulated and replaced by considerable fine pyrite, which is massive in part. Several threads of reddish-brown sphalerite.	211	5.0			.57	.61						
352.0-357.0	2.3	Graphitic schist containing some interbanded silica and sericite schist. Shearing at 70° to core, but contorted in part. Some pyrite and reddish-brown sphalerite present.	212	5.0			.02	nil						
357.0-360.0	2.6	Graphitic schist with mineralization similar to previous section.	213	3.0			.31	1.11						
360.0-370.0	10.0	Light gray sericite schist. Shearing at 80° to core. Some interbanded pyrrhotite in the shear planes.												
370.0-375.0	3.8	Graphitic schist with interbanded silica and pyrite. Shearing at 50° to core.	214	5.0			.40	.57						

DIAMOND DRILL RECORD,

HOLE NO. 5

PROPERTY VANCORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 316.0

STARTED July 27, 1954

LATITUDE 29,400.18

DATUM 4000

COMPLETED August 3, 1954

DEPARTURE 30,529.37

BEARING _____

ULTIMATE DEPTH 512.0

ELEVATION 4,074.77

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-130.0	(39.6)	Casing (overburden)												
130.0-182.0	21.9	Dark grey sericite schist. Pearly lustre on shear planes at 80° to core. A little interbanded granitic schist.												
182.0-204.0	12.6	Light grey sericite replaced by considerable white quartz-carbonate. One 2" section of pyrite, galena and sphalerite at 190.3.												
204.0-209.0	3.9	Well mineralized with fine pyrite, silica and considerable galena and sphalerite in close association.	253	5.0			18.01	12.68						
209.0-253.0	12.0	Dark grey sericite schist containing a little interbanded graphitic schist. Shearing at 80° to core.												
253.0-256.3	3.3	Massive fine pyrite with streaks of reddish-brown sphalerite, galena and vestiges of graphitic and sericite schist.	254	3.3			2.79	4.13						
256.3-296.3	9.2	Light gray to medium grey sericite schist, talcose in part. Shearing at 80° to core, buttoned in part.												
296.3-301.0	4.5	Fine pyrite and silica with vestigial banding at 30° to core. Some galena and sphalerite in association. Remnant of bleached sericite schist from 297.5 to 298.9.	255	4.7			1.51	6.14						
301.0-306.0	5.0	Massive pyrite and silica containing vestiges of sericite schist. A little galena and sphalerite.	256	5.0			1.23	6.64						
306.0-311.0	5.0	Pyrite, silica, galena and sphalerite in close association. Vestiges of bleached sericite schist.	257	5.0			1.68	6.44						
311.0-316.0	5.0	Same as previous section. Fine sugary texture.	258	5.0			4.93	8.05						

DIAMOND DRILL RECORD,

HOLE NO. 53

PROPERTY VANGORDA CREEK, I.T.

SHEET NUMBER 2 SECTION FROM 316.0 TO 462.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
316.0	5.0	Same as previous sections. Sphalerite grains reddish-brown in colour.	259	5.0			3.82	7.45						
321.0	3.3	Fragments of dark grey sericite schist and quartz.												
336.3		No evidence of sulphide-possible fault zone, of low recovery due, to gouge.		1.0										
336.3 - 340.1	3.8	Pyrite and silica well mineralized with galena and sphalerite, the latter reddish-brown in colour.	260	3.8			3.02	9.36						
340.1 - 427.0	72.4	Light grey to medium grey sericite schist with a little interbanded chlorite. Scattered small patches of white quartz. A little pyrrhotite from 381.0 to 387. with sparse associated chalcovrite. Shearing at 65° to core. Sparse pyrrhotite from 390.0 to 417.0, in irregular small threads.												
427.0 - 437.0	9.8	Light grey sericite schist containing sparse pyrite. Shearing at 60° to core.												
437.0 - 442.0	2.0	Graphitic schist and interbanded silica replaced by about 50% fine pyrite.	261	5.0			0.03	ml						
442.0 - 447.0	5.0	Massive fine-grained pyrite containing remnants of graphitic schist.	262	5.0			0.02	ml						
447.0 - 452.0	5.0	Patches of fine pyrite and pyrrhotite in brecciated carbonate matrix. Graphitic schist with interbanded pyrite and silica from 450.0 to 452.0.	263	5.0			0.02	ml						
452.0 - 459.3	2.5	Graphitic schist, broken in buttons on the shear planes at 60° to core. Sparse fine pyrite. Not worthy of assay.												
459.3 - 462.0	2.7	Patches of fine pyrite in partially replaced quartz carbonate. Several small patches of magnetite.	264	2.7			0.11	ml						

DIAMOND DRILL RECORD,

HOLE NO. 34

PROPERTY VANGORDA CREEK Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 101.0 STARTED July 28, 1954
 LATITUDE 27,924.52 DATUM 1000 COMPLETED July 31, 1954
 DEPARTURE 31,296.99 BEARING _____ ULTIMATE DEPTH 360.0
 ELEVATION 4,077.10 DIP Vertical hole PROPOSED DEPTH To test magnetic anomaly

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-26.8		Casing (overburden)												
26.8-45.0	1.7	Fragments of pyrite containing irregular small patches of magnetite.							234	35-40			1.28	ml
45.0-50.0	4.1	Massive pyrite containing small patches of magnetite elongated in the banding at 55° to core. Sparse galena and sphalerite.	220	5.0			3.99	3.82	235	40-45			1.84	1.90
50.0-54.0	3.8	Massive, closely-knit pyrite with magnetite as in previous section.	221	4.0			1.66	1.61	236	45-50	n.a.		ml	0.80
54.0-59.0	5.0	Pyrite, silica, galena and sphalerite showing relic banding from 57.0 to 59.0 at 50° to core. About 20% magnetite from 54.0 to 55.5.	222	5.0			2.28	2.52	237	50-55	n.a.		ml	0.44
59.0-64.0	4.1	Massive pyrite with small patches of magnetite. Remnants of graphitic schist from 63.0 to 64.0.	223	5.0			1.45	1.31	238	55-60	n.a.			
64.0-69.0	2.2	Fragments of silicified graphitic schist containing a little pyrite.	224	5.0			.03	.31	239	60-65	n.a.			
69.0-74.0	2.9	Scattered pyrite grains in a siliceous matrix; massive in part.	225	5.0			Tr	ml	240	65-70			ml	0.30
74.0-79.0	2.6	Same as previous section but mostly in fragments.	226	5.0			Tr	.26	241	70-75			ml	0.41
79.0-101.0	6.0	Fragments of dark grey sericite schist. Pearly lustre on shear planes.							242	75-80			ml	0.30

DIAMOND DRILL RECORD,

HOLE NO. 35

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 63.0 STARTED August 1, 1954
 LATITUDE 28,307.08 DATUM MTD COMPLETED August 4, 1954
 DEPARTURE 31,993.43 BEARING _____ ULTIMATE DEPTH 321.0
 ELEVATION 4,103.11 DIP Vertical hole PROPOSED DEPTH To cross section mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0- 27.0		Casing (overburden)												
27.0- 32.5	1.4	Massive, closely-knit pyrite. Yuggy in part. A little silica present. (From 20-27 granular pyrite was drilled, which did not core so that bedrock is likely at 20.0).	267	5.5			0.37	0.60	298	20-25		1.65		1.10
32.5- 38.0	1.5	Same as previous section, with a few magnetite stringers present.	268	5.5			0.31	ml	299	25-30		2.88		2.11
38.0- 43.0	5.0	Massive pyrite, pyrrhotite, magnetite with a little galena and sphalerite, containing angular fragments of quartz and carbonate.	269	5.0			2.05	3.70	300	30-35		4.27		3.70
43.0- 48.0	5.0	Same as previous section with an increased amount of galena and sphalerite.	270	5.0			3.11	4.70	301	35-40		1.63		8.40
48.0- 53.0	4.1	Pyrite, pyrrhotite, galena and sphalerite (reddish-brown) in a silica matrix. Relict banding at 30° to core.	271	5.0			0.71	3.30						
53.0- 58.0	4.0	Pyrite, pyrrhotite and magnetite in silica, well mineralized with streaks of galena and sphalerite. From 56.6 to 57.0 massive galena with some sphalerite, peppered with fine pyrite and magnetite.	272	5.0			12.33	7.90		8.72				
58.0- 63.0	5.0	Patches of pyrite, pyrrhotite, sphalerite and galena with some magnetite in a silica matrix. Relict banding at 45° to core. Vestiges of silicified schist present.	273	5.0			2.57	3.30						

DIAMOND DRILL RECORD,

HOLE NO. 55

PROPERTY YANGORDA CREEK, Y. T.

SHEET NUMBER 2 SECTION FROM 63.0 TO 203.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
63.0-68.0	5.0	Banded pyrite with a little galena and sphalerite in a siliceous groundmass. Banding at 70° to core.	274	5.0			0.23	0.50						
68.0-73.0	4.4	Magnetite and pyrite in a siliceous matrix.	275	5.0			nil	0.50						
73.0-78.0	5.0	Pyrite in a siliceous matrix with a little associated magnetite and pyrrhotite. Sparse sphalerite.	276	5.0			0.09	0.90						
78.0-83.0	5.0	Same mineralisation as previous section.	277	5.0			0.31	1.40						
83.0-88.0	5.0	Banded pyrite and silica and a little magnetite and reddish-brown sphalerite. Banding obscured in part, by sulphide.	278	5.0			0.20	0.70						
88.0-93.0	5.0	Same mineralisation as previous section.	279	5.0			0.02	0.10						
93.0-98.0	5.0	Pyrite, pyrrhotite with sparse sphalerite in banded vestiges of silicified schist, 50° to core.	280	5.0			0.02	0.20						
98.0-102.0	3.5	Scattered pyrite, pyrrhotite with sparse chalcopyrite in silicified sericite schist.	281	4.0			0.17	0.10						
102.0-193.2	51.5	Light grey to medium grey sericite schist. Talcoses in part. A little interbanded pyrite and pyrrhotite accompanied by silicification. Shearing at 80° to core.												
193.2-198.0	2.5	Graphitic schist with interbanded grey silica and pyrite. Sparse reddish-brown sphalerite.	282	4.8			0.09	nil						
198.0-203.0	2.9	Same mineralisation as previous section. Shearing at 60° to core.	283	5.0			.33	.60						

DIAMOND DRILL RECORD,

HOLE NO. 55

PROPERTY VANGORDA CREEK, I. T.

SHEET NUMBER 3 SECTION FROM 203.0 TO 248.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
203.0-208.0	3.6	Graphitic schist, finely interbanded on the shear planes with silica and pyrite at 45° to core.	284	5.0			1.00	1.39						
208.0-213.0	2.0	Same mineralisation as previous section.	285	5.0			0.29	0.30						
213.0-218.0	3.5	Graphitic schist with pyrite and silica as in previous sections. Shearing at 70° to core.	286	5.0			0.43	0.10						
218.0-223.0	3.6	Graphitic schist as in previous section.	287	5.0			0.11	nil						
223.0-228.0	5.0	Graphitic schist with pyrite banding to 224.5, changing sharply to a grey chert-like matrix, containing fine pyrite and well-defined threads of galena and sphalerite.	288	5.0			0.89	0.60						
228.0-233.0	3.7	Chert-like, siliceous groundmass containing sharply defined threads and small patches of pyrite, galena and reddish-brown sphalerite. Graphitic schist banded with pyrite from 232.0 to 233.0.	289	5.0			3.74	6.42						
233.0-238.0	5.0	Graphitic schist banded with pyrite and a little silica. Sparse galena and sphalerite.	290	5.0			2.78	3.01						
238.0-243.0	2.9	Graphitic schist with irregular patches of pyrite, galena and sphalerite.	291	5.0			3.34	3.48						
243.0-248.0	5.0	Graphitic schist banded with pyrite, silica and sparse galena and sphalerite, at 80° to core.	292	5.0			2.32	2.10						

DIAMOND DRILL RECORD,

HOLE NO. 56

PROPERTY VANGORDA CREEK, I. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 127.0

STARTED August 2, 1954

LATITUDE 29,203.09

DATUM 4000

COMPLETED August 8, 1954

DEPARTURE 31,226.13

BEARING _____

ULTIMATE DEPTH 488.0

ELEVATION 4,127.33

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
0.0-38.0		Casing (overburden). Soft, granular mineralization starting at 80.0. No core.												
38.0-92.0	1.2 M	Massive fine-grained pyrite with several small patches of magnetite. A little galena, sphalerite and silica in association, at the end of the section.	302	4.0		9.45	8.22	335	75-80			1.06	1.60	
92.0-97.0	4.6 M	Uniform, fine-grained fabric of pyrite, silica, galena and sphalerite.	303	5.0		2.48	7.11	336	80-85			Tr	0.80	
97.0-102.0	4.7 M	Same as previous section to 99.0. Balance of sample is magnetite, grey quartz, sparse galena and sphalerite in massive pyrite.	304	5.0		1.66	4.58	337	85-90			Tr	1.22	
102.0-107.0	5.0 M	Fine-grained pyrite with considerable magnetite in irregular patches. A little pyrrhotite, galena and sphalerite. Sparse chalcopyrite.	305	5.0		.86	1.10	338	90-95					
107.0-112.0	5.0	Pyrite and pyrrhotite in close association. Some interstitial quartz from 108.5 to 110.0. Sparse chalcopyrite.	306	5.0		.20	.20							
112.0-117.0	4.9	Mineralization as in previous section with some irregular patches of grey silica and white quartz throughout.	307	5.0		0.86	nil							
117.0-122.0	5.0 M	Pyrite, pyrrhotite, silica and a few threads of reddish-brown sphalerite and galena.	308	5.0		2.03	2.50							
122.0-127.0	5.0 P	Silicified sericite schist containing about 40% pyrite and pyrrhotite with sparse chalcopyrite, galena and sphalerite.	309	5.0		1.51	2.21							

DIAMOND DRILL RECORD,

HOLE NO. 36

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 2 SECTION FROM 127.0 TO 170.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE NO.	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.			AG.	CU.	PB.	ZN.
127.0-132.0	5.0	Same as previous section with some small patches of magnetite and slightly more reddish-brown sphalerite.	310	5.0			0.31	0.50						
132.0-137.0	5.0	Light gray sericite schist, silicified and containing irregular patches of pyrite, pyrrhotite and magnetite with a little chalcocopyrite.	311	5.0			0.26	0.30						
137.0-142.0	5.0	Same mineralization as previous section. Chalcocopyrite present in minute fractures paralleling the core.	312	5.0			0.34	0.89						
142.0-147.0	5.0	Pyrrhotite, pyrite, magnetite and sparse chalcocopyrite and sphalerite in silicified sericite schist.	313	5.0			0.37	1.58						
147.0-152.0	5.0	Light gray sericite schist about 30% replaced by mineralization as in previous section.	314	5.0			0.29	0.70						
152.0-157.0	3.0	Light gray sericite schist with pyrite, pyrrhotite and sparse reddish-brown sphalerite. Shearing at 50° to core.	315	5.0			0.06	0.21						
157.0-162.0	4.5	Light gray, chalky sericite schist, silicified in part. Some fine pyrite and pyrrhotite interbanded in the shear planes at 70° to core.	316	5.0			0.26	0.20						
162.0-166.0	3.0	Light gray sericite schist, silicified in part, with mineralization same as previous section.	317	4.0			nil	nil						
166.0-170.0	3.0	Light gray sericite schist as in previous section. Some web-like threads of reddish-brown sphalerite and galena with white quartz between 169.0 and 169.4.	318	4.0			0.49	1.61						

DIAMOND DRILL RECORD,

HOLE NO. 57

PROPERTY YANGORDA CREEK, Y. T.

SHEET NUMBER 2 SECTION FROM 295.0 TO 488.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
295.0- 297.0	2.0	Graphitic schist with silica and pyrite as in previous section. Pyrite massive in part.	351	2.0			2.19	1.68						
297.0- 374.0	57.0	Graphitic schist containing a little finely-inter-banded silica. Shearing at 80° to core. Some inter-banded talc-sericite schist from 342.0 to 359.0. The graphitic schist is brittle and carbonaceous.												
374.0- 403.0	22.5	Graphitic schist with considerable white quartz in irregular patches and threads. Shearing at 50° to core.												
403.0- 406.5	3.5	Massive pyrite, galena, sphalerite and silica, in close association, replacing graphitic schist.	352	3.5			3.81	5.20						
406.5- 410.0	3.5	Massive fine pyrite and silica replacing graphitic schist. Shearing (vestigial) at 45° to core.	353	3.5			0.20	0.10						
410.0- 473.0	35.1	Graphitic schist with some interbanded grayish-white silica. Several patches of white quartz-carbonate. Shearing between 40° and 70° to core, the latter from 420.0 to 457.0.												
473.0- 478.0	3.9	Graphitic schist with some interbanded silica and some pyrite. Shearing at 60° to core.	354	5.0			1.28	1.71						
478.0- 483.0	5.0	Silicified graphitic schist well mineralized with fine pyrite. No visible galena or sphalerite.	355	5.0			0.69	0.79						
483.0- 488.0	5.0	Graphitic schist, silicified and mineralized as in previous section. A little chalcopyrite and pyrrhotite present.	356	5.0			0.31	0.60						

DIAMOND DRILL RECORD,

HOLE NO. 58

PROPERTY VANGORDA CREEK, I. T.

SHEET NUMBER 1 SECTION FROM _____ TO _____ STARTED August 5, 1954
 LATITUDE 28,265.06 DATUM 4000 COMPLETED August 7, 1954
 DEPARTURE 31,373.08 BEARING _____ ULTIMATE DEPTH 317.0
 ELEVATION 4,064.87 DIP Vertical hole PROPOSED DEPTH To test anomaly in relation to mineraliz

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-15.0	6.5	Casing (over burden)												
15.0-20.0	1.3	Fine grained pyrite, silica, magnetite, sphalerite and galena in close association. Porous in part.	323	5.0			3.34	6.84	339	15-20			1.77	6.25
20.0-25.0	4.4	Massive sulphides as above. Relict banding at 70° to core. Porous from 20.0 to 21.5	324	5.0			3.92	6.71						
25.0-30.0	5.0	Same as previous sections. Magnetite prominent in the banding. Good galena-sphalerite content.	325	5.0			3.46	7.50						
30.0-35.0	3.5	Increased pyrite relative to the magnetite, galena and sphalerite. (reddish-brown).	326	5.0			5.65	7.72						
35.0-40.0	2.7	Pyrite, magnetite, sphalerite and galena in silica matrix. Relict banding at 80° to core.	327	5.0			3.64	8.03	340	35-40			4.62	6.83
40.0-45.0	5.0	Same as previous section - well mineralized with galena and sphalerite. Prominent magnetite content.	328	5.0			4.15	6.18						
45.0-50.0	3.3	Massive fine-grained pyrite. Sparse magnetite.	329	5.0			1.73	2.11	341	45-50			1.14	1.10
50.0-55.0	1.8	Fragments and buttons of massive pyrite. Vestiges of kaolinitized sericite schist.	330	5.0			4.15	5.52	342	50-55			1.06	1.32
55.0-60.0	2.4	Massive pyrite. Relict banding at 50° to core containing rice-size patches of magnetite.	331	5.0			1.43	1.38	343	55-60			0.60	0.88
60.0-65.0	3.7	Massive pyrite to 61.0. Balance of section is graphitic schist, crumpled in part and containing pyrite, pyrrhotite and silica interbanding.	332	5.0			1.00	1.11	344	60-65			2.03	1.37

DIAMOND DRILL RECORD,

HOLE NO. 60

PROPERTY VANGORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 184.0

STARTED August 9, 1954

LATITUDE 29,508.55

DATUM 4000

COMPLETED August 15, 1954

DEPARTURE 30,967.73

BEARING _____

ULTIMATE DEPTH 346.0

ELEVATION 4,094.68

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-141.0	(43%)	Casing (overburden)												
141.0-146.0	5.0	Graphitic schist replaced by fine-grained pyrite and silica. Relict banding at 60° to core. Some reddish-brown sphalerite between 141.0 and 142.0	361	5.0			1.48	2.38						
146.0-151.0	5.0	Pyrite, pyrrhotite and silica containing vestiges of graphitic schist. A little galena and sphalerite present.	362	5.0			1.14	1.23						
151.0-156.0	5.0	Same as previous section. Considerable galena and sphalerite from 154.5 to 156.0.	363	5.0			1.73	2.81						
156.0-161.0	5.0	Graphitic schist replaced by irregular patches of silica, pyrite, galena and sphalerite, closely associated.	364	5.0			3.46	7.32						
161.0-165.0	4.0	Graphitic schist replaced by patches of pyrite with associated reddish-brown sphalerite. Relict banding at 70° to core. Some white quartz-carbonate.	365	4.0			1.50	3.70						
165.0-169.0	1.6	Fragments of graphitic schist with some irregular fine pyrite.	366	4.0			1.38	1.69						
169.0-174.0	2.2	Massive pyrite containing vestiges of graphitic schist. A little reddish-brown sphalerite present with the pyrite.	367	5.0			2.30	3.71						
174.0-179.0	4.4	Massive, fine-grained pyrite in a silica matrix.	368	5.0			1.50	2.10						
179.0-184.0	5.0	Closely-knit pyrite and silica as in previous section.	369	5.0			1.15	0.30						

DIAMOND DRILL RECORD,

HOLE NO. 60

PROPERTY YANCOUDA CREEK, Y. T.

SHEET NUMBER 2 SECTION FROM 184.0 TO 229.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE	FOOTAGE	SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.		AG.	CU.	PB.	ZN.
184.0- 189.0	3.5	Massive pyrite with silica in close association. Core badly broken.	370	5.0			0.71	0.21						
189.0- 194.0	5.0	Pyrite and silica as in previous section. Pyrite massive in part, remainder equigranular in texture.	371	5.0			nil	nil						
194.0- 198.0	4.0	Same as previous section. A few quartz threads.	372	4.0			nil	nil						
198.0- 202.0	1.0	Siliceous groundmass with patches of fine pyrite. Highly fractured.	373	3.0			Tr	0.20						
201.0- 206.0	2.3	Massive pyrite with several patches of grey silica.	374	5.0			nil	nil						
206.0- 210.0	1.0	Fragments of fine pyrite in a grey silica matrix.	375	4.0			nil	nil						
210.0- 214.0	2.5	Silicified sericite schist containing some fine pyrite throughout.	376	4.0			Tr	nil						
214.0- 219.0	4.0	Pyrite and silica as in previous sections. Remnants of graphitic schist from 214.0 to 215.0.	377	5.0			nil	nil						
219.0- 224.0	2.6	Silicified sericite schist containing scattered fine pyrite. Core buttoned in part at about 80°.	378	5.0			Tr	nil						
224.0- 229.0	3.5	Greyish-white sericite schist containing bands of fine pyrite and pyrrhotite in the plane of the shearing at 80° to core. Several small patches of white quartz.	379	5.0			nil	nil						

DIAMOND DRILL RECORD,

HOLE NO. 62

PROPERTY VANCOUVER CREEK, I.T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 427.0

STARTED August 14, 1954

LATITUDE 28,032.69

DATUM 4,000

COMPLETED August 18, 1954

DEPARTURE 32,569.27

BEARING Vertical Hole

ULTIMATE DEPTH 427.0

ELEVATION 4,133.60

DIP _____

PROPOSED DEPTH To test anomaly on strike of main zone.

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0 - 25.0	(76)	Casing (overburden)												
25.0 - 241.0	181.5	Medium greenish-gray sericite schist. Shearing at 50° to core. Several small patches of white quartz containing sections of chlorite and pyrrhotite. A little fine pyrrhotite and pyrite with sparse chalcopyrite from 25.0 to 99.0. Traces of reddish-brown sphalerite at 139.0, 144.0, 210.0 and 230.0. Sphalerite associated with fine pyrrhotite. Some talc present with chlorite in the schist. The schist is gradational to a light gray colour from 178.0 to 241.0 due to decreasing chlorite content.	392	135-140			Tr	0.05						
241.0 - 316.3	50.8	Light gray to greyish-white talc-sericite schist. Shearing at 70° to core. Small patches of pyrrhotite following the schist planes in some cases. Shearing at 80° to core, toward end of section. Some white quartz-carbonate in irregular patterns.												
316.3 - 326.0	1.8	Graphitic schist. Pebbled with dark gray breccia. Shearing obscured.												
326.0 - 353.9	23.8	Strongly talcose sericite schist. Core badly broken soft and sheared at 30° to core. Brecciated in part, the fragments contained in a quartz-carbonate matrix. Shear zone of intense alteration indicated.												
353.9 - 427.0	33.7	Light gray to medium greenish-gray sericite schist. Shearing at 65° to core. A few small patches and threads of pyrrhotite and pyrite.												
Please Note: ^{BY} Caving took place in the zone from 326.0 to 353.9 filling the hole, below this section, with sludge and fragments. It was necessary to aban on the hole at 427.0 for this reason.														

DIAMOND DRILL RECORD,

HOLE NO. 63

PROPERTY VANGORDA CREEK, I. T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 155.0 STARTED August 16, 1954
 LATITUDE 29,354.16 DATUM 4000 COMPLETED August 20, 1954
 DEPARTURE 31,097.41 BEARING _____ ULTIMATE DEPTH 367.0
 ELEVATION 4,112.98 DIP Vertical Hole PROPOSED DEPTH To cross section mineralisation

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
(30) 0.0 - 100.0		Casing (overburden)												
100.0 - 106.0	0.6 M	Fragments of closely-knit massive pyrite and silica	393	6.0			2.05	0.10	416	100-105			1.11	1.78
106.0 - 111.0	0	Sample taken over 6' to first block due to low recovery												
111.0 - 115.0	1.3	Lost core		5.0			5.44		417	105-110			1.54	2.51
111.0 - 115.0	1.3	Fine-grained pyrite, silica galena and sphalerite in close association.	394	4.0			5.18	10.12	418	110-115			5.53	9.11
115.0 - 120.0	4.8	Massive pyrite with irregular patches of white quartz. A little magnetite, pyrrhotite and several irregular threads of galena and sphalerite.	395	5.0			5.10	3.20						
120.0 - 125.0	5.0	Same as previous section with a sharp increase of pyrrhotite from 123.5 to 125.0.	396	5.0			1.31	0.89						
125.0 - 130.0	5.0 M	Massive pyrrhotite containing small irregular patches of white quartz. A little associated chalcopyrite. A little reddish-brown sphalerite between 123.5 and 125.0	397	5.0			2.11	3.58						
130.0 - 135.0	5.0 M	Same as previous section to 132.0. Balance of section is interbanded pyrite, silica, magnetite, galena and sphalerite.	398	5.0			5.27	4.41						
135.0 - 140.0	5.0 M	Sulphides, magnetite and silica as in previous section. Relict banding at 50° to core.	399	5.0			0.74	1.23						
140.0 - 145.0	5.0	Same mineralization as previous two sections	400	5.0			0.43	0.40						
145.0 - 150.0	5.0	Irregular patterns of pyrite and pyrrhotite in a silicified matrix of sericite schist. Sparse chalcopyrite present in fractures.	401	5.0			0.98	0.71						
150.0 - 155.0	5.0	Pyrite and silica with sparse pyrrhotite, magnetite and silica.	402	5.0			0.94	1.50						

DIAMOND DRILL RECORD,

HOLE NO. 64

PROPERTY VANCORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 225.0

STARTED August 18, 1954

LATITUDE 29,271.75

DATUM 4,000

COMPLETED August 25, 1954

DEPARTURE 30,376.41

BEARING _____

ULTIMATE DEPTH 516.0

ELEVATION 4,071.61

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0 - 170.0	(41.6) 5.0	Casing (Overburden)												
170.0 - 158.0		Graphitic schist with some dark grey interbanded sericite schist. Shearing at 80° to core. One botton of pyrite at 145.0.												
158.0 - 177.7	16.3	Light grey talcose sericite schist. A little interbanded chlorite and white quartz. Shearing at 80° to core.												
177.7 - 178.9	1.2	Massive pyrite with some fine pods and threads of magnetite following the relief banding at 50° to core	423	1.2			1.97	2.31						
178.9 - 200.8		Talc sericite schist. A little interbanded chlorite and some white quartz patches. Shearing at 80° to core.												
200.8 - 205.0	4.2	Graphitic schist and silica with some pyrite to 202.5 balance of sample consists of fine pyrite in a siliceous matrix with some closely associated galena and sphalerite. Sparse pyrite.	424	4.2			1.07	3.62						
205.0 - 210.0	3.2	Massive pyrite with vestiges of kaolinized sericite schist. Sparse magnetite, galena and sphalerite. Relief banding at 80° to core.	425	3.0			0.17	3.82						
210.0 - 215.0	3.6	Same mineralization as previous section.	426	3.0			0.06	0.40						
215.0 - 220.0	5.0	Massive pyrite with vestiges of sericite and graphitic schist to 217.5; balance of sample is massive pyrite and pyrrhotite with some closely associated galena and sphalerite.	427	5.0			1.40	2.08						
220.0 - 225.0	5.0	Graphitic schist, crumpled and carbonatized containing irregular patches of massive pyrite and pyrrhotite.	428	5.0			0.26	0.68						

DIAMOND DRILL RECORD,

HOLE NO. 3

PROPERTY VANDORA CREEK, I. T.

SHEET NUMBER 2 SECTION FROM 225.0 TO 380.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS				
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.	
225.0 - 296.0	31.0	Light grey sericite schist. Slightly talcose; a little white quartz containing streaks of chlorite at the margins. Sparse pyrite and pyrrhotite.													
296.0 - 301.0	1.4	Graphitic schist with interbanded grey silica. Scattered fine pyrite.	429	5.0			1.67	0.22							
301.0 - 306.0	2.3	Graphitic schist as in previous section. A little reddish-brown sphalerite and galena.	430	5.0			0.70	ml							
305.0 - 309.0	1.4	Graphitic schist with fine pyrite and silica. Shearing at 60° to core.	431	4.0			1.20	0.70							
309.0 - 315.2	21.3	Light grey talcose sericite schist replaced by irregular patches of white quartz. Shearing at 80° to core.													
315.2 - 350.0	2.4	Graphitic schist with some interbanded silica, pyrite and pyrrhotite. Shearing at 70° to core.	432	4.8			1.14	1.20							
350.0 - 355.0	3.1	Graphitic schist mineralized as in previous section. A little reddish-brown sphalerite and galena.	433	5.0			2.99	2.02							
355.0 - 360.0	5.0	Graphitic schist with interbanded silica, pyrite, reddish-brown sphalerite and galena. Shearing at 80° to core.	434	5.0			2.59	1.43							
360.0 - 365.0	5.0	Graphitic schist brecciated in part and replaced by pyrite, galena, sphalerite and silica.	435	5.0			1.09	2.50							
365.0 - 370.0	5.0	Graphitic schist showing replacement as in previous section. Shearing at 60° to core.	436	5.0			0.98	1.72							
370.0 - 375.0	5.0	Massive pyrite with a little associated magnetite, sphalerite and galena. Remnants of sericite schist with interbanded graphitic schist from 373.0 to 374.5	437	5.0			1.34	0.85							
375.0 - 380.0	2.2	Graphitic schist buttons containing interbanded silica and disseminated pyrite. Massive pyrite from 375.0 to 375.8.	438	5.0			1.93	1.24							

DIAMOND DRILL RECORD,

HOLE NO. 66

PROPERTY YAMORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 341.0

STARTED August 21, 1954

LATITUDE 29,637.36

DATUM 4,000

COMPLETED August 26, 1954

DEPARTURE 31,123.07

BEARING _____

ULTIMATE DEPTH 341.0

ELEVATION 4,095.94

DIP Vertical Hole

PROPOSED DEPTH To test for lateral extent of mineralization.

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0 - 90.0	(27)	Casing (Overburden)												
90.0 - 111.0	9.5	Light grey sericite schist. Shearing at 80° to core; core badly broken on the shear planes.												
111.0 - 116.0	4.0	Greenish-grey, chloritic sericite schist. A few threads and small patches of pyrrhotite; sparse reddish-brown sphalerite in association.	440	5.0										
116.0 - 121.0	4.6	Same as previous section. A little pyrite in association.	441	5.0										
121.0 - 126.0	1.8	Light grey silicified sericite schist containing a little pyrrhotite and chlorite	442	5.0										
126.0 - 129.0	2.0	Irregular patches of pyrrhotite, chlorite and sparse sphalerite in sericite schist.	443	3.0										
129.0 - 190.0	37.6	Light grey to medium grey sericite schist with a little interbanded graphitic schist at irregular intervals. Shearing at 80° to core. Several small patches of white quartz and some associated chlorite.												
190.0 - 341.0	44.4	Light grey to medium grey sericite schist. Shearing at 50° to core. Several small patches of white quartz.												
		PLEASE NOTE: Core size reduced from A to F at 236.0 due to need for casing off some caving with E casing.												
		End of hole at <u>341.0</u>												
		Average recovery <u>41.4%</u>												
		Recovery in mineralizations <u>68.9%</u>												
		Size of cores: Ax (1 3/16") to 236.0 Ex (7") to 341.0 end.												

DIAMOND DRILL RECORD,

HOLE NO. 72

PROPERTY YANGORDA CREEK T. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 70.0

STARTED September 1, 1924

LATITUDE 28,767.35

DATUM 1000

COMPLETED Sept. 4, 1924

DEPARTURE 31,330.54

BEARING _____

ULTIMATE DEPTH 399.0

ELEVATION 4,060.15

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0 - 30.6	(9.3%)	Casing (overburden)												
30.6 - 35.0	2.4	Fragments of fine grained pyrite with some small patches of magnetite and a little galena and sphalerite.	463	4.4			2.15	2.13						
35.0 - 40.0	4.9	Massive pyrite containing several bands of magnetite roughly paralleling the relief banding at 80° to core; a little galena and sphalerite in close association with the magnetite.	464	5.0			0.95	0.60						
40.0 - 45.0	4.8	Some mineralization as previous section. Magnetite bands more irregular than before.	465	5.0			0.70	0.93						
45.0 - 50.0	5.0	Massive pyrite with irregular small patches of silica, magnetite galena and sphalerite in close association.	466	5.0			3.12	5.33						
50.0 - 55.0	5.0	Equigranular fabric of pyrite, silica, galena, sphalerite and a little magnetite. Good lead-zinc content.	467	5.0			2.79	6.41						
55.0 - 60.0	5.0	Some mineralization as previous section to 58.0 Balance of section is massive pyrite with several small stringers of white quartz.	468	5.0			3.43	2.82						
60.0 - 65.0	5.0	Equigranular pyrite and silica containing a few stringers of closely-associated magnetite and reddish-brown sphalerite.	469	5.0			1.53	1.91						
65.0 - 70.0	5.0	Massive pyrite with some irregular patches of silica and magnetite. A little reddish-brown sphalerite and galena.	470	5.0			0.75	0.83						

DIAMOND DRILL RECORD,

HOLE NO. 72

PROPERTY VANGUARD CREEK

SHEET NUMBER 3

SECTION FROM 275.0 TO 295.0

STARTED _____

LATITUDE _____

DATUM _____

COMPLETED _____

DEPARTURE _____

BEARING _____

ULTIMATE DEPTH _____

ELEVATION _____

DIP _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
235.0 - 240.0	5.0	Medium gray sericite schist with irregular patches of pyrite and pyrrhotite. Sparse magnetite galena and sphalerite.	479	5.0			0.01	0.50						
240.0 - 215.0	4.4	Massive pyrite containing small patches of magnetite and vestiges of sericite schist.	480	5.0			0.22	0.10						
215.0 - 250.0	4.5	Graphitic schist with a little associated sericite schist. Some irregular patches of silicea and pyrite. Shearing contorted.	481	5.0			0.25	0.50						
250.0 - 255.0	3.9	Graphitic schist mineralized with pyrite as in previous section. Shearing contorted; bands crumpled in part.	482	5.0			0.28	0.41						
255.0 - 260.0	3.9	Graphitic schist replaced by considerable fine pyrite. Shearing at 65° to core.	483	5.0			0.02	0.20						
260.0 - 265.0	4.0	Graphitic schist with interbedded silicea, pyrite and pyrrhotite.	484	5.0			0.02	0.20						
265.0 - 270.0	3.0	Graphitic schist mineralized as in previous section. Shearing crumpled in part.	485	5.0			0.02	nil						
270.0 - 275.0	2.7	Graphitic schist finely laminated with pyrite and silicea at 60° to the core.	486	5.0			0.01	nil						
275.0 - 280.0	3.9	Graphitic schist and pyrite as in previous section. Traces of reddish-brown sphalerite.	487	5.0			2.82	7.48						
280.0 - 285.0	5.0	Graphitic schist laminated in the shear planes with pyrite, silicea and a little galena and sphalerite.	488	5.0			2.32	4.53						
285.0 - 290.0	3.0	Same as previous section changing to silicea and pyrite at 288.0, for balance of section.	489	5.0			1.48	2.62						
290.0 - 295.0	3.3	Pyrite and silicea from 290.0 to 293.0; some interbedded graphitic schist with pyrite, silicea and sparse sphalerite to 295.0.	490	5.0			1.56	3.21						

DIAMOND DRILL RECORD,

HOLE NO. 7

PROPERTY VANGORDA CREEK Y.T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 306.5

STARTED Sept. 4, 1954

LATITUDE 26,106.62

DATUM 4,000

COMPLETED Sept. 7, 1954

DEPARTURE 34,294.33

BEARING _____

ULTIMATE DEPTH 363.0

ELEVATION 4,066.50

DIP Vertical Hole

PROPOSED DEPTH To test anomaly No. 23

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-														
11.0 (3.3)		Casing (Overburden)												
11.0- 16.0	2.6	Fragments of pyrite and pyrrhotite in a silica matrix. Sparse reddish brown sphalerite and chalcopyrite.	515	5.0			.75	1.19						
16.0- 21.0	3.9	Vestiges of graphitic schist and silica containing fine disseminated pyrite and a little galena and sphalerite.	516	5.0			.33	.50						
21.0- 26.0	2.6	Fragments of massive pyrrhotite and silica. A few small patches of magnetite, in the pyrrhotite.	517	5.0			.89	1.81						
26.0- 31.0	3.5	Massive pyrrhotite and pyrite containing vestiges of graphitic schist to 29.0; vestiges of sericite schist in balance of section.	518	5.0			.50	1.01						
31.0- 196.0	82.7	Medium greenish-gray sericite schist. Fissile shearing, averaging 50° to core. A few small patches of white quartz. Shearing changes to 70° to core toward end of section.												
196.0- 236.0	30.4	Medium gray sericite schist. Fissile shearing at 60° to the core.												
236.0- 255.8	19.8	Medium greenish-gray sericite schist containing a few small stringers and patches of pyrite and pyrrhotite. Sparse sphalerite and chalcopyrite.												
255.8- 258.1	2.3	Greenish-gray sericite schist replaced by small irregular patches of closely-associated pyrite, pyrrhotite, chlorite, quartz and sparse magnetite. & sphalerite.	519	2.3			Tr	int						
258.1- 306.5	39.0	Medium greenish-gray sericite schist shearing at 60° to core. A few threads of pyrite and pyrrhotite between 258.1 and 264.0; sparse pyrite and pyrrhotite in balance of section.												

DIAMOND DRILL RECORD,

HOLE NO. 74

PROPERTY Yagorda Creek, Y.T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 296.0

STARTED Sept. 15, 1954

LATITUDE 28,745.27

DATUM 4,000

COMPLETED Sept.

DEPARTURE 31,614.13

BEARING _____

ULTIMATE DEPTH _____

ELEVATION 4,075.15

DIP Vertical Hole

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-27.0 (8.2)		Casing (Overburden)												
27.0-191.0	69.0	Medium grey sericite schist. Slight greenish cast due to interbanded chlorite. Shearing at 60° to core. A few threads of pyrite and pyrrhotite with sparse reddish brown sphalerite between 141.0 and 191.0, not worthy of assay. One 1/2" patch of quartz and reddish brown sphalerite at 190.8. Core badly broken from 27.0 to 111.0												
191.0-206.0	13.7	Medium grey sericite schist. A few small patches of white quartz. Shearing at 80° to core.												
206.0-209.5	3.5	Massive fine-grained pyrrhotite containing vestiges of sericite schist. Sparse galena, sphalerite and chalcopyrite.	520	3.5			.78	.80						
209.5-213.0	2.7	Same mineralisation as previous section. Vestiges of graphitic schist and sericite schist.	521	3.5			1.23	2.30						
213.0-253.0	34.3	Medium grey sericite schist. Fissile shearing at 70° to core. A few threads of pyrrhotite, and a little interbanded chlorite. Numerous small pink garnets from 231.5 to 234.0												
253.0-296.0	25.4	Medium grey sericite schist. From 279.0 to 281.0 considerable chlorite, pyrrhotite and small garnets replacing the schist. Please note: At 296.0, reduction was made to E core due to the need for casing off some casing with E casing.												

DIAMOND DRILL RECORD,

HOLE NO. 75

PROPERTY VANDORDA CREEK I.T.

SHEET NUMBER 1 SECTION FROM 0.0 TO 72.0
 LATITUDE 28,616.91 DATUM 4,000
 DEPARTURE 31,458.92 BEARING _____
 ELEVATION 4,069.76 DIP Vertical Hole

STARTED Sept. 5, 1954
 COMPLETED Sept. 10, 1954
 ULTIMATE DEPTH 436.0
 PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-25.0		Casing (overburden)												
25.0-30.0		Lost core												
30.0-36.0	0.9	Porous pyrite in a silica matrix. Some galena and sphalerite present in fragments. Sludge sent for assay due to low core recovery.							337 338	25-30 30-35			Tr 2.04	nil 2.78
36.0-38.0	1.2	Fragments of sulphide mineralization containing pyrite, galena and sphalerite in a silica matrix. Relict banding at 90° to core.	322	2.0			2.74	5.41	339	35-40			4.89	6.04
38.0-39.5	1.5	Porous fragments of sulphide containing pyrite and silica, and good galena and sphalerite mineralization.	323	1.5			7.15	7.53						
39.5-50.0	1.4	Fragments of massive pyrite and silica containing a little galena and sphalerite. Sludges sent for assay due to low core recovery.							340 341	40-45 45-50			0.95 4.28	1.20 4.88
50.0-55.0	4.5	Massive pyrite in a silica matrix. A little galena and sphalerite.	324	5.0			87	nil						
55.0-61.0	3.3	Fragments of pyrite and silica containing a little galena and sphalerite. Sample taken over 6.0' due to low core recovery.	325	6.0			86	20						
61.0-66.0	2.2	Banded pyrite and silica containing vestiges of graphitic schist. Porous in part; relict banding at 60° to core.	326	5.0			0.45	nil						
66.0-72.0	4.2	Fragments of massive pyrite in a silica matrix. Sparse galena and sphalerite.	327	6.0			0.52	0.60						

DIAMOND DRILL RECORD,

HOLE NO. 77

PROPERTY YANFORDA CREEK, T. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 166.0

STARTED September 10, 1954

LATITUDE 28,589.62

DATUM L.O.O

COMPLETED September 13, 1954

DEPARTURE 31,742.41

BEARING _____

ULTIMATE DEPTH 348.0

ELEVATION 4,089.86

DIP Vertical Hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-31.0	(9.4)	Casing (overburden)												
31.0 - 36.0	0.7	A few fragments of pyrite in a silica matrix. Sludge sent due to low recovery.							560	30-35			0.42	0.41
36.0 - 41.0	0.9	Same mineralization as previous section, replaced by a little white quartz. Sludge sent due to low recovery.								35-40	No Water Return			
41.0 - 46.0	1.3	Pyrite and silica with a little reddish-brown sphalerite. Relict banding at 90° to core.	545	5.0			2.07	8.33						
46.0 - 51.0	2.7	Pyrite and silica well mineralized with galena and sphalerite. A little magnetite in the relict banding at 60° to core. Porous from 50.0 to 51.0.	546	5.0			3.84	6.10	561	40-50			0.36	1.80
51.0 - 56.0	1.7	Pyrite and silica containing strong, fine-grained equigranular galena and sphalerite.	547	5.0			2.46	8.37						
56.0 - 61.0	3.0	Pyrite and silica with decreasing sphalerite and galena relative to previous section. A few small patches of white quartz.	548	5.0			2.85	2.22	562	50-60			5.11	10.29
61.0 - 63.0	0.6	A few fragments of pyrite and silica. Sludge sent due to low recovery.								60-70			1.12	1.11
63.0 - 67.0	0.6	Fragments of pyrite and silica as in previous section. Sludge sent due to low recovery.												
67.0 - 72.0	5.0	Light grey sericite schist containing irregular patches of pyrite, pyrrhotite and sparse sphalerite. Shearing obscured by the replacing sulphide.	549	5.0			0.31	0.60			Sludges small due to weak water return.			
72.0 - 166.0	53.7	Medium grey sericite schist. A little fine pyrrhotite and pyrite, and several irregular patches of white quartz. Shearing at 80° to core.												

DIAMOND DRILL RECORD,

HOLE NO. 78

PROPERTY VANORONDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 111.0

STARTED September 11, 1954

LATITUDE 28,442.61

DATUM 4,000

COMPLETED September 14, 1954

DEPARTURE 31,592.11

BEARING _____

ULTIMATE DEPTH 346.0

ELEVATION 4,076.06

DIP Vertical hole

PROPOSED DEPTH To cross section mineralization.

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0 - 22.0		(47) Casing (Overburden)												
22.0 - 53.0	4.0	Fragments of sericite schist and white quartz.							581	50-55			0.28	0.91
53.0 - 66.0	2.2	Light gray sericite schist, badly broken. Two fragments of closely-associated pyrite, galena and sphalerite at 62.0. Sludge sent for assay.							582	55-60			0.42	0.82
									583	60-65			1.70	5.21
66.0 - 71.0	5.0	Pyrite and silica well mineralized with galena and sphalerite. Vestigial banding at 10° to core.	550	5.0			2.66	11.21						
71.0 - 76.0	5.0	Massive pyrite with a few small patches of white quartz. A few threads of galena and sphalerite.	551	5.0			2.40	2.10						
76.0 - 81.0	5.0	Pyrite and silica closely associated with fine galena and sphalerite. A few irregular patches of white quartz.	552	5.0			0.62	1.02						
81.0 - 86.0	5.0	Fine grained pyrite and silica, containing irregular threads of white quartz. Sparse galena and sphalerite.	553	5.0			1.09	0.59						
86.0 - 91.0	5.0	Massive pyrite and silica with some associated pyrrhotite and white quartz.	554	5.0			0.42	0.10						
91.0 - 96.0	5.0	Pyrite and silica with a little galena and reddish-brown sphalerite; a little pyrrhotite. Talcose sericite schist from 94.0 to 96.0.	555	5.0			0.78	2.91						
96.0 - 101.0	2.6	Remnants of light gray sericite schist in massive pyrite; a little sphalerite and magnetite in the pyrite.	556	5.0			0.39	1.23						
101.0 - 106.0	2.5	Massive pyrite containing remnants of sericite schist. Sparse chalcopyrite.	557	5.0			0.28	0.60						
106.0 - 111.0	3.5	Massive pyrite with small patches of white quartz and remnants of sericite schist.	558	5.0			0.04	0.30						

16' - 8.42

DIAMOND DRILL RECORD,

HOLE NO. 78

PROPERTY VANGORDA CREEK, V. T.

SHEET NUMBER 2 SECTION FROM 211.0 TO 252.0 STARTED _____

LATITUDE _____ DATUM _____ COMPLETED _____

DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____

ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
111.0 - 113.0	1.6	Irregular patches of pyrite and pyrrhotite in sericite schist.	559	2.0			0.03	0.98						
113.0 - 152.0	24.2	Light gray to medium gray sericite schist with a little interbanded graphitic schist. Shearing at 80° to core. A few small patches of pyrite.												
156.0 - 203.0	27.0	Medium gray sericite schist containing short sections of interbanded graphitic schist. Shearing at 80° to core.												
203.0 - 221.3	19.9	Graphitic schist interbanded with gray silica. Shearing crenulated in part but averaging about 80° to core. Some pyrite disseminated throughout.												
221.3 - 226.0	4.5	Graphitic schist interbanded with gray silica. Considerable fine pyrite and a few small stringers of pyrrhotite - chalcopyrite.	560	5.0			0.25	ml						
226.0 - 231.0	5.0	Graphitic schist with mineralization similar to previous section.	567	5.0			0.33	ml						
231.0 - 236.0	5.0	Graphitic schist interbanded with silica and replaced by irregular patches of pyrite and small patches of pyrrhotite and small stringers of pyrrhotite - chalcopyrite.	568	5.0			0.53	0.20						
236.0 - 237.6	1.6	Graphitic schist and interbanded silica mineralized as in previous section.	569	1.6			0.31	0.50						
237.6 - 243.0	4.4	Graphitic schist and silica with fine interbanded pyrite. Shearing at 80° to core.	570	4.4			0.59	0.89						
243.0 - 247.0	5.0	Graphitic schist, mineralized as in previous section. Shearing contorted.	571	5.0			0.36	ml						
247.0 - 252.0	5.0	Crenulated graphitic schist and silica containing a little pyrrhotite and sparse sphalerite in strong pyrite mineralization.	572	5.0			0.22	0.20						

DIAMOND DRILL RECORD,

HOLE NO. 3

PROPERTY YANORON CREEK, Y. T.

SHEET NUMBER 3 SECTION FROM 252.0 TO 344.0

LATITUDE _____ DATUM _____

DEPARTURE _____ BEARING _____

ELEVATION _____ DIP _____

STARTED _____

COMPLETED _____

ULTIMATE DEPTH _____

PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
52.0 - 57.0	1.7	Graphitic schist, crenulated and interbanded with silica. Some pyrite and pyrrhotite present.	573	5.0		0.33 0.33		Tr						
57.0 - 62.0	2.3	Same as previous section to 59.0. Sericite schist fragments from 259.0 to 262.0.	574	5.0			0.59	0.48						
62.0 - 67.0	1.4	Talcose sericite schist containing several irregular patches of pyrrhotite. Shearing at 80° to core.	575	5.0			0.20	nil						
67.0 - 72.0	1.0	Crenulated graphitic schist and silica. Some pyrite and pyrrhotite disseminated throughout.	576	5.0			0.56	0.20						
72.0 - 77.0	1.0	Mainly grey silica and pyrite with some interbanded graphitic schist. A little pyrrhotite and chalcopyrite. Vestigial banding at 80° to core.	577	5.0			0.67	nil						
77.0 - 82.0	1.4	Grey silica replaced by irregular patches and threads of pyrite. Vestigial banding at 70° to core. Some galena and sphalerite from 281.0 to 282.0.	578	5.0			0.70	1.02						
82.0 - 87.0	2.8	Same as in previous section. Sparse galena and sphalerite.	579	5.0			0.03	1.31						
87.0 - 92.0	3.7	Banded silica and pyrite to 289.8. Graphitic schist and silica containing disseminated pyrite and pyrrhotite from 283.2 to 292.0	580	5.0			0.39	nil						
92.0 - 98.0	2.4	Fragments of brittle graphitic schist. A few quartz threads and sparse pyrite.												
98.0 - 104.0	17.0	Medium grey sericite schist. Shearing at 80° to core. A few quartz threads present.												
		End of hole at 344.0												
		Size of cores Ax (1 3/16")												
		Average recovery 58.9%												

Recovery in mineralisation 82.9%

DIAMOND DRILL RECORD,

HOLE NO. 80

PROPERTY YANFORDA CREEK, Y. T.

SHEET NUMBER 1

SECTION FROM 0.0 TO 68.0

STARTED September 14, 1954

LATITUDE 28,437.85

DATUM 4,000

COMPLETED September 18, 1954

DEPARTURE 31,873.12

BEARING _____

ULTIMATE DEPTH 351.0

ELEVATION 4,095.40

DIP Vertical hole

PROPOSED DEPTH To cross-section mineralization

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
0.0-26.0	(1.9)	Casing (overburden)												
26.0 - 31.0	1.6	Fragments of greyish-white kaolinized sericite schist with irregular patches of pyrite. Relict banding at 80° to core.	584	5.0			0.28	nil	605	25-30			0.89	1.43
31.0 - 36.0	3.0	Fine-grained pyrite with irregular patches of grey silica and remnants of kaolinized sericite schist. A little magnetite from 35.5 to 36.0.	585	5.0			0.64	0.90	606	30-35			2.06	2.68
36.0 - 41.0	3.8	Same as previous section to 36.0. Balance of section is an equigranular fabric of pyrite, silica galena and sphalerite.	586	5.0			1.48	2.83						
41.0 - 46.0	2.9	Closely-associated pyrite, silica, galena and sphalerite. Rock is semi-porous.	587	5.0			3.09	8.73						
46.0 - 51.0	3.7	Pyrite, silica, galena and sphalerite in a fine-grained fabric.	588	5.0			2.29	7.83						
51.0-53.0	1.5	Same mineralization as previous section.	589	2.0			3.29	9.65						
53.0-58.0	0.1	One fragment of pyrite, galena, sphalerite and silica. Sludge sent for assay due to low recovery.							607	30-35			1.82	7.24
58.0-63.0	2.7	Pyrite and silica streaked with galena and sphalerite. Considerable magnetite from 62.0 to 63.0	590	5.0			3.54	4.82						
63.0-68.0	4.8	Pyrite with irregular patches of silica in about equal quantity.	591	5.0			0.61	0.38						

DIAMOND DRILL RECORD,

HOLE NO. 80

PROPERTY VANGORDA CREEK, T. T.

SHEET NUMBER 3 SECTION FROM 169.8 TO 219.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
169.8 - 174.0	4.2	Graphitic schist interbanded with gray silica. Shearing at 80° to core. Considerable pyrite with some associated galena and sphalerite.	603	4.2			1.23	1.12						
		<u>PLEASE NOTE:</u> Due to casing conditions, E casing was run, and reduction to B core made at 174.0.												
174.0 - 179.0	5.0	Graphitic schist with interbanded silica in variable amounts. Considerable fine-grained pyrite. Sparse pyrrhotite and sphalerite.	604	5.0			0.20	nil						
179.0 - 184.0	5.0	Same mineralization as previous section	608	5.0			0.08	Tr						
184.0 - 189.0	2.9	Fine-grained pyrite and silica with vestiges of graphitic schist. Some threads and small patches of pyrrhotite.	609	5.0	↑		0.06	nil						
189.0 - 194.0	5.0	Massive pyrite containing vestiges of graphitic schist. Several small patches of magnetite. Sparse sphalerite.	624	5.0			0.22	nil						
194.0 - 199.0	4.0	Pyrite and silica with vestiges of graphitic schist. Banding at 80° to core.	625	5.0			0.34	0.10						
199.0 - 204.0	5.0	Graphitic schist with a little interbanded pyrite.	626	5.0			0.22	nil						
204.0 - 209.0	3.1	Graphitic schist and interbanded silica containing considerable pyrite, galena and sphalerite.	627	5.0			2.44	6.63						
209.0 - 214.0	4.8	Finely-banded graphitic schist, silica and pyrite at 80° to core. A little reddish-brown sphalerite.	628	5.0			0.98	2.22						
214.0 - 219.0	3.4	Same mineralization as previous section.	629	5.0			0.28	0.20						

DIAMOND DRILL RECORD,

HOLE NO. 480

PROPERTY VANGORDA CREEK, T. T.

SHEET NUMBER 4 SECTION FROM 219.0 TO 274.0 STARTED _____
 LATITUDE _____ DATUM _____ COMPLETED _____
 DEPARTURE _____ BEARING _____ ULTIMATE DEPTH _____
 ELEVATION _____ DIP _____ PROPOSED DEPTH _____

DEPTH FEET	CORE RECOV	DESCRIPTION	CORE SAMPLE NO.	FOOTAGE	CORE ASSAYS				SLUDGE SAMPLE		SLUDGE ASSAYS			
					AG.	CU.	PB.	ZN.	NO.	FOOTAGE	AG.	CU.	PB.	ZN.
219.0 - 224.0	3.5	Considerable fine pyrite replacing banded graphitic schist and silica.	630	5.0			0.31	0.39						
224.0 - 229.0	4.9	Same mineralization as previous section. Banding crenulated.	631	5.0			0.62	0.78						
229.0 - 234.0	5.0	Considerable pyrite and sparse sphalerite replacing graphitic schist banded with silica. Shearing at 45° to core.	632	5.0			0.42	0.40						
234.0 - 239.0	5.0	Graphitic schist containing irregular patches of silica and pyrite.	633	5.0			1.11	1.12						
239.0 - 244.0	5.0	Same as previous section. A little reddish-brown sphalerite.	634	5.0			2.99	3.44						
244.0 - 249.0	5.0	Pyrite and silica in graphitic schist to 246.0. Balance of section is massive pyrite with patches of grey silica.	635	5.0			0.98	1.20						
249.0 - 254.0	5.0	Pyrite and silica with vestiges of graphitic schist. A little sphalerite. Banding at 80° to core.	636	5.0			1.00	4.17						
254.0 - 259.0	5.0	Graphitic schist and silica, largely replaced by fine-grained pyrite.	637	5.0			1.78	2.03						
259.0 - 264.0	4.6	Crenulated graphitic schist to 261.0. Balance of section is equigranular pyrite, silica, sphalerite (reddish-brown) and galena.	638	5.0			1.78	3.03						
264.0 - 269.0	4.5	Pyrite, silica, sphalerite and galena to 266.0. Balance of section is graphitic schist and silica with pyrite and sparse sphalerite.	639	5.0			2.78	7.41						
269.0 - 274.0	5.0	Graphitic schist interbanded with considerable grey silica. Some interbanded pyrite and a little associated galena and sphalerite.	640	5.0			2.33	6.29						

