

Kasawa.

Dec 12/02.

007308

- Coast Plate Complex.
- Tert. Skulkum intrusives - re Au WH Skulkum,
- WH complex intrusives
- Nisling Terrane.
- very devoid of mafic.
- Nisling ^{metaseds.} metamorphics - in soup of intrusives -> equiv to Greenstone.
- Cu mo porphs skarns, PZ veins near Tert.
- Tert stocks popping up throughout on east side.
- gabro ^(#11) has likely equiv. to SE on margins of

~~MA~~ Mavsh. - 21g over 2'
~~Tagish Narrows.~~

Cache Ck., Lebarge. Whitehorse suite.
+ Nansen.

Prospector Mt.

Cache Creek. oceanic no ark sign. in this area?

RGS-response is poor in the area.

Signif Fault Through Lake.

- good exp steep ground.
 - stuff mapped as Nansen is mafic volc in Cache Ck. - what kind of mafic volc.
 - Fair bit of Au in Cks from SW side.
 - Kistiwite is host to Au. Rossbank property.
-

Tagish Narrows.

- placer claims.
- small intrusions in area.

Marsh Pathway

| | fact. | |
|-------|-------|---|
| 2 h | 2 | 1 |
| 1 h-m | 1 | 2 |
| 3 | 3 | 3 |
| | 4 | 4 |
| | 6 | 5 |
| | 5 | 6 |

2
1

Sinafu.

no claims in area.

94-97 mapped Cordy & Stevens.

250 scale
mapping

Catche Ct. up to early R.

mainly chert + shale stst gnt.

- get.

Wiel R granitic rcks.

→ any v. comp.

Photos -

White Goat Mt. - gossams.

Fairly quiet mag. bot slugs of VM rocks.
under the hST.

Srafu Rating.

Tract.

1L

2L

4H

~~3L~~

3L

7m

—

—

—

4

5

~~7~~

6

3

~~2~~

2

1

1

4✓

5✓

7 great occ to south. no data.

no geoch. { 6 no geochem.

8 pod cr ^{Av} not reprod.

3 no geochem. intrusives nearby.

2 LST. no chem. good exp.

1 LST no chem good exp.

4

1

5

7

8

3

6

2

1

Scotty CK.

Kloridite schist - GRT. some QZ stringers S&T.
stromatolite showing QZ vein. - no results.
logline. is to the North. - 20 Km.

Nasir - Kloridite schist.
Wt Suite. - Cretaceous. - logline to North.
Nisling range. - GRT. Tertiary.
ESGA. early R. logline suite

Scotty. Rating.

- | | | |
|---|---|-------------------|
| | | 6 ✓ |
| | | 8 ✓ |
| 1 | <u>Cret.</u> very weak plut. <u>Au</u> sig. mag hi assoz intrusions have Au in silt. | |
| 2 | <u>Cret.</u> very weak. plut <u>Au</u> sig mag " " " Au weaker than #1 but "busy" coverage. | 1 } 2 } 5 } |
| 3 | weak Cu Mo Ni Sb Sn U Zn. <u>no Au</u> poor coverage | 4 } 3 } 7 } |
| 4 | Cu, Mo, W + Sb Sn Zn. but <u>no Au</u> . R plutons @ Minto. better than 3 b.c. of this. | |
| 5 | <u>Cret</u> is intrusive has 2 Au anomalies has strong spot mag hi. | |
| 6 | Klorlike schist. same felsic. N-dipping mag feature. No coverage S. side. Au Cu south Cu Sn Zn. | |
| 8 | Similar to 6 but weaker. mag & geochron. | |

Walsley.

terrible silt responses.

- windy w/ kindly oceanic.
- Karmack's Group. - south side
- Nasira Cup. to North.

Find out from Poggis

- Shaded relief.
- satellite photo.
- good topo.

Wellesley

- | | |
|--|---|
| <p>2 Windy 1/2 OC. weak to mod mag. Au As Sb & Cu Zn. geochem. - mafics & sediments - soils have As Sb Au.</p> | 2 |
| <p>3 poorly exp. strong mag Windy weak Au. Cu Zn w. but all over.</p> | 3 |
| <p>4) 1/2 Carnacks samples of <u>ore</u> <u>dead</u>. no chance in in Windy</p> <p style="text-align: right; margin-right: 50px;"><small>veins poor exposure</small></p> <p style="text-align: right; margin-right: 50px;"><small>no no OC. all <u>quat</u></small></p> | ↓ |
| <p>1 no exposure. in Sed Down Windy if rocks. As Cu</p> | 7 |
| <p>5 both Au/Fu</p> | 4 |
| <p>6 " Some exposure</p> | |
| <p>7 is no exposure. & 2 Cu. but besshi potential Volc. neck hi heat for epithermal.</p> | 6 |
| | 5 |

Pick Handle.

Nishig.
Derah Fault.

CPS₁ Skoli Vale Perm → Permian,
tuff box argill. agg. aug. phytic bed. And.
gwke cgl

PTKR₁ Klone suite UMs'

DMN₃ Dev. Miss QZT. FD augen. sch & mm cug.

Windy McKinley'