






FIG. 2

PROSPECTING LOCATIONS - GUNTER LISHY - JULY 1969.






4 miles = 1 inch.

SUMMARY

- ① - Small amounts scheelite both sides of valley at  contact.  
- Barite in  north of valley.
- ② - Barite north & east in   
- Galena in float, small valley south-east in 
- ③ - Chalcopyrite, pyrrhotite, pentlandite? southwards near  contact
- ④ - Nothing.










LEGEND.

-  CRETACEOUS - Gneiss/diorite
-  UPPER DEVONIAN/  
LOWER MISSISSIPPIAN - Shale, Conglom.
-  SILURIAN/  
DEVONIAN - Dolomite, chert.
-  MIDDLE/UPPER  
CAMBRIAN - Shale, phyllite, Slate
-  LOWER  
CAMBRIAN - Quartz biotite/  
chlorite schist.

105 F - Prospect Field Note

014244



-  Alteration Rock
-  Phyllite
-  Dolomite or Limest.
-  Black Shale
-  Light Grey Dol. Slabby
-  Mineralized

At and near main contact the local rock has altered to Hornfels, Quartzite, Chert, Phyllite and shows local Schistose character. Some sections are mineralized with Pyrrhotite, Pyrite, Arsenic, Copper, Scheelite, Molybdenite and Zink, (except Pyrrhotite accompanied by scheelite none of these) minerals appear in quantity. Barite occurring as veins, stringers and impregnations in Dolomite north east of Camp covers quite an area and might be of interest. All the area outlined, north east of the main contact, has been cut by numerous intrusive dikes with little and mostly no contact alteration.

Locality A9, ... Brownish black Pyrrhotite found in narrow strips at contact between Diaritic intrusive and Phyllite contains some Scheelite.


A7, ... Pyrrhotite occurring in altered rock (Remnant) has traces of Molybdenite, no Scheelite.

B, ... Little below A7 in and around creek Scheelite and Cu. occurs in Pyrrhotite

C, ... Traces of Scheelite in altered rock.


**CAMP I**  
 Dog Camp  
 G. Lohy

P.b. occurs on steep talus slope in Dolomite and Cherty rock found so far only in form of float. Mineralization appears to be spotty and is marked by light brown areas of leaching on a grey background of the host rock. Some Barite occurs nearby.










ALL OF AREA SKETCHED IS CUT BY NUMEROUS INTERSECTING DICES



CAMP 3

G. Lishy

