R.T. Oex

Lshy Prospecting Camp No. 3.

A.. Dark Phtrotitie magnetic mineral found fn form of float on speep talus slope at contact. Source was no examined because of inaccessability. Hostrock dark fine-grained and of rusty weathering, maybe Argelite.

B. Lenticular bedding Quartz vain inPhyllite well mineralized withdark grey metallic mineral. Magnetite, Phyrrhtite and Chalco Pyrite, Exposure occurs in north wall of creek with an approximate dip of 45° striking north westerly. Visibility of vain in height approximately 25 feet with 2 foot width on top and 4-5 foot on creek bottom. Float of the same mineral is found down to main creek to about 500 feet upstream from showing and for 300 feet southe-east along strike on opposite bank. Enrichment patterns in foloat, visible mineralization in rock yp and down stream are suggesting further mineralized lenses in creek bottom.

C. Mineralized Quartz-lens approximately 10 feet x 15 feet outcrops 1/3 of a mile north of Bon a higher elevation on grassy slope. Minerals identified were Chalco Pyrite, Arseno Pyrite, B. mineralization and a colored tarnish identical to Bournite. Float was traced 45° uphill south east for 200feet and lost in over burden on down ward slopes.

D. Sighting north west and down hill along float zone and outcrop of C. the opposite extension of the mineral zone was found in a little creek near valley bottom. As the mineralization occurs in form of float beneath a hanging wall of Shaly rock no description can be given. The up and downhill mineralization occupies a Shale and Phyllite contact zone with the Shale topping the Phyllites. A light grey Shaly or Phyllite rick separates the two Shale zones as seen on the sketch.

G. Lishy.