PLAN VIEW



PERIDOTITE, PYROXENITE AND DUNITE

GABBRO

MASSIVE SULPHIDE SHOWING

FAULT

1987 DRILL HOLE

LONGITUDINAL SECTION

0 100 500 m

STRATIGRAPHY AND MINERALIZATION

UPPER TRIASSIC
NIKOLAI VOLCANICS
basalt flows, minor limestone, shale.

UPPER TRIASSIC
MAPLE CREEK INTRUSIVES
medium-grained diabasic gabbro dykes and small stocks; probably feeders for Nikolai Group basalts.

LOWER TRIASSIC
WELLGREEN COMPLEX
layered ultramafic sills consisting mainly of dunite, peridotite and gabbro.

LOWER PERMIAN
HASEN CREEK FORMATION
siliceous argillite, sandstone, quartzite, limestone and conglomerate.

PENNSYLVANIAN +? PERMIAN
STATION CREEK FORMATION
tuff, volcanic breccia, limestone, argillite, andesite, and basalt flows.



DUNITE

PERIDOTITE

PYROXENITE

GABBRO

QUARTZITE

CHROMITE

DISSEMINATED SULPHIDES

NET TEXTURED SULPHIDES

MASSIVE SULPHIDES

WELLGREEN PROPERTY

016292

M-590-PR-8