DESCRIPTIVE NOTES

PHYSIOGRAPHY

The map area, a corn mesa by three south-east trending valleys, Sage Creek, East McQuesten River and Lynn Creek. Sage Creek empties Bosemoos Lake and drains valleys and basins in the southern fringe of the Ogilvie Mountains. The highest summits in the map area is Bosemoos Mountain at 6000 feet. Sage Ridge separates Sage Creek from the East McQuesten River. The East McQuesten River occupies northeast of the map area, and until Sage Creek, the name is correct, valley, characteristic confluence with the Hoon McQuesten Rivers to the east. South of the East McQuesten River is a series of plateaus that surround the upper Haggart Creek drainage. These are Haggart Dome, Lynn Dome and Plateau. Lynn Creek valley lies south of Potato Hills. Haggart, Dhiso and Lynn Dome areas are only major north south trending valleys in the map area.

GLACIAL HISTORY

In central Yukon, the Quaternary period is characterized by multiple glaciations and interglaciations. The oldest of Cordilleran ice in Yukon consisted of glaciers originating from accumulation zones in the Ogilvie, Wernecke, Selwyn, Pelly, Cassiar and St. Elias mountains. Ice from these mountains combined and flowed up and into central Yukon plateau region. The Dublin Gulch map area was glaciated by ice originating in the Ogilvie and Wernecke mountains. From plateau to youngest the main glacial episodes are pre-Reid, Reid and McConnell glaciations.

The pre-Reid episode represents multiple unidifferentiated glaciations that occurred approximately between 2.9 million and 200,000 years ago. Landforms from these glaciations have been subsequently eroded and are difficult to identify. In the map area dependence of pre-Reid Dome tillers as a result of accelerated erosion of the Reid glacial limit and to rare deposits exposed in erosion gullies. Reid glacial sediments. Because the area is fragmentary, the pre-Reid limit was not mapped. Pre-Reid drift sheets are found at 4500 feet elevation on Bosemoos Mountain and 4800 feet on Haggart Dome. It possible pre-Reid glacial till is exposed on Potato Hills and in Basin Valley. Such deposits lies at 4800 and 4600 feet elevation. Pre-Reid erratics are also noted up to 4800 feet on Potato Hills. All of the major valleys were glaciated by pre-Reid ice in the map area.

The Reid glaciation occurred at least 200,000 years ago (Bender, 1984) and presumably took somewhat four times as the pre-Reid glaciations, but was less extensive. Reid glacial landforms are moderately preserved, and during the glacial limits can be approximately determined from erratic distribution. Reid Cordilleran ice advanced into the map area from the northeast corner and reached a maximum elevation of approximately 4400 feet at 1. The ice split into two valley lobes, one of which advanced into Sage Creek valley and the other flowed down the East McQuesten River valley. South flowing portion the McQuesten Lake/South McQuesten River lobes crossed a divide into the east. McQuesten River at Lynn Creek and the East McQuesten lobe. Ice from the South McQuesten valley also glaciated the valleys of Lynn, Dhiso and Sheep Creek and flowed into Haggart Creek drainage. Ice reached elevations of at least 5800 feet on the south side of Potato Hills, 5300 feet at Sludge Gulch and 5300 feet at the divide between Lynn Creek and Dhiso Creek, and probably the highest position of limits of ice. Although trace of Dublin Gulch drainage was occupied by the Reid ice, the effect of the ice on the valley was likely more of deposition than erosion, particularly in places that have been glaciated. The same area dominated of Haggart Creek to regional ice flow direction. Limited glacial erosion and resulted in contamination of good placers that were predominantly buried between a glacial till and bedrock. The present valley, between Haggart, Dome, Lynn Dome and Potato Hills which provided a topographic and age barrier as the ice overtopped the head waters of Haggart Creek and Dublin Gulch.

Cordilleran ice from the McConnell glaciation advanced into the map area between 17,000 and 30,000 years ago (Bond, 1997). McConnell landforms and the glacial limits are well recognized in the map area. The McConnell Cordilleran ice, which affected the map area originated in the mountains in the Ogilvie Mountains. Ice advanced down the east McQuesten River valley into the Sage Creek valley and terminated near the mouth of Lynn Creek. East McQuesten River ice also advanced Sage Creek valley, terminated at Bosemoos Lake. Ice of former ice margins/ streams near the terminus of each lobe suggests that the glacier was active. It does. This is supported by numerous moraine ridges, kame terraces and the lack of recessional (retreat) moraines. The humid valley moraine in Sage Creek valley, also moderately clear cut ridges, which led to the erection of Bosemoos and Sage Lakes. The South McQuesten ice lobe entered the map area at the head of Lynn and Lynn creeks. Ice flowed from the Lynn Creek terminating lobe in the mouth of the East McQuesten River valley. South McQuesten ice continued southeast, the Lynn Creek terminated accordingly, 3 km upstream from the confluence of Lynn and Shiko creek.

PLACER POTENTIAL

The physiography and geology of Dhiso Gulch include the access to the fine placer influence on the placer potential of Dublin Gulch map area. The second order is the orientation of drainages with respect to ice flow events. Valleys aligned parallel to regional ice flow are broad and contain thick glacial deposits, whereas the Lynn Creek Dublin Gulch map area is deeply incised in a valley that is transverse to Cordilleran ice flow and cross the Potato Hills glacial lobes. Placer occurs as deposits in the Lynn Creek drainage, which is also poorly mineralized. Potato Hills. The limiting factor for Lynn Creek placers to be a orientation parallel to glacial flow. The valley is oriented to position is valley very close. Our sample of about 50 cubic feet from surface to Lynn Hunting at a good source and access to large scale gold mining. Higher gold concentrations may be found at the base of the valley fill where pre-Reid interglacial and glacial sediments may have concentrated gold and other heavy minerals such as cassiterite.

The main placer areas are Haggart Creek, Dublin Gulch, Hunter Gulch and Dil Gulch. In Haggart Creek, gold has been placer mined from the modern alluvial plain and from terrace deposits and terraces buried under alluvial fans. On Dublin Gulch gold was placer mined from the modern alluvial plain and gulch gravels. Hunter Gulch contains a gulch placer deposit. The placer deposit presently being mined at Dil Gulch is at Haggart Creek alluvial terrace buried beneath the Dil Gulch alluvial fan.

Access to claims may be by means of the tributaries of Lynn Creek. Be-ore small streams drain south from Potato Hills into Lynn Creek, each of these (the largest being Rey Gulch) have a large alluvial fan which is deposited at the placer deposit, during the Holocene and Pleistocene. Gold in the commercial form was mineralization in Potato Hills (Yukon Minfile 1980-37) and modified eastern geomorphic relicts 30 000 gold from each creek. Immediately west of Rey Gulch, about on the ocean bluff, of a gold source and economic for placer gold. The gullies are small, however they are large enough in volume to Dublin Gulch on the opposite side of Potato Hills, which has been successfully mined in the past.

Bluffs Creek is a tributary to Lynn Creek on the south side of the valley, 3 km upstream from Rey Gulch. Gold mineralization is reported from the Lyn Claims immediately south of the creek and the area is probably the economic source for placer gold in Bluffs Creek. The size and of the creek is confined to a deeply downcutting confederate bow Lynn Creek with a minimum elevation. Historic flowing boulder deposits nearby, 1 km upstream from the mouth of Bluffs Creek may be the most prospective part of the drainage.

STRATIGRAPHIC SECTIONS

DUBLIN GULCH
YUKON TERRITORY
SCALE 1:50 000

CONTOUR INTERVAL 100 FEET. Elevations in feet above Mean Sea Level (North American Datum 1927) Transverse Mercator Projection

116 A/8 Worm Lake	106 D/9 Elliot Lake	106 D/8
116 A/1 BOND Geoscience Map 1998-2	106 D/4 Dublin Gulch	106 D/5 McQuesten Lake
116 P/16 BOND Geoscience Map 1998-2	106 M/13 BOND Geoscience Map 1998-3	106 M/14 BOND Geoscience Map 1998-4

QUATERNARY

HOLOCENE

ANTHROPOGENIC DEPOSITS - Human made landforms or structures. Rarely forms a prominent unit. The Haggart and Dublin placer mine tailings form the only mappable unit in the area.

A/T — RUCHIVE TAILINGS - tailings, reworked alluvial and glacial gravel, grit to boulder size, sorting bonds contain rocky, silt and fine sands.

ORGANIC DEPOSITS - peat and woody material, occurring as a flat to gently sloping blanket or as a depression, drains sluggishly, moderately, or poorly; one map glacial lake deposits but rarely form a continent geologic unit. Most common in Eagle Creek valley near Bosemoos and Sage Lakes. Permafrost, a commonly present, within 1 m of surface.

PD — fenland: consisting of woody sedge peat, variable thickness.

ALLUVIAL DEPOSITS - sand and gravel with minor silt and colluvial deposited in modern drainages. Common in Eagle Creek valley, East McQuesten River and Lynn Creek valleys.

A/L (A/L) — A/L - alluvial plain and terraces, coarse sand and gravel with minor silt and fine sand occurring as bars, or broken floodplain and low terrace sediments, 0-3 m thick, floodplain deposits subject to periodic flooding. Small valley alluvial plains may not be mapped at this scale. (A/L) alluvial fan, coarse sand, gravel, cobbles, and rhyolitic deposits, up to 2 m thick, modern depositional fan deposits, often dissected and transformed by colluvial processes during stratigraphic or peripheral sedimentation. Av - complex of A/L and Av/L and Af undivided.

PLEISTOCENE and HOLOCENE (UNDIVIDED)

COLLUVIAL DEPOSITS - diamicton, grey, unsorted bedrock, and lenses of sand and silt derived from bedrock and surficial materials by a variety of physical and chemical weathering processes. Transport of mapped deposits was by surface creep, sheetwash, or by mass - wasting processes. Common on all slopes in the map area.

C - C/s — C - colluvium - veneer that conforms to bedrock topography, <1 m thick.
C/s - C/s - colluvium - apron that conforms to bedrock topography, >1 m thick, colluvial aprons thicken in areas between valley sides and valley floor

AEOLIAN DEPOSITS - fine to medium grained well sorted sand and associated by wind point processes Typically consists of glaciofluvial sands that were remobilized by easterly (glacial) winds.

E/v — E/v - eolian veneer <1 m thick.

LATE PLEISTOCENE (WISCONSINAN) - McCONNELL GLACIATION

ALLUVIAL DEPOSITS - angular grey silt with sand, organic silts, and minor colluvial deposited from streams outside the McConnell limit. Rocks and boulders are transported by the increase in precipitation, frost weathering, and lack of vegetation in the unglaciated terrain during glacial periods.

A/f — A/f - alluvial fan, coarse sand lenses, grey cobbles, and rhyolitic - deposits of lesser marked than those of early stage, deposits, 1-2 m thick. Common bay and the McConnell limit, minor fan - like valley shape a larger valley.

GLACIOFLUVIAL DEPOSITS - sorted to massive, poorly to well sorted, grey silt and sand with minor silt and cobbles, deposited by meltwater originating from glacial ice, locally with a veneer of eolian silt (loess). Common near Sage Lake and in the East McQuesten River valley and the McConnell limit.

G fp/ G fp/ — G fp - glaciofluvial plain, 5-10 m thick; G fp - glaciofluvial terrace, <5-10 m thick; G fp - glaciofluvial plain, includes glaciofluvial deposits and minor colluvial deposits associated with ice contact environments. Common near Sage Lake; G fp - glaciofluvial deposits, includes glaciofluvial deposits associated with lateral meltwater channels. Common near Sage and Bosemoos Lakes.

GLACIAL DEPOSITS (till) - unsorted clay, silt, sand, and gravel with minor cobbles, deposited by or from glacier ice and occurs in a variety of different landforms (plains, hummocks, or ridges) moraine surfaces). Most common in Eagle Creek valley and the east McQuesten River valley.

T/v - T/v - till veneer that conforms to bedrock topography, 0-1 m thick; T/p - till plain, gently to moderately sloping plain controlled by bedrock, >1 m thick; T/h - till complex, hummocky or ridged till associated with ice contact sediments, 3-15 m thick. Also includes some veneer and blanket units undifferentiated.

T/v - T/v

MIDDLE PLEISTOCENE - PRE-McCONNELL GLACIATION (UNDIVIDED)

ALLUVIAL DEPOSITS - stratified sand and gravel with minor silt and cobbles deposited by streams outside the McConnell limit. May include some till or clay of alluvial sedimentation that pre-dates the McConnell glaciation and minor deposition result from McConnell glaciofluvial alluvium. Commonly contain placer gold in favourable host rock settings.

A/f (A/f) — A/f - alluvial plain and terraces, stratified coarse sand and gravel with minor silt and fine sand occurring as bars, or broken floodplain and low terrace sediments, 0-3 m thick, typically above buried deposits under Haggart or McConnell alluvium. A/f - alluvium complex of A/f and/or A/f/l buried under >1 deposits from closely many of these cycles, >30 m thick. Common in the Haggart Creek area.

MIDDLE PLEISTOCENE - REID GLACIATION

ALLUVIAL DEPOSITS - angular grey silt with sand, organic silts, and minor cobbles deposited from streams that are outside the Reid glacial limit. Slope and stream sedimentation is enhanced by the increase in precipitation, frost weathering, and lack of vegetation in the unglaciated terrain during the glacial periods. Hilly, ice dammed which minor deposits of these types possibly during the Reid glaciation.

A/f (A/f) — A/f - alluvial fan, coarse sand lenses, gravel, cobbles, and rhyolitic - deposits, 1-40 m thick, identified as the only deposition from Reid and Boomz Creek to A/f. - A complex of coalescing alluvial fans with alluvial plain sediments, may blanket pre-Reid ice glacial deposits.

GLACIOFLUVIAL DEPOSITS - stratified to massive, poorly to well sorted, grey silt and sand with minor silt and cobbles, deposited by meltwater originating from glacial ice, locally with a veneer of eolian silt (loess). Present on Haggart Creek valley and at the mouth of Omega Creek.

G fp/ G fp/ — G fp - glaciofluvial terrace up to 40 m above modern drainages and - 80m thick; G fp - glaciofluvial complex, includes glaciofluvial deposits and minor deposits associated with ice contact environments such as kame, kame moraine, and glaciolacustrine (lake) sediments. Common along the margins of Haggart Creek.

GLACIAL DEPOSITS (till) - unsorted clay, silt, sand, gravel, and cobbles, with minor boulders, deposited by or from glacier ice and occurs as subdued veneer and blanket deposits. Common in the East McQuesten valley, in Haggart Creek valley, and as a veneer near Sheep Creek. Exposed on the left limits of Dublin Gulch.

T/v - T/v — T/v - till veneer that conforms to underlying topography, <1 m thick; T/p - till plain, gently to moderately sloping plain controlled by bedrock or underlying surficial deposits >1 m thick.

EARLY PLEISTOCENE-PRE-REID GLACIATIONS

GLACIAL DEPOSITS (till) - unsorted clay, silt, sand, gravel, cobbles, and minor boulders (bi-silast), deposited by or from glacier ice and occurs as subdued veneer and blanket deposits. A veneer/blanket deposit is present on the Potato Hills and in Basin Bay Dulch.

T/v — T/v - till veneer that conforms to underlying topography, <1m thick.

PLEISTOCENE UNDIVIDED

COLLUVIAL DEPOSITS - angular rock and cryoplanation terraces. Diamicton and rubble derived from bedrock and surficial materials by a variety of colluvial and mass-wasting processes.

C — C/s - mass wasting includes slumping, debris slides, and rockfalls.

C/s — C/s - rubble and/or diamicton occurring as stepped or flat sloped deposits. Formed through mass-wasting processes on bedrock surface mountain summits. Found at the summit of Lynn Dome

PRE-PLIO-PLEISTOCENE

BEDROCK - dominated by Yukon group rocks of the Selwyn Basin that include greenstone, chlorite, carbonates, and meta-clastic rocks. The Robert Service Thrust, a Late Cretaceous thrust fault has emplaced the Yukon Group over the Paleozoic Hart River Fault. Late Cretaceous to Eocene granitic rocks of the Tombstone and Tintina (1998) and Ketza River area, immediately southeast of Lynn Dome and in the valley south across Sheba Creek (Green, 1972). Cretaceous quartz monzonite and granodiorite with minor granite and quartz diorite intrusions are exposed at Dublin Gulch and in the headwaters of Lynn Creek.

R — R - bedrock, primarily prominent ridges, escarpments, and mountain summits.

The surficial geology units are shown as flat tabs-ed by the terrain hazard modifiers. Combined surficial geology units are used where for reasons of scale, two or more deposits cannot be delineated individually. The dominant unit (>50 percent) is shown first, and the subordinate unit (<50 percent) second, as in till/colluvium and river - flood deposits are the surficial units and each appears in the same relative position from the surficial geology.

The age of glacial deposits are designated by the following subscripts: - McConnell Glaciation, e.g. Gh1 - Reid Glaciation, e.g. Gh1 - Pre-Reid Glaciation, e.g. Gh1

SYMBOLS

Geological boundary

McConnell glacial limit (defined, approximate)

Reid glacial limit (defined, approximate)

Crevasse (McConnell, Reid, Pre-Reid)

Glacial striae (McConnell, Reid, Pre-Reid)

Moraine ridge

Esker

Glacial meltwater channel: McConnell, Reid, Pre-Reid

Angular unconformity

Stream sediment (80 mesh) gold assay (ppb) — J997-38 37

TERRAIN HAZARDS

Mass Movement

- slow to moderate failure (i.e. creep or slump)
- moderate to extreme failure (i.e. slides and avalanches)

Arctic, Alpine, and Periglacial Processes

IP - permafrost within 1 m of surface

K - cryoturbation

S - Solifluction

Ⓐ - Nivation

Fluvial Processes

B - braided, multiple unstable channels, prone to rapid changes (flooding)

A - anastomosing, multiple stable channels, gradual changes in channel/bar morphology

M - meandering, 1-3 channels, gradual channel changes caused by meander migration (lateral)

Marginal Processes

P - pooling, subsurface channel erosion

G - gullying, areas of rapid surface erosion

REFERENCES

BENDER, G.W., 1984 - Age of the Haselton Basalt Creek basins from thermoluminescence dating of bracketing ages at Klaklenka - Bridges of the Science between North America and the Russian Far East, North West Science conference, 29-31 August, 1984 - Anchorage, Alaska and 30 August - 2 September, 1984, Vladivostok, Russia.

BOND, J.D., 1997 - The Glacial History and Placer Gold Potential of the Hoon McQuesten River (116 A/1), Dublin Gulch (106 D/4) and Hanna Hill (106 D/5) Map areas, Mayo Mining District, Central Yukon. In Gabrielse, H.P. and Roots, C.F. (editors), 1997, Yukon Exploration and Geology, Volume II, Exploration and Geological Services Division, Northern Affairs Program, Yukon Region, p.90-98.

GREEN, L.H., 1971 - Geology of Mayo Lake, Scougale Creek and McQuesten Lake map areas, Yukon Territory. Geological Survey of Canada, Memoir 357.

MURPHY, J.B. and HEON, D., 1993 - Geology and tectonic occurrences of Sheba Creek map area (116 P/16), western Selwyn Basin, Yukon. In Yukon and Exploration and Geology 1994 - 95 Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 58-71.

RECOMMENDED CITATION

BOND, J.D., 1997 - Surficial geology of Dublin Gulch, Canada, Yukon, NTS 106 D/4. Exploration and Geological Services Division, Indian and Northern Affairs Canada. Geoscience Map 1998-6 (D). 1:50 000 scale map.

Digital cartography and drafting by Willi ten Hanken, Yukon Geology Program.

Any revisions or additional geological information which may be used should be reported by the Yukon Geology Program.

Copies of this map, the accompanying report and Yukon Minfile may be purchased from Geoscience Information and Sales, Exploration and Geological Services Division, Indian and Northern Affairs Canada, Room 102-300 Main St. Whitehorse, Yukon Y1A 2B5 Ph: 867-667-3266 Fax: 867-667-3267.

Keep the map scale in each area to one and map colours from fading.

This map is released May, 1998, subsequent revision dates are

STRATIGRAPHIC SECTIONS

Indian and Northern Affairs Canada
Exploration and Geological Services Division
Yukon Region

Geoscience Map 1998–6

SURFICIAL GEOLOGY OF DUBLIN GULCH CENTRAL YUKON (106 D/4)

by
J.D. Bond
Yukon Geology Program