

Indian and Northern Affairs Canada | Affaires indiennes et du Nord Canada | Canada | canada/yukon economic development agreement | Yukon Government | Énergie, Mines et Ressources Canada | Energy, Mines and Resources Canada

GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA

LEGEND

NEOGENE

Q — Glacial drift, moraine and river deposits (unconsolidated)

CRETACEOUS

Kr — Rhyolite, trachyte intrusions

Kg — Biotite hornblende granite, quartz monzonite[1]

TRIASSIC

TRJd — Hornblende-clinopyroxene diorite[2] to gabbro; typically foliated meta-diorite

TRIASSIC AND ?JURASSIC

TJs — Recessive, buff to grey weathering, calcareous slate, siltstone and sandstone; dark grey, finely crystalline limestone

MISSISSIPPIAN

"Keno Hill quartzite"[3]

Mq — Dark grey, thick-bedded to massive, vitreous quartzite; minor interbeds of carbonaceous schist; lenses of chloritic phyllite (interpreted as metavolcanic tuff); includes abundant smaller meta-diorite intrusions (TJd)

UPPER DEVONIAN AND MISSISSIPPIAN

(DMe continuous with Earn Group; correlative DMp and DMv (new unit) that includes the former "Lower schist"

DMe — Upper part: Black- and dark brown-weathering, sandstone, chert grit and thickly bedded chert-pebble conglomerate (circle pattern), light gray-weathering fissile mudstone and beige sandstone; Lower part: Black siltstone and mudstone with lenses of chert pebble conglomerate, chert breccia with baritic matrix and dark grey, fetid limestone. At base: Brown-weathering, fine-grained sandstone and siltstone with chert pebbles

DMp — Dark blue-grey- to black-weathering, carbonaceous and locally baritic siliceous slate and chert; local fine-grained, dark grey- to orange-weathering, dark grey quartz arenite; local pebble to cobble conglomerate (circle pattern) with flattened clasts of chert and cobbles of quartz arenite; local chert sandstone associated with dark brown-weathering, medium grey to black slate; local bedded barite

DMv — Dark grey- to orange-weathering, foliated, aphanitic, dark grey to waxy green felsic meta-volcanic rocks and foliated quartz porphyry[2]

CAMBRIAN(?) TO LOWER DEVONIAN(?)

Road River Group

€Dr — Light grey- to blue-weathering, black or light grey chert, interbedded with black mudstone and dark brown and black quartz siltstone

MIDDLE ORDOVICIAN(?)

€O — Olive- and brown-weathering siltstone with black laminae, brown sandstone with thin interbeds of black bioturbated chert

UPPER PROTEROZOIC(?) AND LOWER CAMBRIAN(?)

Hyland Group (P€m, P€h, P€s, P€p; stratigraphic order unknown, except P€m may be youngest)

P€m / P€m' — Maroon and green mudstone and siltstone, locally with white quartz sandstone interbeds; P€m' is foliated and phyllitic equivalent.

P€h — Grey-weathering, fine- to coarse-grained, locally calcareous, quartz meta-sandstone (psammite); pale green to grey phyllite. Minor white to grey, crystalline limestone and phyllitic limestone (hachure pattern)

P€s — Brown-weathering medium- to coarse-grained quartz-mica schist; compositional layering indicated by carbon or iron content, or friability (minor carbonate cement). Some layers contain 5-10% quartz granules. Forms well-stratified members >100m thick within P€h

P€p — Dark brown-weathering, black carbonaceous slate and minor interbedded, fine grained dark grey quartzite

P€v — Chloritic phyllite and meta-diorite

AGE UNKNOWN; POSSIBLY LATE PALEOZOIC

?l — Massive; white weathering, grey, non-crystalline and locally fetid limestone (Van Cleeves Hill, 10 km NW of Mayo Lake)

?q — White-weathering, grey vitreous quartzite, locally graded with siltstone laminae (5 km NW of Kalzas Twins)

Footnotes:

1 K-Ar age determinations from biotite separates (Ma):

Roop Lakes pluton	81.0	Leech et al. 1963
Mt. Haldane porphyry	89.0 ± 2.6	Stevens et al. 1982
Kalzas Twins hornfels	90.1 ± 1.4	Hunt et al. 1987
Two Buttes plug	97.1 ± 3.6	Stevens et al. 1981

2 U-Pb zircon-baddeleyite age from a mafic sill intruding "Keno Hill quartzite" (in Ogilvie Mountains): 232.2 ± 1.5 Ma (Mortensen and Thompson, 1990)

3 Visean (middle Mississippian) conodonts from limestone lenses within "Keno Hill quartzite" in Dempster Highway area (M.J. Orchard, pers. comm., 1987)

4 Similar locally porphyritic felsic metavolcanic rock at MARG (3 km NE of Mayo map area) has been dated by U-Pb on zircon as Late Devonian or Early Mississippian (J. Mortensen, pers. comm., see Turner and Abbott, 1991).

5 Late Middle Devonian to Late Devonian U-Pb zircon age from foliated felsic schist 10 km NE of Mount Roop (J. Mortensen, pers. comm., see Gordey, 1990)

Regional Bedrock Geology maps

Larsen Creek 116A *Green, 1972*	Nash Creek 106D *Green, 1972* 2 5 4 6	Nadaleen R. 106C *Blusson, 1974*
McQuesten 115P *Bostock, 1964*	1 8 3 7 Mayo 105M *Bostock, 1947*	Lansing 105N *Blusson, 1974*
Carmacks 115I *Tempelman-Kluit, 1984*	Glenlyon 105L *Campbell, 1967*	Tay River 105K *Gordey and Irwin, 1987*

Numbered squares indicate 1:50 000 geology maps:

1,2 - Poole, 1965
3,4,5 Green, 1971
6 - Abbott, 1990
7 - Gordey, 1990
8 - Kindle, 1962; Murphy and Roots, 1992

CROSS SECTION NOTES

(These notes refer to circled numbers on the cross section)

1. Dotted form lines indicate orientation of foliation (S1). The foliation is folded by small and large scale folds.
2. Thin, solid form lines schematically indicate folded bedding. Observed folds have axial planes parallel with regional foliation. Inclined short, solid lines at surface of profile are true dips, where measured near line of section.
3. Bottom part of section is not to scale.
4. Robert Service Thrust surface was deformed both during and after motion on the Tombstone thrust. This interpretation is based upon:
   a. Rocks that appear to overlie the "Keno Hill quartzite" vary in composition and degree of deformation (Green, 1971, p. 55). Alternating layers of quartzite (Keno Hill) and schist (Hyland Group) near Sourdough Hill and Mount Hinton could be separated by light folds or splays of the Robert Service Thrust (Roots and Murphy, 1992).
   b. Small lenses of DMp within Hyland Group south of Tiny Island Lake (Gordey, 1990) are interpreted as structural inliers of the footwall of Robert Service Thrust.
   c. Recognition of Keno Hill quartzite, in addition to DMv and DMp (Earn Group) in NW corner of map. This may be an inverted succession of footwall rocks, which implies that the overlying Robert Service fault surface is overturned.
   d. All Hyland Group rocks in the central part of the map area, and as far south as Avalanche Creek, are highly strained but no higher than lower greenschist grade. These rocks appear to have been deformed after they were raised to shallow depth; perhaps by northward thrust motion (Robert Service Thrust) over a ramp which currently underlies the northern limit of unstrained Paleozoic strata.
5. Mayo Lake anticline, recognized by Bostock (1947) and Green (1971), plunges SE at 25°. It probably formed after the Tombstone Thrust, and may be structurally related to the Roop Lakes pluton.
6. Isoclinal folds with both north- and south-vergence and 10-300 m amplitude are visible in outcrops along rapids of Stewart River.
7. Three granitic stocks and fine-grained felsic intrusions form a west-northwest trend across the map south of Stewart River.
8. A prominent break in slope in the Avalanche Creek area is here interpreted as a gently dipping thrust. The hanging wall consists of nearly flat-lying sandstone (cliffs 200 m high) but a topographic bench at their base exposes underlying steeply dipping, strongly deformed phyllite and schist. The contact has not been located. If it is a thrust, large klippen exist in Mayo map area. The contact might, however, be an angular unconformity, with implications for timing of deformation in the Hyland Group.
9. The metamorphic aureole (dark hornfels; may contain cordierite) obscures relations around McArthur pluton. The intrusive contact is vertical where observed in north-facing cirques.
10. These rocks have not been visited by the authors.

REGIONAL SETTING

This new synthesis incorporates the early work of Bostock (1947) in addition to the larger scale mapping in the north quarter of the area (see diagram), combined with new field observations in 1990 and 1991.

Mayo map area contains the eastern ends of two laterally extensive structural sheets that moved northward, first on the Robert Service thrust (Green, 1972) and later on the Tombstone thrust (Roots, 1991; Roots and Murphy, 1992). The southern structural sheet is the hanging wall of the Robert Service Thrust. It includes Hyland Group clastic rocks (Gordey, in press) and middle Paleozoic basinal strata. These units can be traced from the Macmillan Pass area (180 km to the southeast) across Mayo map area and are truncated by the Tintina Fault west of the the McArthur Group mountains. The Hyland Group is increasingly foliated and recrystallized northward (toward Mayo Lake); this may indicate that the Robert Service Thrust is relatively shallow beneath the central part of the map-area.

The northern third of the map-area is the hanging wall of the Tombstone Thrust (Mortensen and Thompson, 1990). It includes the Keno Hill quartzite and older Earn Group schist and phyllite that were mobilized northwest, then northeast (Roots and Murphy, 1992). Abundant Triassic mafic intrusions lie within this structural sheet, and all rocks are highly strained. The Tombstone Thrust may intersect the surface near the north edge of the map-area because rocks to the north (in the footwall) are only weakly deformed (Abbott, 1990).

Several quartz monzonite stocks intrude the structural sheets. In age and composition the stocks are similar to the Selwyn Plutonic Suite (Anderson, 1987), but metallic mineralization is less common. Ag-Pb-Zn vein mineralization in the Keno Hill camp is derived from a mesothermal system driven by the Roop Lakes pluton (Lynch, 1986, 1989) along faults and fractures of a northeast-trending sinistral brittle shear zone (Roots and Murphy, 1992).

SYMBOLS

- Geological contact (defined, approximate, assumed)
- Bedding (tops unknown, known, vertical, overturned)
- Foliation (inclined, vertical)
- Lineation (mineral streaking)
- Antiform or anticline (trace of axial plane; upright, overturned)
- Synform or syncline (trace of axial plane; upright, overturned)
- Fault, displacement unknown (exposed, inferred)
- Thrust fault (approximate, assumed, overturned)
- Mineral occurrence (Yukon Minfile number) — #80

MINERAL OCCURRENCES[1]

1. UNITED KENO HILL (Ag,Pb,Zn veins; Holdings not shown; incl. of Keno and Galena hills)
2. FAITH (Ag,Pb - Vein)
3. DUNCAN (Ag,Pb - Vein)
4. GOLD QUEEN (Ag,Pb - Vein)
5. SILVER BASIN (Pb,Ag,Au - Vein)
6. NABOB (Ag,Pb - Vein)
7. MONUMENT (Ag,Pb - Vein)
8. COMSTOCK (Ag,Pb - Vein)
9. APEX (Ag,Pb,Zn - Vein)
10. VANGUARD (Ag,Pb - Vein)
11. HOMESTAKE (Ag,Pb - Vein)
12. CHRISTINE (Ag,Pb - Vein)
13. MO (Pb,Ag - Vein)
14. MAYBURN (Ag,Pb,Zn - Vein)
15. HOGAN (Ag,Pb - Vein)
16. RUNER (Ag,Pb - Vein)
17. WERNECKE (Ag,Pb,Zn - Vein)
18. FORMO (Ag,Pb,Zn - Vein)
19. NOMAD
20. PADDY (Ag,Pb,Zn - Vein)
21. EAGLE (Ag,Pb,Zn - Vein)
22. FISHER (Pb,Zn,Ag - Vein)
23. PARENT
24. CREAM and JEAN (Pb,Zn,Ag - Vein)
25. NORD (Ag,Pb - Vein)
26. GERLITZKI (Ag,Pb,Zn - Vein)
27. TITAN (Ag,Pb - Vein)
28. SHANGHAI (Ag,Pb,Zn - Vein)
29. WAYNE (Ag,Pb,Zn,Au,W - Vein)
30. ARGENT
31. STREBCHUK (Sn,Ag,Pb,W - Vein)
32. MT. HALDANE (Ag,Pb,Zn - Vein)
33. LAYSER (Ag,Pb - Vein)
34. COBALT (Ag,Pb - Vein)
35. PATTERSON
36. ETTA
37. GORDON (Sb,Ag - Vein)
38. TWO BUTTES (W,Mo - Ppy)
39. SIDESLIP (Cu - Skn)
40. GREAT HORN (W,Cu,Zn - Skn)
41. RAM
42. HOT SPRING (Ag,Pb - Vein)
43. LOST WERNECKE COPPER (Cu - Un)
44. ROOP (W - Skn)
45. ABLE
46. MOON (Ag,Pb - Vein)
47. MT. ALBERT (Ag,Pb - Vein)
48. McKIM (Ag,Pb - Vein)
49. VACA
50. NERO (Ag,Pb - Vein)
51. FRIEBEN (Cu,W - Skn)
52. MT. HINTON (Au,Ag - Vein)
53. AVENUE (Ag,Pb - Vein)
54. CHANCE (Sb - Vein)
55. YONO (Ag,Pb - Vein)
56. SUNDOWN (Ag,Pb - Vein)
57. GUSTAVUS (Ag,Pb - Vein)
58. HALFWAY
59. RANKIN
60. NEWRY
61. CHRISTAL (Ag,Pb,Zn - Vein)
62. SEGSWORTH (Ag,Pb,Zn - Vein)
63. IRONCLAD (Ag,Pb,Zn - Vein)
64. WALLINGHAM
65. NADAR
66. KALZAS (W,Sn - Vein)
67. CORKERY
68. WEASEL
69. GAMBLER (Ag,Pb - Vein)
70. HAVRENAK (Au,Ag - Vein)
71. DRILL (W - Vein)
72. BELEY (Zn - Vein)
73. BEMA (Au,Ag - Vein)
74. BIRGIT
75. WHITEMAN
76. GOLDROCK
77. T-BIRD
78. FEED
79. TINY ISLAND (Ba - Form)
80. GORDEY (Ba - Form)

[1] Numbered with Yukon MINFILE reference numbers. (105M)

REFERENCES

Abbott, J.G.
1990: Geological map of Mt. Westman map area (106D/1), Yukon; Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1990-1.

Anderson, R.G.
1987: An overview of some Mesozoic and Tertiary plutonic suites and their associated mineralization in the northern Canadian Cordillera; in Granite-related mineral deposits: Geology, Petrogenesis and Tectonic Setting; Eds. R.P. Taylor and D.F. Strong; Canadian Institute of Mining and Metallurgy, Geology Division, Special volume 39.

Blusson, S.L.
1974: Geology of Nadaleen River, Lansing, Niddery Lake, Bonnet Plume Lake and Mount Eduni map areas, Yukon Territory; Geological Survey of Canada, Open File 205 (scale 1:250 000).

Bostock, H.S.
1936: Carmacks district, Yukon; Geological Survey of Canada, Memoir 189.

1947: Mayo, Yukon Territory, Geological Survey of Canada, Map 890A.

1964: Geology, McQuesten, Yukon Territory; Geological Survey of Canada, Map 1143A, (scale 1:253 440).

Campbell, R.B.
1967: Geology of Glenlyon map area, Yukon; Geological Survey of Canada, Memoir 352 (includes Map 1221A, scale 1:253 440).

Gordey, S.P.
1990: Geology of Tiny Island Lake map area (105M/16), Yukon; Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1990-2.

in press: Evolution of the northern Cordilleran miogeocline, Nahanni map-area (105I), Yukon Territory and District of Mackenzie; Geological Survey of Canada, Memoir 428.

Gordey, S.P. and Irwin, S.E.B.
1987: Geology, Sheldon Lake and Tay River map areas, Yukon Territory; Geological Survey of Canada, Map 19-1987 (3 sheets), (scale 1: 250 000).

Green, L.H.
1971: Geology of Mayo Lake, Scougale Creek and McQuesten Lake map areas, Yukon Territory (105M/15, 106D/2, 106D/3); Geological Survey of Canada, Memoir 357 (includes map 1270A - Mayo Lake, scale 1:50 000).

1972: Geology of Nash Creek, Larsen Creek and Dawson map areas; Geological Survey of Canada, Memoir 364, (includes maps 1282A - Nash Cr., and 1283A - Larsen Cr., scale 1:250 000).

Hughes, O.L.
1982: Surficial geology and geomorphology, Geological Survey of Canada, Maps 2,3,4,5-1982 (4 maps, scale 1:100 000 for Mayo map area)

Hunt, P.A. and Roddick, J.C.
1987: A compilation of K-Ar ages, Report 17, in Radiogenic and Isotopic Studies, Report 1; Geological Survey of Canada, Paper 1987-2.

Kindle, E.D.
1962: Geology, Keno Hill, Yukon Territory; Geological Survey of Canada, Map 1105A (scale 1:63 360).

Lynch, J.V.G.
1986: Mineral Zoning in the Keno Hill silver-lead-zinc mining district, Yukon; in Yukon Geology, vol. 1, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 89-97.

1989: Large-scale hydrothermal zoning reflected in the tetrahedrite-freibergite solid solution, Keno Hill Ag-Pb-Zn district, Yukon; Canadian Mineralogist, vol. 27, p. 383-400.

Leech, G.B., Lowdon, J.A., Stockwell, C.H., and Wanless, R.K.
1963: Age determinations and geologic studies, including isotopic studies, Report 4; Geological Survey of Canada, Paper 63-17.

Mortensen, J.K. and Thompson, R.I.
1990: A U-Pb zircon-baddeleyite age for a differentiated mafic sill in the Ogilvie Mountains, west-central Yukon Territory; in Radiogenic Age and Isotopic Studies: Report 3, Geological Survey of Canada, Paper 89-2, p. 23-28.

Murphy, D.C. and Roots, C.F.
1992: Geology of Keno Hill map area (105M/14); Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon Region, Open File 1992-3.

Poole, W.H.
1965: Mount Haldane (105M/13) and Dublin Gulch (106D/4) map areas; in Report of Activities, Field, 1964; Geological Survey of Canada, Paper 65-1, p. 32-34.

Roots, C.F.
1991: A new bedrock mapping project near Mayo, Yukon; in Current Research, Part A, Geological Survey of Canada, Paper 91-1A, p. 255-260.

Roots, C.F. and Murphy, D.C.
1992: New developments in the geology of Mayo map area, Yukon Territory; in Current Research, Part A, Geological Survey of Canada, Paper 92-1A, p. 163-171.

Stevens, R.D., DeLabio, R.N., and Lachance, G.R.
1981: Age determinations and geological studies: K-Ar isotopic ages, Report 15; Geological Survey of Canada, Paper 81-2.

1982: Age determinations and geological studies: K-Ar isotopic ages, Report 16; Geological Survey of Canada, Paper 82-2.

Tempelman-Kluit, D.J.
1984: Geology of Laberge and Carmacks map-areas, Yukon Territory; Geological Survey of Canada, Open File 1101 (scale 1:250 000).

Turner, R.J.W. and Abbott, J.G.
1990: Regional setting, structure, and zonation of the MARG volcanogenic massive sulphide deposit, Yukon; in Current Research, Part E, Geological Survey of Canada, Paper 90-1E, p. 31-41.

ACKNOWLEDGEMENTS

This project was funded in 1990 by the Cordilleran Division, Geological Survey of Canada and in 1991 by the Canada-Yukon Economic Development Agreement. Discussions with Grant Abbott of Northern Affairs Program and Steve Gordey of the Geological Survey of Canada were helpful in forming our interpretation of this region. The authors are grateful to Lasha Reid, Archer Cathro and Associates, and Trans North Air, Ltd. for logistics, and Peter Daubeny, Stefan Meinke, Fred Roots and David Lucas for field assistance.

Indian and Northern Affairs Canada
Exploration and Geological Services Division
Yukon Region

Open File 1992-4

GEOLOGY OF MAYO MAP AREA (105M)

by

Charles F. Roots and Donald C. Murphy
Geological Survey of Canada

This Open File map and Open File 1992-3 (Geology of Keno Hill) by D.C. Murphy and C. Roots are products of the Canada-Yukon Cooperation Agreement on Mineral Resources 1991-1996, administered by:
Department of Economic Development,
Government of Yukon
Box 2703 - F3,
Whitehorse, Yukon Y1A 2C6 (403) 667-5384

These Open file maps and other geological publications are available from:
Exploration and Geological Services Division
Northern Affairs Program
200 Range Road
Whitehorse, Yukon Y1A 3V1 (403) 667-3204