










# LANDSCAPE EVOLUTION

**LANDSCAPE EVOLUTION**

Sedimentation into the Old Crow Basin during the Neogene (~23.2 Ma) was dominated by shallow rivers and lacustrine systems. The basin was later occupied by the Laurentide Ice Sheet in the late Pleistocene. Fluvial incision along the Porcupine and Crow rivers and recent basin denudation was likely controlled by combined uplift in the surrounding mountains and relative depression of the Old Crow Basin. The basin was later occupied by the Laurentide Ice Sheet in the late Pleistocene. Fluvial incision along the Porcupine and Crow rivers and recent basin denudation was likely controlled by combined uplift in the surrounding mountains and relative depression of the Old Crow Basin. The basin was later occupied by the Laurentide Ice Sheet in the late Pleistocene. Fluvial incision along the Porcupine and Crow rivers and recent basin denudation was likely controlled by combined uplift in the surrounding mountains and relative depression of the Old Crow Basin.



# TERRAIN CLASSIFICATION SYSTEM

The surficial geology map was classified using the Terrain Classification System for British Columbia (Howes and Kent, 1997), with minor modifications to meet standards set by the Yukon Geological Survey. For example, we have added notes on permafrost processes deposited to accommodate the wider variety of permafrost features found in Yukon. We have also added an age classification to distinguish materials deposited during different Pleistocene glacial advances.

**COMPOSITE SYMBOL DELIMITERS:**

- Due to scale limitations, up to 4 terrain units may be included in a single map unit label (e.g., sgFpOptM-Xs). Each component is separated by a dash (-) to indicate relative proportions between the components ("1" = 1/4, "2" = 1/2, "3" = 3/4, "4" = 100%).
- 1 - terrain units on either side of the symbol are of approximately equal proportion
- "1" - terrain units before the symbol is more extensive than the one(s) following
- "2" - terrain units before the symbol is considerably more extensive than the one(s) following
- "3" - terrain units before the "1" symbol stratigraphically overlies the one(s) following

**TEXTURE**

Texture refers to the size, shape and sorting of particles in clastic sediments, and the proportion and degree of decomposition of plant fibers in organic sediments. Texture is indicated by up to three lower case letters, placed immediately below the surficial material designator, listed in order of decreasing abundance.

**Specific clastic textures**

- b - blocks: angular particles >256 mm in size
- B - boulders: rounded particles >256 mm in size
- c - cobbles: rounded particles 64-256 mm in size
- P - pebbles: rounded particles <64 mm in size
- s - sand: particles >0.0625-2 mm in size
- z - silt: particles 2 µm-0.0625 mm in size
- cl - clay: particles <2 µm in size

**Common clastic textural groupings**

- c - mixed fragments: a mixture of rounded and angular particles >2 mm in size
- a - angular fragments: a mixture of angular fragments >2 mm in size (i.e., a mixture of blocks and rubble)
- g - gravel: a mixture of two or more size ranges of rounded particles >2 mm in size (e.g., a mixture of boulders, cobbles and pebbles); may include interstitial sand
- r - rubble: angular particles between 2 and 256 mm; may include interstitial sand
- m - mud: a mixture of silt and clay; may also contain a minor fraction of fine sand
- sh - shells: a sediment consisting predominantly of shells and/or shell fragments

**Organic terms**

- f - fibric: the least decomposed of all organic materials; contains amounts of well-preserved fibre (40% or more) that can be identified to botanical origin upon rubbing
- h - humic: organic material at a stage of decomposition intermediate between fibric and humic
- u - humic: organic material at an advanced stage of decomposition; it has the lowest amount of fibre, the highest bulk density, and the lowest saturated water-holding capacity of organic materials; fibres that remain after rubbing constitute less than 10% of the volume of the material

**SURFICIAL MATERIAL AGE**

TIME PERIOD	APPROXIMATE AGE
<Hol.	<10,000 years ago
<Pl.	15,000 to <2 million years ago
<P.	>2 million years ago

**SURFICIAL MATERIALS**

Surficial materials are non-lithified, unconsolidated sediments. They are produced by weathering, sediment deposition, biological accumulation, and human and volcanic activity. In general, surficial materials are of relatively young geological age and constitute the parent material of most (pedological) soils. Note that a single polygon will be coloured only by the dominant surficial material, but other materials may exist in that unit.

**Holocene**

- A** - Anthropogenic (A): Surficial materials modified by human activities such that their original physical properties have been significantly altered. Applied to areas within the map containing significant quantities of painted rock on the surface (e.g., sawpiling, logjams, building pads).
- O** - Organic (O): Material derived from decomposition of organic matter and consists of peat with fibric to mesic decomposition. This is the cool part of Yukon organic materials accumulation on top of many surficial deposits including colluvial slopes, ice deposits, and fluvial terraces and floodplains.
- L** - Lacustrine (L): Modern sediments that have settled from suspension in bodies of standing water are limited to thin deposits in small bodies of standing water on the floodplain of the Porcupine River. Sediments consist of stratified fine sand, silt and clay deposited on the lake bed from suspension.
- E** - Eolian (E): Sediment transported and deposited by wind. The dominant eolian material in the map area is loess, which is predominantly silt in texture with a smaller fraction of fine sand. Eolian all deposits are found on both upland and lowland surfaces in the study area, but occur most commonly along the edge of the bluff above terraces (topsoil bays). In cryoblasted and colluvial areas, which are extensive in the map area, loess is reworked into the soil profile and mixed with underlying sediments. Loess is not extensive in the map area, and is indicated by the "L" surficial symbol with other geological material.

**Holocene and Pleistocene**

- C** - Colluvium (C): Material transported and deposited by downslope, gravity-driven processes such as creep, solifluction, landslides and snow avalanches. Colluvium is the dominant surficial material above elevations of ~200 m a.s.l. in the map area. It commonly has a stratified structure with a highly variable texture and composition controlled by the parent material, transport mechanisms, and travel distance. Colluvium on uplands and shallow gradient slopes is generally derived from weathered bedrock, with minor contributions of loess, resulting in a silt-rich matrix containing angular, local bedrock clasts. Colluvium on gentle upland slopes is dominated by slow colluvial processes such as sheetwash, solifluction and creep, and is characterized by greater accumulations of organic permafrost. On steeper slopes on the valley side, colluvium is generally coarse grained, as it has incorporated pre-Pleistocene Fluvial materials and has been deposited by rapid mass wasting processes such as debris falls, slumps, and talus-landslide slides. Colluvial aprons found on the lower slopes of the escarpment and the upland side of the floodplain terrace commonly contain ice-rich permafrost and are primarily composed of reworked glacial materials.

**Pleistocene and Older**

- F** - Fluvial (F): Sediments transported and deposited by modern streams and rivers, found in floodplains, fans and terraces. Fluvial deposits typically consist of well-sorted sand and gravel comprising subangular to rounded clasts. Fluvial fans, fan-shaped landforms at the base of fluvial and colluvial fans and gravel deposits in the Old Crow Basin, are formed by fluvial fans. Fluvial fans are composed of fluvial and colluvial materials. Fluvial fans are composed of fluvial and colluvial materials. Fluvial fans are composed of fluvial and colluvial materials.
- FA** - Active fluvial (FA): Sandy and gravelly materials subject to regular flooding.
- F'** - Fluvial (F'): Sandy pebble and cobble gravel deposited by streams having a fluvial source graded to a former base level of the Porcupine River (possibly >200 m a.s.l.). Pleistocene and older fluvial deposits in the map area range from <10 cm in thickness, and are characterized by well-sorted, pebble-cobble gravel with laterally and vertically discontinuous beds of massive, planar and ripple cross-bedded sand. Gravel units range from cobble to pebble-dominated and can have both open-work and matrix-supported facies. Discontinuous sand and silt units are usually 2-3 cm thick and poorly exposed.
- L'** - Lacustrine (L'): Stratified fine silt, clay and sand deposited on the bed of a lake that existed prior to Pleistocene changes in regional drainage. Neogene and older lacustrine sediments are found in the lowlands that of the Porcupine River in the Old Crow Basin, Northwest Territories. Geological Survey of Canada Bulletin, 458, 45 p.
- D** - Weathered Bedrock (D): The uplands surrounding and making up Berry Hill are uniformly mapped as "weathered bedrock". Weathered bedrock frequently consists of a matrix of blocks and boulders created through frost shattering, colluviation, and chemical weathering processes. Weathered bedrock typically contains a component of loess-derived silt and is subject to sorting and mixing from cryoturbation and other periglacial processes. Permafrost is present in both bedrock and weathered bedrock in the map area.
- R** - Bedrock (R): Bedrock in the map area consists of clastic and carbonaceous sedimentary rocks. The upper Neoproterozoic Katherine Formation in the map area is a massive, very fine grained, thin to thick bedded, brown, greenish grey and white orthoquartzite sandstone with minor recessive intervals of shale. Comprising Berry Hill and the gentle slopes below it, this distinctive white rock is the primary source for crush, rip-up and other aggregate needs for the community of Old Crow. Lying unconformably below this unit is the dark grey siltstone and shale of the Jurassic-aged Porcupine River member of the Husky Formation which outcrops on the west bank of the lower Crow River just above the community of Old Crow.

**LEGEND**

**GROUND OBSERVATION SITES:**

- permafrost field station
- permafrost borehole (labelled with site number)

**GEOLOGICAL FEATURES:**

- cryopreservation terrace
- landslide depositional
- direction of landslide movement
- permafrost

**GEOLOGICAL BOUNDARIES:**

- defined
- approximate
- assumed
- limit of mapping

**TOPOGRAPHIC FEATURES:**

- contours
- streams

**RECOMMENDED CITATION**

Kennedy, K.E. 2016. Surficial geology, Old Crow, Yukon: parts of NTS 1160/12. Yukon Geological Survey, Energy, Mines and Resources, Government of Yukon, Open File 2016-16, 1:10,000 scale.

Any revisions or additional geological information known to the user will be welcomed by the Yukon Geological Survey.

Paper copies of this map may be obtained from Yukon Geological Survey, Room 102 - 300 Main St., Whitehorse, Yukon, Y1A 2B5. Phone: 867-667-2201. E-mail: geology@gov.yk.ca.

A digital PDF (Portable Document Format) file of this map may be downloaded free of charge from the Yukon Geological Survey website: <http://www.geology.gov.yk.ca>.

**SELECTED REFERENCES**

Berkeik, B.E., Kennedy, K., Forster, D., Lewkowicz, A., Roy, L.P., Joy Grandpé, L., Grandmont, K., Drulis, S., Colpron, M., Light, E. and Williams, T. 2016. Old Crow landscape hazard mapping for climate change adaptation planning. Northern Economic Development Agency's (CANMET) Strategic Investments in Northern Economic Development (SINED) program. Vuntut Gwitchin Government and the Vuntut Gwitchin First Nation are gratefully acknowledged for their support and participation in this project.

EBA Terra Tech Company. 2012. Old Crow Landslide Assessment, Impacts and Recommendations. Unpublished report prepared for the Vuntut Gwitchin Government, June, 2012, 16 p.

Howes, D.E. and Kent, E. 1997. Terrain Classification System for British Columbia (Version 2). Recreational Fisheries Branch, Ministry of Environment and Sustainable Resources, Province of British Columbia, Victoria, BC, 45 p.

Hughes, O.L. 1972. Surficial geology of northern Yukon Territory and northwestern District of Mackenzie, Northwest Territories. Paper 69-36, Geological Survey of Canada, Department of Energy, Mines and Resources, 11 p. (1 map sheet).

Hughes, O.L., Plon, J., Vaillette, J.J., Zolai, S.C. and Pettapiece, W.S. 1973. Surficial Geology and Geomorphology of Old Crow (NTS 1160 and 1160-act). Geological Survey of Canada Open File 167, 1 map sheet (unlimited manuscript), scale: 1:25,000.

Kennedy, K.E., Friesse, D.G., Zauza, G.D. and Laurin, B. 2010. Late Glacial Maximum age for the northwest Laurentide maximum from the Eagle River valley and delta complex, northern Yukon. Quaternary Science Reviews, vol. 29, p. 1558-1560.

Kokke, S.V. and Joergensen, M.T. 2013. Advances in Permafrost and Periglacial Processes. vol. 24, p. 108-119.

Lewkowicz, A.G. 2007. Dynamics of active-layer detachment failures. Footham Peninsula, Ellesmere Island, Nunavut, Canada. Permafrost and Periglacial Processes, vol. 18, p. 89-103.

Schwager, C.E. 1989. The Old Crow and Bluefish Basins, Northern Yukon: Development of the Quaternary History. In: Late Cenozoic History of the Interior Basins of Alaska and the Yukon, D.L. Carter, D.L. Hamilton and J.P. Colwell (eds.), Geological Survey of Canada, vol. 1026, p. 30-33.

Thomson, R.M. and Dixon, E.J. 1983. Alluvial history of the Porcupine River, Alaska: Role of glacial lake overflow from Northwest Canada. Geological Society of America Bulletin, vol. 94, no. 5, p. 576-589.

Zauza, G.D., Duk-Robki, A., Schwager, C.E. and Mollen, E.R. 2004. Late Pleistocene chronology of glacial Lake Old Crow and the north-west margin of the Laurentide Ice Sheet. In: Quaternary Glaciations - Extent and Chronology, J. Ehlers and P.L. Gibbard (eds.), Part II, p. 347-362.

Open File 2016-16

**Surficial Geology, Old Crow, Yukon**  
Parts of NTS 1160/12  
1:10 000 scale

by  
K.E. Kennedy

Yukon Geological Survey  
Energy, Mines and Resources  
Government of Yukon

Yukon  
Energy, Mines and Resources